G lassy R atchets For C ollectively Interacting P articles

C.Reichhardt, C.J.Olson Reichhardt, and M.B.Hastings

Center for Nonlinear Studies and Theoretical Physics Division, Los A lam os National Laboratory, Los A lam os, NM 87545

(M arch 22, 2024)

We show that ratchet e ects can occur in a glassy media of interacting particles where there is no quenched substrate. We consider simulations of a disordered binary assembly of colloids in which only one species responds to a drive. We apply an asymmetric ac drive that would produce no net dc drift of an isolated overdam ped particle. When interacting particles are present, the asymmetric ac drive produces a ratchet e ect. A simple model captures many of the results from simulations, including ux reversals as a function of density and temperature.

PACS numbers: 82.70 Dd, 05.40.-a, 05.70 Ln

In ratcheting system s, a dc current or ux arises upon application of a periodic drive, such as when the underlying potential substrate is ashed, or, for a xed substrate, when alternating forces are applied [1]. Ratchet e ects, which can be determ inistic or stochastic in nature, have been studied in a wide variety of system s, including biological motors [2], ux motion in superconductors [3], granular transport [4], and colloidal transport. In many of these system s, an underlying asymmetric potential acts to break the left-right spatial symmetry. A lternative ratchet system s that have recently been investigated [5] have symmetric substrates, and rely on ac drives to provide symmetry breaking. In each of these cases, the particles move through some form of xed substrate.

In this work we study a ratchet e ect that arises due to the collective interactions between driven and non-driven particles in a system that does not contain a xed substrate. The particular system we consider is a binary mixture of interacting repulsive particles which form a disordered or glassy media. These particles, referred to as the passive species, interact only with other particles, and do not respond to an applied ac drive. W e introduce a sm all num ber of a separate species of particles, term ed the active species, which respond both to the other particles and to the ac drive. Using num erical simulations we dem onstrate that a ratchet e ect can be achieved for the active particles in this system when the coupling between the active and passive particles is strong enough and a symmetry breaking is introduced by means of the ac drive. This is in contrast to a recent theoretical proposal of an asymmetric drive ratchet which transports undriven superconducting vortices [6]. In addition, for speci c ac drives, we nd current reversals as a function of density and tem perature. W e also show that the m agnitude of the ratchet e ect goes through a maximum as a function of the number of active particles, due to the collective string-like m otions of the particles.

W e explicitly demonstrate a ratchet e ect form ixtures of colloids. Our system is similar to recent studies of binary colloidalm odels driven with ac elds where one species of the colloids moves in the direction of the applied drive while the other species moves in the opposite direction [7]. These studies have found interesting

collective e ects in which the moving particles organize into lanes. In extensive studies of driven di usive models [8], where the two species of oppositely moving particles are placed on a lattice, a rich variety of nonequilibrium phases were observed. Continuum models, with di erent portions of particles moving in opposite directions, have also been studied in the context of pedestrian ow s [9]. A rem arkable phenom ena term ed \freezing by heating," where the system can jam at higher tem peratures, has recently been observed in these types of system s [10]. Since individual colloids or groups of colloids can be m anipulated easily with optical traps [11], colloidal system s are ideal for experimentally realizing a system in which only a fraction of the particles respond to the applied drive. The active species could be created using charged or magnetic colloids, while the passive particles would be charge-neutral or non-magnetic. For example, in recent experim ents, m agnetic colloids were driven with ac [12] and dc drives [13] through a glassy assembly of other non-magnetic colloids. It should also be possible to drive a fraction of the colloids using optical or magnetic traps.

W e consider a binary two-dimensional (2D) assembly of colloids using the same simulation method as in our previous works [14,15]. The overdam ped equation of motion for colloid i is

$$\frac{\mathrm{d}r_{i}}{\mathrm{d}t} = f_{ij} + f_{AC} + f_{T} ; \qquad (1)$$

 $_{j\in i}^{N}$ r $_{i}V$ (r $_{ij}$) is the interaction force where $f_{ij} =$ from the other colloids, $f_{\!A\,C}\,$ is the force due to an applied driving eld, and f is a thermal force from Langevin kicks. The colloids interact via a Yukawa or screened Coulomb interaction potential, $V(r_{ij}) =$ $(q_i q_j = jr_i)$ r_j) exp(jr_i r_j). Here q_i is the charge on colloid i, 1 = is the screening length, and $r_{i(j)}$ is the position of particle i(j). The system size is measured in units of the lattice constant a, which is set to $a = a_0$ for m ost of this work, giving a particle density = $1=a_0^2$. We take the screening length = $3=a_0.0$ ur system is periodic in both the x and y directions. The total num ber of particles is $N = N_A + N_B + N_D$, with N_A colloids of passive species A and charge q_{A} , and N $_{B}$ colloids of passive species B and charge q_B . W e add N $_D$ active colloids



FIG.1. (a) Velocity v vspower-law exponent from Eq.2 for $f_1 = 1.0$. Ratcheting in the positive direction (v > 0) occurs for > 1. From top left to bottom left, $f_2=f_1 = 1.0$ (at line), 0.9, 0.8, 0.7, 0.6, and 0.5. (b) v vs for xed $f_1=f_2 = 0.5$ and, from bottom to top, $_2=_1 = 3.25$, 2.5, 2, 1.5, 1.25, and 1.

with charge q_D that respond to an applied ac drive. For the results presented here, we $x \in q_B = 1=2$. We have also considered other ratios, including the crystalline case when $q_A = q_B = 1$. To initialize the system, we start at a high temperature and anneal either to T = 0 or to a

xed T > 0. The binary particles of species A and B form a disordered assembly, with the active colloids scattered random ly throughout. A fler annealing, we apply a two-step ac drive during a single cycle of period . The stage is a fast drive where a constant force of f_1 is applied for a time $_1$ in the positive x-direction. The second stage is a slow drive where a force $f_2 < f_1$ is applied for a time $_2 > _1$ in the negative x-direction. The ac drive has the properties $f_{1\ 1} = f_{2\ 2}$, with $= _1 + _2$, so that the particle experiences zero average force during a single cycle. The average drift velocity v_d can be obtained after m any cycles. We conduct a series of simulations for di erent particle densities, system sizes, tem peratures, q_b , ac am plitudes, frequencies, and N_D.

We rst consider a simple model that predicts that a ratchet e ect can arise in such a system. Recently it was shown theoretically and in simulations that, for a single colloid driven with a dc drive through a glassy assem bly of other colloids [14], the colloid velocity follows a power law V / f with = 1:5 for certain parameters. The power law appears in regimes where the driven colloid produces collective disturbances of the surrounding colloids on length scales of the order of a, the lattice constant. W hen the density of the surrounding colloids is low, these collective distortions are lost and the velocity of the driven colloid is linear with the drive. Linear behavior also occurs when the charge q_D of the driven particle is much smaller than that of the surrounding particles. Recent experiments on driven magnetic colloids have also found power law velocity-force relations for the driven colloid with exponents = 1.5 or higher in system s with high density [12]. In low density system s, the velocity is linear with drive.

We nd that for a group of driven particles in a glassy media, the velocity-force curve is also of a power law



FIG.2. Simulation image. Black circles: active particles that respond to the ac drive. Open circles: passive particles form ing the glassy background. Here, $N_D = N = 0.016$.

form . In a system with a power law velocity-force relationship, ratcheting can occur under the application of an asymmetric ac drive. During a single cycle the colloids move a distance $x = f_{1 \ 1} \quad f_{2 \ 2}$. The net drift velocity is then $v_d = x = (1 + 2)$. Using the condition $f_{1 \ 1} = f_{2 \ 2}$ the drift velocity can be expressed as

$$v_{d} = \frac{f_{1} f_{2} f_{2} f_{1}}{f_{2} + f_{1}}$$
(2)

If = 1, as in the case of a single overdam ped particle, then $v_d = 0$, as expected. For any value of > 1, how – ever, the drift velocity will be positive, $v_d > 0$. Ratcheting can still occur if f_1 is close to f_2 as long as > 1. Thism odel predicts that the ratchet e ect become smore e cient for larger values of . In Fig. 1(a) we plot v_d vs

from Eq. (2), with $f_1 = 1$ and ratios of $f_2=f_1 = 0.5$ to 1. For $f_2=f_1 = 1$, $v_d = 0$ for all , as expected. For $f_2 < f_1$ and > 1, ratcheting in the positive direction $(v_d > 0)$ occurs. v_d increases with larger and larger $f_2=f_1$. The velocity reverses for < 1. This is an unphysical limit for the colloid system, where the lowest possible value is = 1. How ever, some physical system s such as non-New tonian uids could have < 1.

Since the velocity becomes linear with the drive at low densities or high drives, it should be possible to select ac drive parameters that reverse the ratchet e ect and give current reversals for > 1. This can be achieved by setting $_1 < _2$, where the inequality is not too large. In the highly nonlinear regime the particle will still ratchet in the positive direction; how ever, upon approaching the linear regime, the particle will ratchet in the negative direction. In Fig.1(b) we show v vs from Eqn. (2) for $f_2=f_1=0.5$ and increasing $_2=_1$, where a crossover from a positive to negative ratchet e ect occurs at $_c$. Here $_c$ increases with increasing $_2=_1$.

W e next consider simulation results. In Fig.2 we show a real space im age of the simulated system with N $_{\rm D}$ = 11



FIG.3. Particle velocity v vs time t= . Dotted line: v for an isolated overdam ped particle under an asym m etric ac drive with $f_1 = 0.56$, $f_2=f_1 = 0.25$, and $_1=_2 = 0.25$. Solid line: v in a system of interacting particles. Inset: D rift velocity v_d vs t= for (dotted line) a single overdam ped particle, and (solid line) the interacting particle system .

active particles (black circles) and N_A + N_B = 678 passive particles (open circles) after m any cycles of the ac drive. The active particles, which were originally scattered throughout the system, have clustered into a lane, and they also drift over time in the positive x-direction.

In Fig. 3 we plot the velocity vs time of a single overdam ped colloid in the absence of any other particles (dashed curve), showing short large positive pulses and long sm all negative pulses of velocity. Here we consider the case of $f_2=f_1 = 0.25$ with $f_1 = 0.56$. The average velocity of the isolated colloid is zero in a single period. The solid line shows the velocity of a driven colloid in the interacting particle system with N = 370 and N $_{\rm D}$ = 11, in a regim e where the dc velocity-force curves have a pow erlaw form . Here, the magnitude of the particle velocity v during both the positive and the negative portions of the ac drive cycle is sm aller than that of the single particle, and shows considerable uctuations. The integrated velocity during the positive portion of the ac cycle is on average larger in magnitude than the integrated velocity during the negative portion of the cycle. To show this more clearly, in the inset of Fig. 3 we plot the tim eaveraged drift velocity v_d for each system over many ac cycles. For the single particle case (low er dashed curve), $v_d = 0$, while for the interacting particle system, $v_d > 0$, with an long time average drift velocity of $\langle v_d \rangle$ 0:03, indicating that a positive ratchet e ect is occurring.

W e next consider the dependence of the ratchet e $\,$ ect on various system parameters. In Fig. 4 (a) we plot the drift velocity v_d vs the fraction of driven particles, N $_D$ =N , for $\,$ xed ac drive parameters f_1 = 0:56, f_2 = f_1 = 0:25, and $_1$ = $_2$ = 0:25, with a= a_0 = 1. v_d goes through a



FIG.4. (a-c) D rift velocity v_d obtained from simulations with $f_{1-1} = f_{2-2}$ and $f_2=f_1 = 0.25$. (a) v_d vs the fraction of driven particles $N_D = N$ for a density of $= 1=a_0^2$. (b) v_d vs particle density for xed $N_D = N = 0.03$. (c) v_d vs temperature $T=T_m$ for $= 1=a_0^2$ and $N_D = N = 0.03$. (d-e) R atchet reversals for a system with $f_{1-1}=f_{2-2} = 0.9$: (d) v_d vs density . (e) v_d vs temperature $T=T_m$ for $= 1.26=a_0^2$.

maximum near $N_D = N = 0.03$, and then slow ly falls o for higher N_D = N before reaching $v_d = 0$ just below N_D = N = 1:0. We have also considered di erent system sizes by varying $a=a_0$. We obtain the same behaviors, indicating that the ratcheting is not a nite size e ect. It is simple to understand why v_d ! 0 in the lim it of N_D = N ! 1, since here all the particles move in unison with the ac drive, and the plasticity responsible for the nonlinear velocityforce response is lost. For a single driven particle N_D = 1, we found a dc velocity-force scaling exponent of = 1:5. As the number of driven particles N_D increases, increases to = $2 \text{ near N}_{D} = N = 0.03$, and then decreases back toward = $1 \text{ as } N_D = N$ increases further. This result is consistent with the predictions of Eq. 2, where a larger ratcheting e ect is expected for larger exponents

In Fig. 4 (b) we consider a system with $x \text{ ed } N_D = N = 0.03$ and ac drive parameters, and plot v_d vs the system density measured in units of $1=a_0^2$. For low densities < 0.5, the active particles interact only weakly with the passive particle background, and the dc velocity-force

curves are close to linear (1) so little ratcheting occurs. For large densities > 1.3, the ratcheting drift velocity v_d decreases since it becomes more di cult for the active particles to pass the passive particles. We have also xed the system density and varied the charge q_b of the active particles. The behavior is similar to Fig. 4 (b): the ratchet e ect is lost at small q_j when the velocity-force curve becomes linear. In Fig. 4 (c) we show the e ect of a nonzero temperature on the ratchet e ect. We plot v_d vs $T=T_m$, where T_m is de ned as the temperature at which a completely ordered single-species colloid assembly with charge q_b and density = $1=a_0^2$ m elts. For high T, the velocity increases linearly with force, so the ratchet e ect is lost.

Equation 2 predicts that a reverse ratchet e ect should occur for < 1, as shown in Fig.1(a), or for $f_{1-1} \notin f_{2-2}$. The reverse ratchet e ect can occur in systems where the media responds easily to slow moving particles but becomes still er under faster perturbations. For example, in granular media the grains respond in a luid manner to a slow moving object, while for a fast moving object the granular material jams and acts like a solid. Many polymer systems show a similar response. In order for a ratchet e ect to occur in a substrate-free system, there must be a combination of the nonlinear behavior of the particle motion and the left-right symmetry breaking by the ac drive. In general the ratchet e ect we observe should occur for any media that has a nonlinear viscosityfrequency response.

In Fig. 4 (d,e) we consider two cases where ux reversals occur with $f_{1}_{1}=f_{2}_{2}=0.9$. For the parameters chosen here, the drift velocity of an isolated particle would be $v_d = 0.016$. In Fig. 4 (d) we plot v_d vs density for $N_D = N = 0.03$. For low density < 0.3, the velocity-force response curve is linear (=1) and the particle m oves in the negative direction as expected for this ac drive. As the density increases, the velocity-force response curve becom es nonlinear, and v_d crosses over to a positive value $v_d > 0$ as the ratcheting e ect reaches a maximum near = $1=a_0^2$. The drift velocity drops at high density just as in Fig. 4(b). In Fig. 4(e) we show V vs $T=T_m$ for the same system with a xed density of $= 126 = a_0^2$. For low tem peratures, the velocity-force response curve is nonlinear, and a ratchet e ect with positive velocity occurs. As the tem perature increases, the drift velocity v_d drops and crosses back to a negative value as the system enters the linear response regime. v_d saturates to v_d 0:01, slightly smaller in magnitude than the value expected for a single isolated particle. We have also considered the e ect of changing q_{1} at xed density and T = 0, and observe behavior sim ilar to that shown in Fig. 4 (d).

In conclusion we have investigated a \glassy ratchet" e ect, which occurs in a glassy system of interacting particles without a quenched substrate. We considered the speci c case of a disordered assembly of colloids with overdam ped dynam ics where only one colloid species couples to an ac drive. The ac drive is asymmetric, with a large positive drive applied for a short time followed by a sm all negative drive applied for a longer time, and obeys the constraint that the net applied force on the driven colloids is zero in a single cycle. A ratchet e ect occurs when the driven and the non-driven colloids are coupled, producing a nonlinear velocity-force response. We nd that the dc drift velocity has a maximum value as a function of the fraction of driven particles, tem perature, and particle density. A simplem odel for the ratchet e ect agrees well with our simulation results, including the occurrence of ratchet reversal regimes. The system considered here can be realized experim entally for m agnetic or charged colloids driven through non-m agnetic or uncharged colloids. Our results may also have applications for new types of electrophoresis devices.

A cknow ledgm ents: W e thank E.W eeks, D G.G rier, and H.Lowen for useful discussions. This work was supported by the USD epartm ent of Energy under Contract No.W -7405-ENG -36.

- For reviews, see P.Reim ann, Phys.Rep 361, 57 (2002);
 R D.Astum ian and P.Hanggi, Physics Today, 55, 33 (2002).
- [2] R D. A stum ian and M. Bier, Phys. Rev. Lett. 72, 1766 (1994).
- [3] C.-S. Lee et al., Nature 400, 337 (1999).
- [4] Z. Farkas, P. Tegzes, A. Vukics, and T. Vicsek, Phys. Rev. E 60, 7022 (1999); JF. W am baugh, C. Reichhardt, and C.J. O lson, ibid. 65, 031308 (2002).
- [5] O. Usm ani, E. Lutz, and M. Buttiker, Phys. Rev. E 66, 021111 (2002); C. Reichhardt, C.J. Olson, and M.B. Hastings, Phys. Rev. Lett. 89, 024101 (2002); R.Guantes and S.M iret-Artes, Phys. Rev. E 67, 046212 (2003).
- [6] S. Savel'ev and F. Nori, Nature Mater. 1, 179 (2002).
- J.D zubiella and H.Lowen, J.Phys.Condens.M atter 14, 9383 (2002); J.D zubiella, G P.Ho m ann, and H.Lowen, Phys.Rev.E 65, 021402 (2002); J.Chakrabarti, J.D zubiella, and H.Lowen, Europhys. Lett, 61, 415 (2003).
- [8] B. Schm ittm ann, K. Hwang, and R.K.P. Zia, Europhys. Lett. 19, 19 (1992); I. Vilfan, R.K.P. Zia, and B. Schm ittm ann, Phys. Rev. Lett. 73, 2071 (1994);
 B. Schm ittm ann and R.K.P.Zia, Statistical Mechanics of Driven Di usive Systems, (New York, Academic Press, 1995)
- [9] D. Helbing, I. Farkas, and T. Vicsek, Nature 407, 487 (2000); D. Helbing, Rev. M od. Phys. 73, 1067 (2001).
- [10] D. Helbing, I.J. Farkas and T. Vicsek, Phys. Rev. Lett. 84, 1240 (2000); H E. Stanley, Nature 404, 718 (2000).
- [11] D.G.Grier, Nature, in press.
- [12] R E. Courtland, P. Havdas, and E R. W eeks, to be published.
- [13] S.W u et al., to be published.
- [14] M B. Hastings, C J. O lson Reichhardt, and C. Reichhardt, Phys. Rev. Lett. 90, 098302 (2003).
- [15] C. Reichhardt and C.J. Olson, Phys. Rev. Lett. 88, 248301 (2002).