

## Bogoliubov sound speed in periodically modulated Bose-Einstein condensates

E. Taylor and E. Zaremba

Department of Physics, Queen's University, Kingston, Ontario K7L 3N6, Canada.

(Dated: March 22, 2024)

We study the Bogoliubov excitations of a Bose-condensed gas in an optical lattice. Of primary interest is the long wavelength phonon dispersion for both current-free and current-carrying condensates. We obtain the dispersion relation by carrying out a systematic expansion of the Bogoliubov equations in powers of the phonon wave vector. Our result for the current-carrying case agrees with the one recently obtained by means of a hydrodynamic theory.

PACS numbers: 03.75.Kk, 03.75.Lm, 05.30.Jp, 67.40.Db

## I. INTRODUCTION

The possibility of creating optical lattices in trapped Bose-condensed gases has provided an opportunity to study superfluids in novel situations. The presence of the lattice leads to a variety of solid state effects associated with the coherent motion of the atoms in a periodic potential. For example, the oscillation frequency of the centre of mass motion of the condensate is reduced [1] as a result of the enhanced effective mass of the atoms tunnelling between potential wells. Furthermore, when subjected to a uniform force, as provided by gravity [2] or alternatively by accelerating the optical lattice itself [3], Bloch oscillations of the condensate have been observed. Reducing the amplitude of the lattice potential leads to a breakdown of these oscillations as a result of Landau-Zener tunnelling between bands [3]. All of these observations are essentially a manifestation of the superfluidity of the Bose condensate in an optical lattice. Another aspect of equal interest is the breakdown of superfluidity as recently observed [4] in a study of the centre of mass motion of a trapped condensate moving through an optical lattice. When the amplitude of the oscillation exceeded a critical value, dissipation was seen to set in.

In this paper we study a Bose-condensed gas subjected to a uniform optical lattice in a regime where the dynamics of the condensate wave function is well described by the time-dependent Gross-Pitaevskii (GP) equation. In particular, we are concerned with small amplitude collective modes which at long wavelength are phonon-like excitations. The relevant physical parameters determining the properties of the excitation are the optical potential amplitude,  $V_0$ , the lattice constant,  $d$ , the mean density,  $n$ , of the gas, and the magnitude of the supercurrent. The problem has been addressed theoretically in a number of papers using a variety of techniques and approximations.

Berg-Sørensen and Mølmer [5] were the first to investigate phonon excitations within an optical lattice. They solved the Bogoliubov equations numerically for a one-dimensional model and established that the long wavelength excitations are phonon-like, having an energy dispersion that is linear in the wave vector of the mode. They also obtained an analytic expression for the sound

speed,  $s$ , which is based on a combined weak potential and slowly varying approximation. These calculations were extended by Wu and Niu [6] to the case in which the condensate carries a current. This work is noteworthy for having pointed out that the modes exhibit both energetic and dynamic instabilities for sufficiently large currents. The former instability is associated with the Landau criterion for the breakdown of superfluidity, while the latter is related to the onset of dissipation as observed in [4]. The recent paper by Machholm et al. [7] explores these instabilities further. Similar results were obtained by Bronski et al. [8] by considering a special form of the lattice potential, while Konotop and Salamo [9] used a different approach to establish that the dynamic (or modulational) instability leads to the generation of solitons.

When the potential wells are sufficiently deep, the condensate is well-localized on each lattice site and a tight-binding description becomes useful. Javanainen [10] used this picture within a many-body formulation to derive the phonon dispersion throughout the Brillouin zone for a one-dimensional lattice. This calculation is in fact equivalent to one based on the Bogoliubov equations [11] or the discrete version of the time-dependent GP equation [12]. The virtue of these methods is that they provide analytical expressions for the dispersion relation, although an accurate a priori determination of the tight-binding parameters involves further numerical calculation. Smorzi et al. [12] extended the results of Javanainen [10] by deriving the phonon dispersion for a current-carrying state and found a dynamic instability that is responsible for a so-called 'superfluid-insulator' transition.

The phonon dispersion at long wavelengths was addressed in [13] by deriving an energy functional involving density and phase fluctuations which vary slowly in space. The approach is closely allied to the effective mass approximation used in solid state physics [14] and recently applied to Bose gases in optical lattices [15], and to multiple-scale analysis [9, 16]. The phonon sound speed is found to be [17]

$$s = \frac{\hbar}{m} \frac{\partial \mu}{\partial n}$$

where  $m$  is an effective mass and  $\mu$  is the chemical potential. The precise definition of the effective mass as

$(m)^{-1} = \sim^2 \partial^2 \sim = \partial^2 K^2$ , where  $\sim$  is the energy per particle of the  $N$ -particle system, seems first to have appeared later [7, 18]. This is an important point since there are other plausible candidates for the effective mass. In fact, we shall see that a somewhat different, but equivalent, definition can be given. In this regard, we note that the effective mass appearing in both the effective mass [15] and multiple-scale [9] theories is that corresponding to the bare optical potential. In other words, the effect of the interactions on this parameter is not included, and therefore the use of the dynamical equations obtained in these theories will not in general give the correct Bogoliubov sound speed.

Our purpose in this paper is to obtain the long wavelength phonon dispersion directly from the Bogoliubov equations defining the collective modes. This is achieved by developing a systematic expansion of these equations in powers of the phonon wave vector  $q$ . We do this first for the current-free state (Sec. III), confirming the result for the sound speed given above. We then consider a current-carrying state (Sec. IV) and obtain the analogous phonon dispersion in this case, reproducing the result obtained in [7] by means of a hydrodynamic analysis. Our expansion technique can be viewed as a justification of the assumptions on which the hydrodynamic approach is based. Furthermore, it provides explicit perturbative expressions for the various physical quantities that appear in the theory (for example, the effective mass).

In Sec. II we present the theoretical background required for the calculation of small amplitude collective excitations in an optical lattice. For the most part we consider a three dimensional optical potential with cubic symmetry, although we also touch on systems with one dimensional modulation as well as radially confined condensates. The underlying periodicity of the optical potential implies that the Bogoliubov equations admit solutions having a Bloch function form. This aspect accounts for the use of a Bloch function basis in solving these equations in both the current-free (Sec. III) and current-carrying (Sec. IV) states. However, different calculational methods are used in the two cases and these are therefore presented separately. We also examine various physical limits (Thomas-Fermi, weak potential, weak coupling and tight-binding) in order to make contact with previous work. As stated previously, our main result for the phonon dispersion aims the result which follows from the insightful use of hydrodynamic equations to describe the dynamics of long wavelength fluctuations [7, 13].

## II. BASIC THEORY

We consider an extended 3D BEC subjected to standing wave light fields that give rise to a periodic external potential having the property  $V_{\text{opt}}(\mathbf{r} + \mathbf{R}) = V_{\text{opt}}(\mathbf{r})$ , where  $\mathbf{R}$  is a Bravais lattice vector. For the most part we restrict ourselves to a cubic lattice for which

$\mathbf{R} = d(n_1\hat{x} + n_2\hat{y} + n_3\hat{z})$ , with  $n_i$  an integer.

We base our analysis on the time-dependent Gross-Pitaevskii (GP) equation for the condensate wave function,  $\psi(\mathbf{r};t)$ ,

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r};t) = \frac{\hbar^2 \nabla^2}{2m} \psi + V_{\text{opt}}(\mathbf{r}) \psi + g |\psi|^2 \psi \quad (\mathbf{r};t): \quad (1)$$

This equation admits stationary solutions of the form

$$\psi(\mathbf{r};t) = \psi(\mathbf{r}) e^{-iEt/\hbar}$$

where  $\psi(\mathbf{r})$  satisfies the time-independent GP equation

$$\frac{\hbar^2 \nabla^2}{2m} \psi + V_{\text{opt}} \psi + g |\psi|^2 \psi = E \psi \quad (2)$$

with the normalization

$$\int_V d^3r |\psi(\mathbf{r})|^2 = N;$$

where  $N$  is the total number of particles in the volume  $V$ . Also of interest is the total energy of the system given by

$$E_{\text{tot}} = \int_V d^3r \left[ \frac{\hbar^2 \nabla^2}{2m} \psi + V_{\text{opt}} \psi + \frac{g}{2} |\psi|^4 \right] \quad (3)$$

The energy parameter  $E$  is the chemical potential and is related to  $E_{\text{tot}}$  by  $E = \partial E_{\text{tot}} / \partial N$ .

Often the ground state solution of the GP equation is of interest but we shall also consider states which have a superfluid flow. These states have a Bloch function form

$$\psi_{\mathbf{n}\mathbf{k}}(\mathbf{r}) = \frac{1}{N} \sum_{\mathbf{n}} e^{i\mathbf{k} \cdot \mathbf{r}} \tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r})$$

where  $\mathbf{n}$  is a band index and  $\mathbf{k}$  is a wave vector restricted to the first Brillouin zone. The factor  $\frac{1}{N}$ , where  $n$  is the mean density, is introduced in the definition of  $\tilde{w}_{\mathbf{n}\mathbf{k}}$  so as to give the normalization

$$\frac{1}{V} \int_V d^3r |\tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r})|^2 = 1; \quad (4)$$

where  $V$  is the Wigner-Seitz volume. The condensate density is then  $n_c(\mathbf{r}) = |\psi(\mathbf{r})|^2 = |\sum_{\mathbf{n}\mathbf{k}} \tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r})|^2$ . The Bloch function  $\tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r})$  is in general complex and is the self-consistent solution of

$$\frac{\hbar^2 (\nabla + i\mathbf{k})^2}{2m} \tilde{w}_{\mathbf{n}\mathbf{k}} + V_{\text{opt}} \tilde{w}_{\mathbf{n}\mathbf{k}} + g |\sum_{\mathbf{n}\mathbf{k}} \tilde{w}_{\mathbf{n}\mathbf{k}}|^2 \tilde{w}_{\mathbf{n}\mathbf{k}} = E \tilde{w}_{\mathbf{n}\mathbf{k}} \quad (5)$$

We assume that  $\tilde{w}_{\mathbf{n}\mathbf{k}}$  has the periodicity of the lattice,  $\tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r} + \mathbf{R}) = \tilde{w}_{\mathbf{n}\mathbf{k}}(\mathbf{r})$ , although it should be noted that period-doubled states also exist [19]. The chemical potential,  $\mu_{\mathbf{n}\mathbf{k}}(n)$ , is implicitly a function of the mean density and depends on the particular Bloch state being considered.

The super uid current density in this state is

$$\begin{aligned} \mathbf{j}_s(\mathbf{r}) &= \frac{\hbar}{2m} \nabla \cdot (\mathbf{r} \cdot \nabla \psi_{nk}) \\ &= \frac{\hbar}{m} \nabla \cdot (\mathbf{r} \cdot \nabla \psi_{nk}) + \frac{\hbar}{2m} \nabla \cdot (\mathbf{r} \cdot \nabla \psi_{nk}) \end{aligned}$$

and has the property  $\mathbf{r} \cdot \nabla \psi_{nk} = 0$ . Introducing the super uid velocity according to the relation

$$\mathbf{j}_s(\mathbf{r}) = n_s(\mathbf{r}) \mathbf{v}_s(\mathbf{r}); \quad (6)$$

we have

$$\mathbf{v}_s(\mathbf{r}) = \frac{\hbar}{m} (\mathbf{k} + \nabla \phi_{nk}(\mathbf{r})) \quad (7)$$

where  $\phi_{nk}(\mathbf{r})$  is the phase of the Bloch function  $w_{nk}$ . The spatially-averaged super uid velocity is

$$\langle \mathbf{v}_s \rangle = \frac{1}{V} \int d^3r \mathbf{v}_s(\mathbf{r}) = \frac{\hbar}{m} (\mathbf{k} + \langle \nabla \phi_{nk} \rangle); \quad (8)$$

In one dimension, the periodicity of  $w_{nk}(\mathbf{x})$  implies  $\langle \nabla \phi_{nk} \rangle = 2\pi l/d$ , where  $l$  is an integer. By continuity of the phase with  $k$ , we expect  $l$  to have a fixed value for a given band, and  $\langle \mathbf{v}_s \rangle = (\hbar/m)(\mathbf{k} + \mathbf{G})$  where  $\mathbf{G}$  is some reciprocal lattice vector. For the lowest band, we show in Appendix A that  $\mathbf{G} = 0$ . Thus, we arrive at the somewhat surprising conclusion that  $\langle \mathbf{v}_s \rangle = \hbar \mathbf{k}/m$ . We suspect that similar results apply in higher dimensions but have not been able to show this explicitly.

The average super uid velocity should be distinguished from the velocity determining the average current density

$$\begin{aligned} \langle \mathbf{j}_s \rangle &= \frac{1}{V} \int d^3r \nabla \cdot (\mathbf{r} \cdot \nabla \psi_{nk}) \\ &= \frac{n}{m} \int d^3r \nabla \cdot (\mathbf{r} \cdot \nabla \psi_{nk}) \\ &= n \mathbf{v}_{nk}; \end{aligned} \quad (9)$$

This velocity is given by [20]

$$\mathbf{v}_{nk} = \frac{1}{\hbar} \nabla \epsilon_{nk}(\mathbf{k}); \quad (10)$$

where  $\epsilon_{nk} = E_{\text{tot}}/N$  is the energy per particle in the state characterized by the mean density  $n$  and quasimomentum  $\mathbf{k}$ .

In one dimension, the average current density vanishes at the zone boundary  $k = \pi/d$  if the mean density is below a critical value  $n_c$  [20]. On the other hand, the average super uid velocity is  $\langle \mathbf{v}_s \rangle = \hbar \mathbf{k}/m$ . This is not a contradiction since the local super uid velocity in (7) is averaged differently when calculating the average current density.

Dynamical states of the condensate are determined by the time-dependent GP equation (1). For small amplitude excitations, the condensate wave function is expressed as

$$\psi(\mathbf{r};t) = [\psi_{nk}(\mathbf{r}) + \psi_i(\mathbf{r};t)] e^{i\mathbf{k} \cdot \mathbf{r} - i\epsilon_{nk}t} \quad (11)$$

and the GP equation is expanded to first order in the deviation  $\psi_i(\mathbf{r};t)$ . By writing

$$\psi_i(\mathbf{r};t) = u_i(\mathbf{r}) e^{iE_i t} + v_i(\mathbf{r}) e^{-iE_i t}; \quad (12)$$

where  $E_i$  is allowed to be complex, one obtains the following Bogoliubov equations for the quasiparticle amplitudes  $u_i$  and  $v_i$ ,

$$\begin{aligned} \hat{L} u_i(\mathbf{r}) - g_{nk}^2(\mathbf{r}) v_i(\mathbf{r}) &= E_i u_i(\mathbf{r}) \\ \hat{L} v_i(\mathbf{r}) - g_{nk}^2(\mathbf{r}) u_i(\mathbf{r}) &= -E_i^* v_i(\mathbf{r}) \end{aligned} \quad (13)$$

where the operator  $\hat{L}$  is defined as

$$\hat{L} = \frac{\hbar^2 \nabla^2}{2m} + V_{\text{opt}} + 2g \psi_{nk}^2 \quad (14)$$

Each distinct solution labelled by the index  $i$  corresponds to a collective excitation of the condensate and  $E_i$  represents the excitation energy of the mode. The orthonormality of the quasiparticle amplitudes is specified by

$$\int d^3r [u_i u_j - v_i v_j] = \delta_{ij}; \quad (15)$$

Since the operator  $\hat{L}$  has the translational symmetry of the lattice, the Bogoliubov equations admit solutions of the form

$$\begin{aligned} u_i(\mathbf{r}) &= e^{i(\mathbf{q} + \mathbf{k}) \cdot \mathbf{r}} \tilde{u}_i(\mathbf{r}) \\ v_i(\mathbf{r}) &= e^{i(\mathbf{q} - \mathbf{k}) \cdot \mathbf{r}} \tilde{v}_i(\mathbf{r}) \end{aligned}$$

where  $u_i(\mathbf{r})$  and  $v_i(\mathbf{r})$  have the periodicity of the lattice. These functions satisfy

$$\begin{aligned} \hat{L}_{q+k} u_i(\mathbf{r}) - g n w_{nk}^2(\mathbf{r}) v_i(\mathbf{r}) &= E_i u_i(\mathbf{r}) \\ \hat{L}_{q-k} v_i(\mathbf{r}) - g n w_{nk}^2(\mathbf{r}) u_i(\mathbf{r}) &= -E_i^* v_i(\mathbf{r}) \end{aligned} \quad (16)$$

with

$$\hat{L}_{q+k} = \frac{\hbar^2}{2m} (\mathbf{r} + i\mathbf{q} + i\mathbf{k})^2 + V_{\text{opt}} + 2g n \psi_{nk}^2 \quad (17)$$

Our notation emphasizes that  $\mathbf{q}$  and  $\mathbf{k}$  play distinct roles in the Bogoliubov equations: the former characterizes the Bloch-like character of the quasiparticle amplitudes while the latter corresponds to the quasimomentum of the condensate wave function.

In the following, we shall also make use of the Hamiltonian

$$\hat{H}_k(\mathbf{q}) = \frac{\hbar^2}{2m} (\mathbf{r} + i\mathbf{q} + i\mathbf{k})^2 + V_{\text{opt}} + g n \psi_{nk}^2 \quad (18)$$

which for  $\mathbf{q} = 0$  is just the Hamiltonian determining the time-independent condensate wave function  $w_{nk}$ .

### III. PHONON DISPERSION FOR A STATIONARY CONDENSATE

We begin by considering the simpler situation in which there is no superfluid flow ( $\mathbf{k} = 0$ ). In this case, the ground state solution of the time-independent GP equation can be taken to be real, and  $n w_{n=0, \mathbf{k}=0}^2(\mathbf{r}) = n w_{n=0, \mathbf{k}=0}^2(\mathbf{r}) = n_c(\mathbf{r})$ . It is then convenient to introduce the functions  $\psi_i = u_i - v_i$  and to combine the Bogoliubov equations into a single equation for  $\psi_i$  [21],

$$\hat{h}_0^2 \psi_i + 2gn_c \hat{h}_0 \psi_i = E_i^2 \psi_i \quad (19)$$

where  $\hat{h}_0$  is the Hamiltonian

$$\hat{h}_0(\mathbf{q}) = \frac{\hbar^2}{2m} (\mathbf{r} + i\mathbf{q})^2 + V_{\text{opt}} + gn_j w_{0,0}^2 \quad (20)$$

in which the mean field,  $gn_j w_{0,0}^2$ , of the current-free condensate appears.

To solve (19), we introduce a complete set of Bloch states which are solutions of the equation

$$\hat{h}_0(\mathbf{q}) w_{n\mathbf{q}} = E_n(\mathbf{q}) w_{n\mathbf{q}} \quad (21)$$

This is a linear Schrödinger equation but the solution for  $\mathbf{q} = 0$  and  $n = 0$  coincides with the self-consistent GP solution  $w_{0,0}$ . By definition, the band energies,  $E_n(\mathbf{q})$ , are referred to  $w_{0,0}$ , so that  $E_0(0) = 0$ . The functions  $w_{n\mathbf{q}}$  satisfy the periodicity property  $w_{n\mathbf{q}}(\mathbf{r} + \mathbf{R}) = w_{n\mathbf{q}}(\mathbf{r})$  and the orthonormality relation

$$\frac{1}{V} \int d^3r w_{n\mathbf{q}} w_{n'\mathbf{q}} = \delta_{nn'} \quad (22)$$

In addition, at  $\mathbf{q} = 0$  they are chosen to be real.

Since  $\psi_i$  is itself a Bloch function, it can be expanded as

$$\psi_i(\mathbf{r}) = \sum_n c_n(\mathbf{q}) w_{n\mathbf{q}}(\mathbf{r}) \quad (23)$$

The label  $i$  represents a band index and the Bloch wave vector  $\mathbf{q}$ . However in the following, we will only be interested in the lowest excitation band and will therefore drop the label for convenience. Substituting this expansion into (19) we obtain

$$\sum_{n^0} M_{nn^0}(\mathbf{q}) E_{n^0}(\mathbf{q}) c_{n^0}(\mathbf{q}) = E^2(\mathbf{q}) c_n(\mathbf{q}) \quad (24)$$

where

$$M_{nn^0}(\mathbf{q}) = \frac{1}{V} \int d^3r w_{n\mathbf{q}}(\mathbf{r}) 2gn_c(\mathbf{r}) w_{n^0\mathbf{q}}(\mathbf{r}) + E_n(\mathbf{q}) \delta_{nn^0} \quad (25)$$

We have displayed explicitly the  $\mathbf{q}$ -dependence of all the variables.

For a cubic lattice, we anticipate a particular eigenvalue  $E(\mathbf{q})$  which has a linear dispersion of the form

$$E(\mathbf{q}) = \hbar s q +$$

Our objective is to derive an explicit expression for the Bogoliubov sound speed  $s$ . From Eq. (19) it is clear that in the  $q \rightarrow 0$  limit, the eigenvector corresponding to this particular eigenvalue will be

$$c_n(0) = \delta_{n0} \quad (27)$$

where  $n = 0$  labels the lowest Bloch band, since only this band has a vanishing energy ( $E_0(0) = 0$ ). As a function of  $\mathbf{q}$ , the lowest band energy behaves as

$$E_0(\mathbf{q}) = \frac{\hbar^2 q^2}{2m_0} + O(q^4) \quad (28)$$

which defines the effective mass  $m_0$  of this band. We emphasize that this band mass is determined by the linear Schrödinger equation (21). More will be said about this later. The phonon eigenvector is a continuous function of  $\mathbf{q}$  and, as we shall see, behaves as  $c_n(\mathbf{q}) = \delta_{n0} + O(q^2)$  for small  $\mathbf{q}$ .

To obtain an expression for  $s$  we separate the  $n = 0$  equation in (24)

$$E^2 c_0(\mathbf{q}) = \sum_{n^0 \neq 0} M_{00n^0}(\mathbf{q}) E_{n^0}(\mathbf{q}) c_{n^0}(\mathbf{q}) + \sum_{n^0 \neq 0} M_{0n^0}(\mathbf{q}) E_{n^0}(\mathbf{q}) c_{n^0}(\mathbf{q})$$

from the  $n \neq 0$  equations

$$E^2 c_n(\mathbf{q}) = \sum_{n^0} M_{nn^0}(\mathbf{q}) E_{n^0}(\mathbf{q}) c_{n^0}(\mathbf{q}) + \sum_{n^0 \neq 0} M_{nn^0}(\mathbf{q}) E_{n^0}(\mathbf{q}) c_{n^0}(\mathbf{q})$$

Since  $E_0(\mathbf{q})$  and  $E^2$  are both proportional to  $q^2$ , the latter equation shows that  $c_n(\mathbf{q}) / q^2$  for  $n \neq 0$ , as claimed. Thus, to order  $q^2$ , these equations can be replaced by

$$E^2 c_0(0) = \sum_{n^0} M_{00n^0}(0) E_{n^0}(0) c_{n^0}(0) + \sum_{n^0 \neq 0} M_{0n^0}(0) E_{n^0}(0) c_{n^0}(0) \\ 0 = \sum_{n^0} M_{n0}(0) E_{n^0}(0) c_{n^0}(0) + \sum_{n^0 \neq 0} M_{nn^0}(0) E_{n^0}(0) c_{n^0}(0) \quad (29)$$

Solving for  $E^2$ , we obtain

$$E^2 = \frac{\hbar^2}{2m_0} \left( \sum_{n^0} M_{00n^0}(0) E_{n^0}(0) c_{n^0}(0) + \sum_{n^0 \neq 0} M_{0n^0}(0) E_{n^0}(0) c_{n^0}(0) \right) \quad (30)$$

where all quantities within the square brackets are understood to be the  $q = 0$  values. The prime on the summation indicates that the terms  $n = 0$  and  $n^0 = 0$  are excluded from the sum. The matrix  $\tilde{M}_{nn^0}$  is the matrix obtained by deleting the first row and first column of  $M_{nn^0}$ . We note that this combination of matrix elements can in fact be written as

$$M_{00} \sum_{n^0} M_{0n^0}(\tilde{M}^{-1})_{nn^0} M_{n^0 0} = \frac{1}{(\tilde{M}^{-1})_{00}} \quad (31)$$

Thus we find that the square of the sound speed is given by

$$s^2 = \frac{1}{2m_0(M^{-1})_{00}} : \quad (32)$$

We next relate the sound speed to variations of the chemical potential with mean density. Writing for simplicity  $w_0 = w_{0,0}$  and  $\phi_0 = \phi_{0,0}$ , we have

$$\frac{\hbar^2}{2m} r^2 w_0 + V_{\text{opt}} w_0 + gnw_0^3 = \phi_0 w_0 : \quad (33)$$

The derivative of this equation with respect to  $n$  is

$$\hat{h}_0(0)w_{0,n} + gw_0^3 + 2gnw_0^2w_{0,n} = \phi_{0,n}w_0 \quad (34)$$

where we use the notation  $(\cdot)_{,n}$  to denote a derivative with respect to  $n$ . Taking the inner product of (34) with  $w_0$  and noting that  $\hat{h}_0 w_0 = 0$ , we find

$$\phi_{0,n} = \frac{g}{Z} \int d^3r w_0^4 + \frac{2gn}{Z} \int d^3r w_0^3 w_{0,n} : \quad (35)$$

To solve (34) for  $w_{0,n}$ , we note that the normalization condition in (22) implies

$$\int d^3r w_{0,n} w_0 = 0 : \quad (36)$$

Thus,  $w_{0,n}$  is orthogonal to  $w_0$  and has the expansion

$$w_{0,n} = \sum_{n \neq 0} a_n w_n \quad (37)$$

in terms of the (real)  $q = 0$  Bloch functions  $w_n$  with  $w_{n,q=0} = 0$ . Substituting this expansion in (34) yields

$$\sum_{n \neq 0} M_{nn^0} a_n = \frac{1}{2n} M_{nn^0} : \quad (38)$$

Using the expansion (37) for  $w_{0,n}$  in (35) with the expansion coefficients defined by (38), we find

$$\phi_{0,n} = \frac{1}{2n} \sum_{nn^0} M_{nn^0} \sum_{nn^0} M_{nn^0}^{-1} M_{nn^0} : \quad (39)$$

Comparing this with (31), we see that (32) is equivalent to

$$s = \frac{\overline{\frac{n}{m_0}}}{\overline{\frac{n}{m_0}}} = \frac{\overline{\frac{n}{m_0}}}{\overline{\frac{n}{m_0}}} \quad (40)$$

This result for an optical lattice was first given by Menotti et al. [17] on the basis of general dynamical considerations. We see here that it follows directly from the Bogoliubov equations and also applies in the case of a 3D lattice with cubic symmetry. The small- $q$  expansion can be viewed as a systematic way of implementing the slowly varying ansatz used by Kramers et al. [13]. The expression for  $s$  has the same form as for a homogeneous gas, with  $m_0$  replacing the bare mass  $m$  and the density derivative of the chemical potential,  $\phi_{0,n}$ , replacing the interaction parameter  $g$ . In other words, at long wavelengths the condensate behaves as a gas of particles of mass  $m_0$  with a compressibility,  $\chi$ , given by  $\chi^{-1} = n(\partial \phi_{0,0} / \partial n)$ .

## A. Thomas-Fermi Limit

The Thomas-Fermi (TF) approximation is valid when the density varies in space on a length scale much larger than the local coherence length  $\xi = \hbar^2 / 2mgn$ . In this situation, the density is well approximated by

$$n_0(r) = \frac{1}{g} (\phi_0 - V_{\text{opt}}(r)) \quad (41)$$

except in regions where  $V_{\text{opt}}' > \phi_0$ . This does not occur if  $\phi_0 > |V_{\text{opt}}|_{\text{max}}$ , and we then have

$$\phi_0 = gn + \bar{V}_{\text{opt}} ; \quad (42)$$

where  $\bar{V}_{\text{opt}}$  is the mean value of the optical potential in the unit cell. Thus,  $\phi_{0,n} = g$  as for a homogeneous gas. Since the effective potential,  $V_{\text{opt}} + gn_0$ , in the GP equation is a constant in the TF limit, we would expect the band mass,  $m_0$ , to be close to the free particle mass,  $m$ . One can in fact show that the deviation of  $m_0$  from  $m$  is proportional to  $V_0^2 (\hbar^2 / md)^4$ , where  $V_0$  is the amplitude of the potential modulation. Since we are assuming that  $\hbar^2 / md \ll 1, m_0 \approx m$  and the TF sound velocity is

$$s_{\text{TF}} = \sqrt{\frac{\hbar^2}{m}} ; \quad (43)$$

as for a uniform gas. It should be noted that this result is valid even when the amplitude of the density modulation is of order the mean density  $n$ , provided only that the inequality  $\hbar^2 / md \ll 1$  is everywhere satisfied.

If  $\phi_0 < |V_{\text{opt}}|_{\text{max}}$ , the Thomas-Fermi density develops 'holes' in regions where  $V_{\text{opt}} > \phi_0$ . For a one-dimensional modulation the density is disjoint, as is the case in two or three dimensions for sufficiently small density. In this situation long wavelength propagating phonon excitations cannot exist within the TF approximation since the necessary fluctuations in the number of atoms from one lattice cell to the next cannot occur. In reality, the GP density in regions where  $V_{\text{opt}} > \phi_0$  is small but finite and phonon-like excitations continue to exist. However, increasing the localization of the density in the potential minima leads to larger effective masses and eventually the sound speed  $s$  tends to zero. This behaviour cannot be described within the TF approximation.

## B. Weak-Coupling Limit

The  $g$ -dependence of  $\phi_{0,n}$  appears explicitly in (35) and implicitly through the wave function  $w_0$  which satisfies (33). To extract the dependence in the limit  $g \rightarrow 0$ , we expand the wave function as  $w_0 = w_0^{(0)} + g(w_0 = \phi_0/g)_{g=0} + \dots$ . Since  $\phi_0$  depends on  $g$  through the combination  $gn$ , we see that  $g(w_0 = \phi_0/g) = n(w_0 = \phi_0/n)$ . Thus, the combination  $nw_{0,n}$  appearing in (35) is proportional to  $g$  in the small- $g$  limit and the second term on the right

hand side of (35) is of order  $g^2$ . We then have

$$\begin{aligned} \epsilon_{0,m} = & \frac{g}{Z} \int d^3 r w_0^{(0)4} \\ & + \frac{6g^2}{Z} \int d^3 r w_0^{(0)3} (n w_{0,m} = g)_{g=0} + \end{aligned}$$

As discussed in Ref. [18], the first term accounts for the effect of the lattice on the compressibility,  $\chi$ , which decreases with increasing localization of the wave function  $w_0^{(0)}$ . The second term shows that  $\epsilon_{0,m}$  deviates from a linear dependence on  $g$  as the strength of the interaction is increased. An explicit expression for this quadratic correction can be obtained from the equivalent expression for  $\epsilon_{0,m}$  in (39) in which the two terms respectively correspond to the two integrals in (35). From the definition of the  $M_{nn^0}$  matrix in (25), we see that

$$\begin{aligned} \frac{1}{2n} M_{00} &= \frac{g}{Z} \int d^3 r w_0^{(0)4} \\ &+ \frac{4g^2}{Z} \int d^3 r w_0^{(0)3} (n w_{0,m} = g)_{g=0} + \\ \frac{1}{2n} M_{0n} &= \frac{g}{Z} \int d^3 r w_0^{(0)3} w_n^{(0)} + \\ \bar{M}_{nn^0}^{-1} &= \frac{n n^0}{n^{(0)}} + O(g) : \end{aligned}$$

Thus,

$$\begin{aligned} \epsilon_{0,m} = & \frac{g}{Z} \int d^3 r w_0^{(0)4} \\ & g^2 \frac{X}{n} \frac{6n}{n^{(0)}} \frac{1}{Z} \int d^3 r w_0^{(0)3} w_n^{(0)2} + \end{aligned}$$

Since the excitation energies are positive, we see that  $\epsilon_{0,m}$  has a negative curvature, which agrees with the numerical results in Ref. [18]. The interaction has the effect of increasing the width of the wave function which counteracts the localizing effect of the lattice potential.

### C. Weak Potential Limit

It is also of interest to obtain an expression for the sound speed in the case of a weak optical potential where perturbation theory applies. For simplicity we consider a weak one-dimensional periodic potential  $V_{\text{opt}} = V_0 \cos(Gz)$ , where  $G = 2\pi/d$ , applied to an otherwise three-dimensional system. The relevant GP equation is now one-dimensional,

$$\frac{\sim^2}{2m} \frac{d^2 w_0}{dz^2} + V_0 \cos(Gz) w_0 + g n w_0^3 = \epsilon_0 w_0 : \quad (46)$$

In treating the optical potential as a perturbation, we expand the wave function as  $w_0 = w_0^{(0)} + w_0^{(1)} + w_0^{(2)} +$  and chemical potential as  $\epsilon_0 = \epsilon_0^{(0)} + \epsilon_0^{(1)} + \epsilon_0^{(2)} +$

where the superscript here denotes the order in  $V_0$ . The properly normalized wave function in the absence of the potential is  $w_0^{(0)} = 1$  and  $\epsilon_0^{(0)} = g n$ . To first order in  $V_0$ ,  $\epsilon_0^{(1)} = 0$  and

$$(44) \quad w_0^{(1)} = \frac{V_0}{n_G^{(0)} + 2gn} \cos(Gz) ; \quad (47)$$

where  $n_G^{(0)} = \sim^2 G^2 / 2m$ . In calculating the second order contribution,  $w_0^{(2)}$ , to the chemical potential, the second order wave function,  $w_0^{(2)}$ , need not be determined, but the normalization condition  $\int_{d=2}^{R_{d=2}} dz w_0^{(0)} w_0^{(2)} = (1=2) \int_{d=2}^{R_{d=2}} dz w_0^{(1)2}$  is required. Thus, the chemical potential correct to second order in  $V_0$  is found to be

$$\epsilon_0 = g n \frac{n_G^{(0)} V_0^2}{2(n_G^{(0)} + 2gn)^2} + \quad (48)$$

Taking the derivative with respect to  $n$ , we have

$$\epsilon_{0,m} = g + \frac{2g V_0^2 n_G^{(0)}}{(n_G^{(0)} + 2gn)^3} : \quad (49)$$

The weak-coupling limit of this result to lowest order in  $g$  is  $\epsilon_{0,m} = g(1 + 2(V_0 = n_G^{(0)})^2)$ . This can be shown to agree with the expansion of the first term in (45) to second order in  $V_0$ .

To complete the calculation of the sound speed, we require an equivalent expression for the effective mass. This can be obtained by solving

$$(45) \quad \frac{\sim^2}{2m} \frac{d}{dz} + i q \sim^2 w_{0q} + V_0 \cos(Gz) w_{0q} = \epsilon_0(q) w_{0q} ; \quad (50)$$

where  $w_0$  is the solution of (46). Since the correction to the effective mass is second order in  $V_0$ , it is sufficient to consider the mean-field potential  $(g n w_0^2)$  to first order in  $V_0$ . We must then solve

$$\frac{\sim^2}{2m} \frac{d}{dz} + i q \sim^2 w_{0q} + V_0 \cos(Gz) w_{0q} = \epsilon_0(q) w_{0q} ;$$

where  $V_0^0 = V_0 n_G^{(0)} = (n_G^{(0)} + 2gn)$ . A gain treating  $V_0^0$  perturbatively, the effective mass for the lowest band is found to be

$$\frac{1}{m_0} = \frac{1}{m} \left( 1 - \frac{2V_0^2}{(n_G^{(0)} + 2gn)^2} \right) \quad (51)$$

Inserting (51) and (49) into (40), and discarding the quartic term in  $V_0$ , we arrive at the following expression for the sound speed

$$s = \frac{v_{\text{u}}}{c} \frac{gn}{m} \left( 1 - \frac{4gn V_0^2}{(n_G^{(0)} + 2gn)^3} \right) : \quad (52)$$

When  $\mu_G^{(0)} = gn$  (or  $d = 1/2$ ), this expression reduces to

$$s = \frac{v_{\text{out}}}{m} \frac{gn}{1 + \frac{V_0^2}{2(gn)^2}} \quad (53)$$

in agreement with the approximation to the sound speed obtained by Berg-Sørensen and Mølmer [5].

To make contact with Ref. [18], we introduce the recoil energy  $E_R = \hbar^2 k^2 / 2m = \mu_G^{(0)} / 4$  and define  $2V_0 = E_R$  (the parameter is called  $\beta'$  in Ref. [18]). Eq. (52) can then be rewritten as

$$\frac{s}{s_0} = 1 - \frac{2}{128(1 + \beta'^2)^3} \quad (54)$$

where  $s_0 = \frac{v_{\text{out}}}{m} \frac{gn}{gn=m}$  is the sound speed for the homogeneous gas and  $\beta'$  is the ratio  $gn/E_R = (d/\lambda)^2$ . This shows that  $s$  decreases quadratically with the strength of the optical potential, which is consistent with the numerical results in Ref. [18]. This expression is valid if  $2V_0 = \mu_G^{(0)}$ , or  $d = 1/2$ , however, this constitutes a rather limited range of the values of  $d$  of physical interest.

#### D. Radially Confined Condensates

As a final application of the results derived in this section we consider a condensate that is confined in the radial direction. To be specific, we assume a potential of the form

$$V(r) = V_{\text{opt}}(z) + V_\perp(r) \quad (55)$$

where  $V_\perp(r) = m \omega_\perp^2 r^2 / 2$ , that is, harmonic confinement in the radial ( $r$ ) direction. The optical potential is periodic in the axial direction with periodicity  $d$ . This potential approximates the situation of a long cigar-shaped trap with an axial standing light wave.

Although the geometry is quite different from that considered earlier, the previous analysis can be carried over with minor modification. The ground state GP solution has the property  $\psi_0(z+d) = \psi_0(z)$ , and has the normalization

$$\frac{1}{d} \int_{-\infty}^{\infty} dz \int_{-\infty}^{\infty} dr |\psi_0(r)|^2 = 1; \quad (56)$$

which defines the mean linear density  $\bar{n}$  along the length of the condensate. As before, it is convenient to define  $\bar{n}_0(r) = \bar{n} \psi_0(r)$ .

The Bogoliubov excitations in the present situation have a Bloch wave character along the axis and are obtained from (19) with the Hamiltonian

$$\hat{H}_0(q) = \frac{\hbar^2 q^2}{2m} + \frac{\partial}{\partial z} + iq \frac{\partial}{\partial z} + V + g \int \bar{\psi}_0^\dagger \psi_0 : \quad (57)$$

The eigenstates of  $\hat{H}_0(q)$  are now labelled by the set of quantum numbers  $f; n; m; g$ , where  $n$  is a one-dimensional band index,  $m$  is the azimuthal quantum number associated with the  $z$ -component of angular momentum, and  $g$  labels the different radial excitations. Of interest here are axially symmetric solutions ( $m = 0$ ) since these have the character of the phonon mode of interest.

The analysis after (19) is followed step by step, the only change being the integration volume used in the normalization of the states. We thus find that the sound speed is given by

$$s = \frac{v_{\text{out}}}{m_0} \frac{\partial \epsilon_0}{\partial \mu} \quad (58)$$

where  $m_0$  is the effective mass of the lowest band ( $n = 0$  and  $m = 0$ ).

Eq. (58) is of course valid in the absence of the optical potential. Treating the condensate in the TF approximation, we find  $\epsilon_0 = \frac{\hbar^2}{2m} \bar{n}^2$  and  $\partial \epsilon_0 / \partial \mu = \bar{n}^2 / 2 = \bar{n}^2 / 2$ . This gives a sound speed

$$s = \frac{v_{\text{out}}}{2m} \frac{gn_0(0)}{2m} \quad (59)$$

where  $n_0(0) = \bar{n}_0 = g$  is the density at the centre of the trap. This result was first obtained in Ref. [22] using a different method. In the weak coupling limit, (45) applies. In this case, the condensate wave function is a gaussian,

$$\psi_0^{(0)} = \frac{1}{\sqrt{\pi}} \exp \left( -\frac{m \omega_\perp^2 r^2}{2\hbar} \right);$$

and one again obtains the result in (59) for the sound speed, as found previously [23].

When the optical potential is strong the condensate becomes localized on each site. In this situation, the tight-binding approximation is a useful method for dealing with the system [10, 11, 24]. Approximating the condensate wave function as  $\psi(r) = \sum_i c_i f_i(r)$  where  $f_i(r)$  is a function localized on the  $i$ -th site and normalized to unity, the energy of the system is given approximately by

$$E_{\text{tot}} = \sum_i \epsilon_0 |c_i|^2 - t \sum_i (c_i c_{i+1} + c_{i+1} c_i) + \frac{1}{2} g \sum_i |c_i|^4;$$

where  $\epsilon_0$  is an on-site energy,  $t$  is a hopping matrix element connecting the amplitudes on nearest-neighbour sites and  $g$  is an effective interaction strength. For the ground state,  $|c_i|^2 = \bar{n}$ , where  $\bar{n}$  is the number of atoms per site. We then have

$$E_{\text{tot}} = (\epsilon_0 - 2t)N + \frac{1}{2} g N \bar{n};$$

Assuming that the parameters  $\epsilon_0$ ,  $t$  and  $g$  are density independent in the extreme tight-binding limit, we

have  $\epsilon_0 = \epsilon_{\text{tot}} = \epsilon_N = (\epsilon_0 - 2t) + g$  and  $\epsilon_0 = \epsilon = g$ . Within the same approximation the band energy is  $\epsilon(q) = \epsilon_0 + g - 2t \cos(qd)$ , which gives an effective mass  $m_0 = \hbar^2 / 2td^2$ . Thus the Bogoliubov sound speed from (58) is  $v_{\text{tb}} = \frac{g}{2t} \frac{td^2}{\epsilon_0} = \frac{g}{2t}$ , which is the result obtained by Javanainen [10].

#### IV. PHONON DISPERSION FOR A MOVING CONDENSATE

We turn next to the derivation of the phonon dispersion relation for the case where the condensate is "moving" through the optical lattice. We address this problem by directly solving the Bogoliubov equations in (16) for a specific condensate wave function  $w_{nk}$  in the limit of small wave vectors  $q$ . The structure of these equations is quite different when  $k \neq 0$  and the method used in the previous section to determine the dispersion relation can no longer be applied. In fact, the analysis is much more intricate as will soon become apparent. The results we obtain confirm the more intuitive hydrodynamic approach presented recently [7], which describes the dynamics of the system in terms of slowly varying hydrodynamic variables (density and momentum). By including small length scale variations, our approach in a sense provides a "microscopic" derivation of the hydrodynamic equations that one would expect to be valid in the long wavelength limit.

Since the solution in the small- $q$  limit is required, we rewrite (16) so as to display the  $q$ -dependent terms explicitly:

$$\begin{aligned} \hat{h}_k + \frac{\hbar^2 q^2}{2m} (p + \hbar k) + \frac{\hbar^2 q^2}{2m} + g n \hat{w}_{0k} \int u \\ g n w_{0k}^2 v = E u \\ \hat{h}_k + \frac{\hbar^2 q^2}{2m} (p - \hbar k) + \frac{\hbar^2 q^2}{2m} + g n \hat{w}_{0k} \int v \\ g n w_{0k}^2 u = -E v; \quad (60) \end{aligned}$$

where  $p = (\hbar/i)r$  is the momentum operator. The GP Hamiltonian here is  $\hat{h}_k (q = 0)$  as defined in (18). In these equations, we have adopted the index  $n = 0$  for the condensate wave function. We will usually think of this state as the lowest Bloch state solution of the GP equation, although it in principle could correspond to an arbitrary excited band. For simplicity we have dropped the index on the quasiparticle amplitudes  $u$  and  $v$  and the excitation energy  $E$  as we will only be considering the phonon-like excitation.

It is clear that for  $q = 0$  (60) admits a solution with  $u \propto w_{0k}$ ,  $v \propto w_{0k}$  and  $E = 0$ . For finite  $q$  we seek solutions in the form of an expansion in eigenfunctions of  $\hat{h}_k$ , namely,

$$u = \sum_n a_n(q) w_{nk}$$

$$v = \sum_n b_n(q) w_{n-k} \quad (61)$$

where

$$\hat{h}_k w_{nk} = \epsilon_{nk} w_{nk} \quad (62)$$

According to this definition,  $\epsilon_{0k} = 0$ . The functions  $w_{nk}$  are an orthonormal set with normalization given by (22). Although we use the same notation, it should be noted that these functions are distinct from those defined in (21). In addition, we may assume  $w_{n-k}(r) = w_{nk}(r)$  and  $\epsilon_{n-k} = \epsilon_{nk}$ .

Substituting these expansions into (60), we obtain the matrix equations

$$\begin{aligned} \epsilon_{nk} + \frac{\hbar^2 q^2}{2m} a_n + \sum_{n^0} \frac{\hbar^2 q}{m} P_{nn^0} a_{n^0} \\ + \sum_{n^0} (A_{nn^0} a_{n^0} - B_{nn^0} b_{n^0}) = E a_n; \quad (63) \end{aligned}$$

$$\begin{aligned} \epsilon_{nk} + \frac{\hbar^2 q^2}{2m} b_n + \sum_{n^0} \frac{\hbar^2 q}{m} P_{nn^0} b_{n^0} \\ + \sum_{n^0} (A_{nn^0} b_{n^0} - B_{nn^0} a_{n^0}) = -E b_n; \quad (64) \end{aligned}$$

where we have defined the matrices

$$A_{nn^0}(k) = \frac{1}{Z} \int w_{nk} g n \hat{w}_{0k} \hat{w}_{n^0k} d^3 r; \quad (65)$$

$$B_{nn^0}(k) = \frac{1}{Z} \int w_{nk} g n w_{0k}^2 w_{n^0k} d^3 r; \quad (66)$$

$$P_{nn^0}(k) = \frac{1}{Z} \int w_{nk} (p + \hbar k) w_{n^0k} d^3 r; \quad (67)$$

Due to the inversion symmetry of the lattice, the Bloch states at the zone centre ( $k = 0$ ) can be chosen to be simultaneous eigenstates of parity. Although parity is not a good quantum number for states with nonzero  $k$ , each band can nevertheless be assigned a parity index  $\pi_n = \pm 1$  such that

$$w_{n-k}(r) = \pi_n w_{nk}(r); \quad (68)$$

This property is proved explicitly in one dimension in Appendix A and can also be shown to follow to lowest order in  $k$  by means of  $k$ -perturbation theory. Together with the conjugation (time-reversal) property  $w_{nk}(r) = w_{n-k}(r)$ , we have

$$w_{nk}(r) = \pi_n w_{nk}(\pi r); \quad (69)$$

This important property is used throughout the following discussion. For example, it can be used to show that the matrices in (65-67) satisfy the following relations

$$\begin{aligned} A_{nn^0}(k) &= A_{n^0n}(k) = A_{nn^0}(-k) = \pi_n \pi_{n^0} A_{nn^0}(k) \\ B_{nn^0}(k) &= B_{nn^0}(-k) = \pi_n \pi_{n^0} B_{nn^0}(k) = \pi_n \pi_{n^0} B_{n^0n}(k) \\ P_{nn^0}(k) &= P_{n^0n}(k) = P_{nn^0}(-k) = \pi_n \pi_{n^0} P_{nn^0}(k) \end{aligned} \quad (70)$$



Note that  $A$  and  $P$  are hermitian while  $B$  is not. In addition, we see that  $A_{nn^0}(0)$  and  $B_{nn^0}(0)$  are real and nonzero only for pairs of states having the same parity index, while  $P_{nn^0}(0)$  is purely imaginary and only couples states with opposite parity.

In solving (63) and (64), it is convenient to define the following linear combinations:  $c_n = \frac{1}{2}(a_n + b_n)$  and  $d_n = \frac{1}{2}(a_n - b_n)$ . Introducing these variables into (63) and (64), and making use of the relations in (70), we obtain the equations

$$\begin{aligned} \mu_{nk} + \frac{\tilde{q}^2}{2m} c_n + \sum_{n^0} A_{nn^0} \mathcal{B}_{nn^0} c_{n^0} \\ + \sum_{n^0} \frac{\tilde{q}}{m} q_{nn^0} d_{n^0} = E d_n; \end{aligned} \quad (71)$$

$$\begin{aligned} \mu_{nk} + \frac{\tilde{q}^2}{2m} d_n + \sum_{n^0} A_{nn^0} + \mathcal{B}_{nn^0} d_{n^0} \\ + \sum_{n^0} \frac{\tilde{q}}{m} q_{nn^0} c_{n^0} = E c_n; \end{aligned} \quad (72)$$

where we have defined the hermitian matrix  $\mathcal{B}_{nn^0} = B_{nn^0} - A_{nn^0}$ . As in our earlier analysis, we anticipate that  $E(q)$  will depend linearly on  $q$  for  $q \neq 0$ , but due to the quasimomentum  $k$  of the condensate, it no longer depends simply on the magnitude  $q$ . To extract this dependence we systematically expand the coefficients  $c_n(q)$  and  $d_n(q)$  as a series in powers of  $q$ . Specifically, we write

$$\begin{aligned} c_n(q) &= h(q)(c_n^{(0)} + c_n^{(1)} + c_n^{(2)}) \\ d_n(q) &= h(q)(d_n^{(0)} + d_n^{(1)} + d_n^{(2)}) \end{aligned} \quad (73)$$

where the superscript indicates the order of  $q$  in the respective terms (here, order signifies similar powers of the vector magnitude  $q$  [25]). The factor  $h(q)$  contains the nonanalytic behaviour of the coefficients which is required in order to satisfy the normalization condition in (15). For a homogeneous system,  $h(q) \propto q^{-1/2}$ , and we expect a similar dependence in the case of a lattice. In the following, it is sufficient to note that this factor is the same for both coefficients and can therefore be ignored in developing a systematic  $q$ -expansion.

We noted earlier that  $u/w_{0k}$  and  $v/w_{0k}$  for  $q \neq 0$  which, according to (61), implies that  $a_n(q) \propto n_0$  and  $b_n(q) \propto n_0$  in this limit. If the state  $w_{0k}$  is an even parity state ( $\ell_0 = +1$ ), as we assume in the following, we must then have  $c_n^{(0)} = n_0$  and  $d_n^{(0)} = 0$ . With this information, (71) gives to  $O(q^0)$  the equation

$$\mu_{0k} + (A_{n0} - \mathcal{B}_{n0}) = 0 \quad (74)$$

which is satisfied since  $\mu_{0k} = 0$  and  $\mathcal{B}_{n0} = A_{n0}$ . To  $O(q^1)$ , (71) gives

$$\mu_{nk} c_n^{(1)} + \sum_{n^0} (A_{nn^0} - \mathcal{B}_{nn^0}) c_{n^0}^{(1)} = 0; \quad (75)$$

Similarly, (72) gives

$$\sum_{n^0} N_{nn^0} d_{n^0}^{(1)} = E_{n0} \frac{\tilde{q}}{m} q_{nn^0}; \quad (76)$$

where we have defined the hermitian matrix

$$N_{nn^0}(k) = A_{nn^0}(k) + \mathcal{B}_{nn^0}(k) + \mu_{nk} - \mu_{n^0k}; \quad (77)$$

Eq. (75) is homogeneous and indicates that  $c_n^{(1)} \neq 0$ . On the other hand, (76) can be solved for  $d_n^{(1)}$  in terms of the unknown excitation energy  $E$ . To determine the latter, we must consider (71) to  $O(q^2)$ , obtaining

$$\begin{aligned} \mu_{nk} c_n^{(2)} + \frac{\tilde{q}^2}{2m} n_0 + \sum_{n^0} (A_{nn^0} - \mathcal{B}_{nn^0}) c_{n^0}^{(2)} \\ + \sum_{n^0} \frac{\tilde{q}}{m} q_{nn^0} d_{n^0}^{(1)} = E d_n^{(1)}; \end{aligned} \quad (78)$$

Setting  $n = 0$  in this equation, and noting that  $\mu_{0k} = 0$  and that  $A_{0n} = \mathcal{B}_{0n}$ , we find

$$\frac{\tilde{q}^2}{2m} + \sum_n \frac{\tilde{q}}{m} q_{nn} d_n^{(1)} = E d_0^{(1)}; \quad (79)$$

Since  $d_n^{(1)}$  is itself linear in  $E$  according to (76), we see that (79) is implicitly a quadratic equation for  $E$  which can be solved to determine the excitation energy to lowest order in  $q$ . However, to do so directly would not reveal the interesting dependences on various physical parameters that in fact emerge. As seen in the  $k = 0$  analysis, the excitation energy could be related to the variation of the chemical potential with mean density  $n$ . This remains a quantity of interest in the present case, but we must also consider variations of the chemical potential with  $k$ . We thus turn next to the determination of  $\partial \mu_{0k} / \partial n$  and  $\partial \mu_{k1} / \partial k_1$ .

#### A. Determination of $\partial \mu_{0k} / \partial n$

The chemical potential  $\mu_{0k}$  is determined by the self-consistent solution of the GP equation,  $\hat{h}_k w_{0k} = 0$ , with  $\hat{h}_k$  given by (18) with  $q = 0$ . The variation of this equation with respect to  $n$  gives

$$\hat{h}_k w_{0k;n} + g \int w_{0k}^2 w_{0k} + g n w_{0k;n} w_{0k} + w_{0k} w_{0k;n} w_{0k} = \mu_{0k;n} w_{0k}; \quad (80)$$

where  $w_{0k;n} \equiv \partial w_{0k} / \partial n$ . Taking the derivative of the normalization (22) with respect to  $n$ , and using (69), we find that

$$\int w_{0k} w_{0k;n} d^3r = 0; \quad (81)$$

Thus,  $w_{0k;n}$  is orthogonal to  $w_{0k}$  and has the expansion

$$w_{0k;n} = \sum_n w_{nk}; \quad (82)$$

where as before, the prime on the summation indicates that the  $n = 0$  term is excluded from the sum. Inserting this expansion into (80), and taking the inner product with respect to  $w_{nk}$ , we obtain

$$\frac{1}{2n} N_{n0} + \sum_{n^0}^X N_{nn^0} N_{n^0} \quad 0k;n \quad n^0 = 0; \quad (83)$$

where we have noted that  $N_{nn} = N_{nn}$  as a result of the symmetry property (68). Setting  $n = 0$  in (83), we find

$$0k;n = \frac{1}{2n} N_{00} + \sum_n^X N_{0n} N_{nn} : \quad (84)$$

The set of equations for  $n \neq 0$  has the solution

$$n = \frac{1}{2n} \sum_{n^0}^X (\mathbb{P}^{-1})_{nn^0} N_{n^0 0}; \quad (85)$$

where  $\mathbb{P}$  is the reduced matrix obtained by deleting the first row and first column from  $N$ . Thus we find that

$$0k;n = \frac{1}{2n} N_{00} + \sum_{nn^0}^X N_{0n} (\mathbb{P}^{-1})_{nn^0} N_{n^0 0} : \quad (86)$$

This is analogous to (39) and reduces to it in the  $k = 0$  limit since the  $M$  and  $N$  matrices are then the same (note that  $w_{n0}$  are defined to be real).

#### B. Determination of $0k;i$

We follow a similar method to obtain  $0k;i$ . Taking the derivative of  $\hat{h}_k w_{0;k} = 0$  with respect to  $k_i$ , we have

$$\frac{\hbar}{m} p_i + \tilde{w}_{k_i} + g n w_{0k;i} w_{0k} + w_{0k} w_{0k;i} \quad 0k;i \quad w_{0k} + \hat{h}_k w_{0k;i} = 0; \quad (87)$$

where  $w_{0k;i} = \partial w_{0k} / \partial k_i$ . Noting the orthogonality of each vector component  $w_{0k;i}$  with  $w_{0k}$ , we have the following expansion

$$w_{0k;i} = \sum_n^X n_{in} w_{nk}; \quad (88)$$

where the expansion coefficients,  $n_{in}$ , define the Cartesian components of a vector  $n$ . Inserting (88) into (87), and taking the inner product with  $w_{nk}$ , we obtain

$$\frac{\hbar}{m} (P_i)_{n0} + \sum_{n^0}^X N_{nn^0} n_{in^0} \quad 0k;i \quad n^0 = 0: \quad (89)$$

where, as before, we have used  $n_{in} = n_{in}$ . An expression for  $0k;i$  can be found by setting  $n = 0$  in (89):

$$0k;i = \frac{\hbar}{m} (P_i)_{00} + \sum_n^X N_{0n^0} n_{in^0} : \quad (90)$$

The set of equations for  $n \neq 0$  yield the solution vector,

$$n = \frac{\hbar}{m} \sum_{n^0}^X (\mathbb{P}^{-1})_{nn^0} (P_i)_{n^0 0}; \quad (91)$$

and thus,

$$0k;i = \frac{\hbar}{m} (P_i)_{00} + \sum_{nn^0}^X N_{0n} (\mathbb{P}^{-1})_{nn^0} (P_i)_{n^0 0} : \quad (92)$$

#### C. Excitation Energy

These results will now be used to obtain an expression for the excitation energy  $E$  from (76) and (79). Setting  $n = 0$  in (76) we have

$$\sum_{n^0}^X N_{0n^0} d_{n^0}^{(1)} = E \frac{\hbar}{m} q \quad E_0 \quad N_{00} d_0^{(1)} \quad (93)$$

while for  $n \neq 0$  we see that

$$d_n^{(1)} = \sum_{n^0}^X (\mathbb{P}^{-1})_{nn^0} \frac{\hbar}{m} q \quad E_0 + N_{n^0 0} d_0^{(1)} \quad (94)$$

The quantities on the right hand side of this equation are in fact related to the expansion coefficients  $n$  in (85) and  $n$  in (91). We find the simple relation

$$d_n^{(1)} = 2n d_0^{(1)} + q \quad n : \quad (95)$$

Using this result in (93), we have

$$N_{00} + 2 \sum_n^X N_{0n} n \quad d_0^{(1)} = E \frac{\hbar}{m} q \quad E_0 \quad \sum_n^X N_{0n} q \quad n \quad (96)$$

which with (84) and (90) can be written as

$$2 \sum_n 0k;n d_0^{(1)} = E \quad q \quad 0k : \quad (97)$$

We see that  $d_0^{(1)}$  and  $E$  are now related to each other through physically meaningful and calculable parameters.

We now substitute (95) into (79) to obtain

$$\frac{\hbar^2 q^2}{2m} + \sum_n^X \frac{\hbar}{m} (q \quad E_n) (q \quad n) = E \frac{\hbar}{m} q \quad E_0 \quad \sum_n^X \frac{\hbar}{m} q \quad E_n 2n \quad n \quad d_0^{(1)} : \quad (98)$$

With (85), the sum on the right hand side becomes

$$\begin{aligned} \sum_n^X \frac{\hbar}{m} P_{0n} 2n \quad n &= \sum_{nn^0}^X \frac{\hbar}{m} P_{0n} (\mathbb{P}^{-1})_{nn^0} N_{n^0 0} \\ &= \sum_{nn^0}^X N_{0n} (\mathbb{P}^{-1})_{nn^0} \frac{\hbar}{m} P_{n^0 0} \\ &= \sum_n^X N_{0n} n : \end{aligned} \quad (99)$$

In going from the first to the second line, we have used the fact that all the matrices have the transposition property  $M_{n'n} = M_{nn'}^0$ . With the expression for  $\epsilon_{0k;i}$  in (90), (98) thus becomes

$$\frac{\tilde{\omega}^2 q^2}{2m} + \sum_n \frac{\tilde{\omega}^2}{m} (q - \epsilon_n) (q - \epsilon_n) = (E - q - \epsilon_{0k}) d_0^{(1)} \quad (100)$$

Eliminating  $d_0^{(1)}$  from (97) and (100), we finally obtain

$$E = q + \frac{\epsilon_{0k}}{2n_{0k;n}} + \frac{\tilde{\omega}^2 q^2}{2m} + \sum_n \frac{\tilde{\omega}^2}{m} (q - \epsilon_n) (q - \epsilon_n) : (101)$$

The sign of the square root is chosen to be positive to give a positive excitation energy in the  $k \rightarrow 0$  limit. The natural quantity to interpret is the summation within the square root.

#### D. Effective Mass Tensor

The square bracket in (101) involves the tensor

$$\frac{1}{m} \epsilon_{ij} = \frac{1}{m} \epsilon_{ij} + \frac{2}{m} \sum_n \epsilon_{i)0n} (\epsilon_{j})_n = \frac{1}{m} \epsilon_{ij} + \frac{2}{m^2} \sum_n \epsilon_{i)0n} (\epsilon_{j})_{nn^0} (\epsilon_{j})_{n^0} : (102)$$

This expression is similar to the usual effective mass tensor defined on the basis of  $k$  perturbation theory, although the structure of the summation is different. To make contact with the  $k$  perturbation expression we consider the  $N_{nn^0}$  matrix in the  $k \rightarrow 0$  limit. Quite generally, this matrix has the block structure

$$N(k) = \begin{pmatrix} A_{++} + B_{++} + D & A_+ & B_+ \\ A_+ + B_+ & A & B \\ A_+ + B_+ & A & B + D \end{pmatrix} \quad (103)$$

where the blocks are defined according to the parity index of the various states. For example, the block in the upper-left corner contains matrix elements between states with a positive parity index,  $\epsilon = +1$ . The diagonal matrix  $D$  contains the energy eigenvalues  $\epsilon_{nk}$  on its diagonal. In the limit  $k \rightarrow 0$ , we have  $B(k=0) = A(k=0)$  and  $A_{nn^0} = 0$  if  $n \neq n^0$ . Thus, in this limit we have

$$N(k=0) = \begin{pmatrix} 2A_{++} + D & 0 \\ 0 & D \end{pmatrix} \quad (104)$$

that is,  $N(k=0)$  is block-diagonal, which of course is also true of its inverse. Since,  $P_i$  only connects states of opposite parity in the  $k \rightarrow 0$  limit, we thus see that

$$\lim_{k \rightarrow 0} \frac{1}{m} \epsilon_{ij} = \frac{1}{m} \epsilon_{ij} + \frac{2}{m^2} \sum_n \epsilon_{i)0n} (\epsilon_{j})_{n0} : (105)$$

This is precisely the effective mass tensor obtained by means of  $k$  perturbation theory [27] as applied to the Hamiltonian  $\hat{H}_0(q)$  in (20). The tensor defined in (102) is a generalized effective mass tensor in that it depends on the presence of a superfluid flow ( $k \neq 0$ ). Also because of this, it is no longer diagonal despite the cubic symmetry of the optical lattice.

To complete the identification of  $(m^{-1})_{ij}$ , we consider variations of the GP equation with respect to the condensate wave vector  $k$ . The second derivative of  $\hat{H}_k w_{0k} = 0$  yields the equation

$$\hat{H}_{k;ij} w_{0k} + \hat{H}_{k;i} w_{0k;j} + \hat{H}_{k;j} w_{0k;i} + \hat{H}_k w_{0k;ij} = 0 : (106)$$

Here,

$$\hat{H}_{k;i} = \frac{\tilde{\omega}^2}{m} (p_i + \tilde{k}_i) + g n_{c;i} \epsilon_{0k;i} ; (107)$$

$$\hat{H}_{k;ij} = \frac{\tilde{\omega}^2}{m} \epsilon_{ij} + g n_{c;ij} \epsilon_{0k;ij} : (108)$$

The inner product of (106) with  $w_{0k}$  gives

$$\frac{1}{Z} \int w_{0k} (\hat{H}_{k;ij} w_{0k} + \hat{H}_{k;i} w_{0k;j} + \hat{H}_{k;j} w_{0k;i} + \hat{H}_k w_{0k;ij}) d^3 r = 0 : (109)$$

The first integral is

$$\frac{1}{Z} \int w_{0k} \hat{H}_{k;ij} w_{0k} d^3 r = \frac{\tilde{\omega}^2}{m} \epsilon_{ij} \epsilon_{0k;ij} + \frac{g}{n} n_{c;n_{c;ij}} d^3 r : (110)$$

while the integral of the next two terms gives the result

$$\begin{aligned} \frac{1}{Z} \int w_{0k} (\hat{H}_{k;i} w_{0k;j} + \hat{H}_{k;j} w_{0k;i}) d^3 r \\ = \frac{\tilde{\omega}^2}{m} \sum_n \epsilon_{i)0n} (\epsilon_{j})_n + (\epsilon_{j})_{0n} (\epsilon_{i})_n \\ + \frac{g}{n} n_{c;i} n_{c;j} d^3 r : (111) \end{aligned}$$

We thus find

$$\epsilon_{0k;ij} = \frac{\tilde{\omega}^2}{m} \epsilon_{ij} + \frac{2}{m} \sum_n \epsilon_{i)0n} (\epsilon_{j})_n + \frac{g}{2n} (n_c^2)_{ij} d^3 r : (112)$$

With this result we see that the effective mass tensor defined in (102) can be expressed as

$$\frac{1}{m} \epsilon_{ij} = \frac{1}{\tilde{\omega}^2} \frac{\partial^2}{\partial k_i \partial k_j} \epsilon_{0k} + \frac{g}{2n} n_c^2 d^3 r : (113)$$

#### E. Relation to Energy Density

This last result can be related to the total energy (3) in the state  $\epsilon_{0k}$ . Defining the energy per particle as  $E_{\text{tot}} = N \sim \langle n; k \rangle$ , we have

$$\begin{aligned} \langle n; k \rangle = \frac{1}{Z} \int w_{0k} \left[ \frac{\tilde{\omega}^2}{2m} (r + ik)^2 + V_{\text{opt}} \right] w_{0k} d^3 r \\ + \frac{gn}{2} \int w_{0k}^4 d^3 r \end{aligned} \quad (114)$$

Comparing this with  $\epsilon_{0k}$ , we see that

$$\epsilon(n; k) = \epsilon_{0k} \frac{gn}{2} \int_0^Z \int_{\Omega_k} d^3r \quad (115)$$

which is the expression in brackets in (113). Thus the effective mass tensor is given by

$$\frac{1}{m_{ij}} = \frac{1}{\epsilon^2} \frac{\partial^2 \epsilon}{\partial k_i \partial k_j} : \quad (116)$$

For a cubic lattice,  $\epsilon_{0k}$  has an expansion of the form  $\epsilon_{0k} = \epsilon_{00} + \frac{1}{2} k^2 = 2m_0 + \dots$ . Similarly,  $\epsilon(n; k) = \epsilon(n; 0) + \frac{1}{2} k^2 = 2m_0 + \dots$ . However, as proved by (105), the parameter is in fact the band mass  $m_0$  defined by the Hamiltonian (20). In other words, the correct effective mass parameter can be extracted without solving the GP equation for  $w_{0k}$  (with  $k \neq 0$ ) self-consistently. We note in passing that direct differentiation of (115) establishes the relation  $\hbar j_i = r_k (n) = \epsilon$  [20].

With these results, the phonon energy (101) can be written in a compact form. Denoting the mean energy density as  $\epsilon = n\epsilon$ , the Bogoliubov excitation energy at long wavelengths is given by

$$E = \epsilon_{ni} q_i + \frac{p}{\epsilon_{mn} \epsilon_{ij} q_i q_j} \quad (117)$$

where we use a repeated summation convention on the Cartesian indices  $i$  and  $j$ . This is precisely the expression given by Machholm et al. [7] who argued that the dynamics of the system at long wavelengths could be based on a hydrodynamic analysis. Since their approach arrives at (117) in a more economical fashion, it is useful to summarize the essential assumptions on which it is based.

The central assumption is the existence of an average phase fluctuation,  $\hbar(r; t)$ , that varies slowly in space and time. Expanding this average phase as

$$\hbar(r + r'; t + t') = \hbar(r; t) + r' \cdot \nabla \hbar(r; t) + \frac{1}{2} \nabla^2 \hbar(r; t) r'^2 + \dots; \quad (118)$$

one identifies  $r \cdot \nabla \hbar$  with the local wave vector,  $\hbar k$ , and  $\partial \hbar / \partial t$  with  $\hbar \omega$  where  $\hbar \omega$  is the local chemical potential. The equation of motion for the local wave vector is thus

$$\frac{\partial \hbar k}{\partial t} = r \cdot \nabla \hbar \omega : \quad (119)$$

The second hydrodynamic equation is the continuity equation

$$\frac{\partial \hbar n}{\partial t} + r \cdot \nabla \hbar j = 0; \quad (120)$$

where  $\hbar j$  is the local current density. The current density and chemical potential are then assumed to be given by the usual expressions for a uniform system, namely,

$$\hbar j = \frac{1}{m} r \cdot \nabla \hbar \omega; \quad \hbar \omega = \frac{\partial \epsilon}{\partial \hbar n} \quad (121)$$

where  $\epsilon(\hbar n; \hbar k)$  is the average energy density for a uniform optical lattice, viewed as a function of the local density  $\hbar n$  and wave vector  $\hbar k$  [26]. By expanding the variables as  $\hbar n = n + \delta n$  and  $\hbar k = k + \delta k$ , one obtains a pair of equations for the fluctuations which admit wave-like solutions with frequency  $\omega = E/\hbar$  and wave vector  $q$ . The dispersion relation found is identical to (117).

It is clear that the assumptions made in the hydrodynamic approach are completely justified. The average energy density  $\epsilon$  is the fundamental quantity determining the excitation energy at long wavelengths, as confirmed by our systematic  $q$ -expansion. The additional information provided by the expansion technique are the perturbative expressions for  $\partial \epsilon_{0k} / \partial n$ ,  $r_k \epsilon_{0k}$  and  $(1/m)_{ij}$  as given by (86), (92) and (102), respectively.

## F. Discussion

For small  $k$ ,  $\epsilon(n; k) = \epsilon(n; 0) + \frac{1}{2} k^2 = 2m_0 + \dots$ , and the Bogoliubov excitation energy is

$$E = \hbar \omega + \hbar q \cdot \frac{\partial \epsilon}{\partial n} \frac{n}{m_0} + s q \quad (122)$$

where  $s$  is the sound speed for the condensate at rest. This result was given previously by Kramer et al. [18]. The energy first becomes negative when the superfluid flow satisfies  $k > m s = \hbar \omega$ , where  $1/m = \partial \epsilon / \partial n = \partial \epsilon / \partial n_0 = \partial \epsilon / \partial n$ , which defines the Landau criterion for energetic instability at long wavelengths in an optical lattice. The region of energetic instability was mapped out for arbitrary  $q$  by Wu and Niu [6, 28] and Machholm et al. [7]. In this region the energy of the superfluid state is no longer a local maximum. As a result, transitions to lower energy states can occur spontaneously provided a means of conserving energy and quasimomentum is available.

The excitation energy given by (117) becomes imaginary when the argument of the square root is negative. This signals a dynamic instability whereby the amplitude of the condensate fluctuation grows (or decays) in time. Of the two factors in the square root,  $\epsilon_{ij}$ , or equivalently the effective mass tensor,  $(1/m)_{ij}$ , is the most physically relevant.

It is instructive to examine the latter in the weak potential limit. We consider for simplicity the one-dimensional situation discussed in Sec. IIIC. Repeating the perturbative analysis in Sec. IIIC for the case  $k \neq 0$ , we find

$$\epsilon(n; k) = \frac{1}{2} gn + \frac{V_0^2}{2 \epsilon_G^{(0)} + 2gn} + \frac{4 \epsilon_k^{(0)}}{4 \epsilon_k^{(0)}} \quad (123)$$

We note that this expression becomes singular at a wave vector  $k_c$  satisfying  $\epsilon_G^{(0)} + 2gn - 4 \epsilon_k^{(0)} = 0$ , which gives

$$k_c = \frac{q}{k_0^2 + G^2} = 4 \quad (124)$$

where  $k_0 = m_0 s_0 \sim \dots$ . The singularity is indicating the breakdown of nondegenerate perturbation theory, but provided  $k$  is not too close to  $k_c$ , we can use (123) to evaluate the effective mass. To second order in  $V_0$  we have

$$\frac{1}{m_0(k)} = \frac{1}{m} \left[ 1 - \frac{2V_0^2}{\epsilon_0^{(0)} + 2gn} \frac{\epsilon_k^{(0)2}}{4} + \frac{32\epsilon_k^{(0)}V_0^2}{(\epsilon_0^{(0)} + 2gn)^3} + \dots \right] \quad (125)$$

At  $k = 0$  we recover the result (51) found in Sec. III C. For  $m_0^{-1}$  to go to zero,  $k$  must be close to  $k_c$ . With  $k = k_c + \delta k$ , we find

$$\frac{k}{k_c} \sim \left( \frac{V_0}{\epsilon_0^{(0)} + 2gn} \right)^{2/3}; \quad (126)$$

that is, the wave vector  $k$  approaches  $k_c$  as  $V_0 \rightarrow 0$ . We note that at  $k = k_c + \delta k$ , the perturbative correction to the energy in (123) is still small so that the perturbation theory estimate of where  $m_0^{-1}$  goes to zero is reasonable. We thus expect a dynamical instability to set in when  $k \rightarrow k_c$  in the weak potential limit.

This condition for the dynamical instability is the  $q = 0$  limit of the result given in Refs. [6] and [7]. The Bogoliubov excitations of wave vector  $q$  in an homogeneous gas with current  $\hbar j_i = n \hbar k_i$  have the energies

$$E_{\pm}(q) = \frac{\hbar^2 k q}{m} \pm \sqrt{\frac{n g \hbar^2 q^2}{m} + \frac{\hbar^4 q^4}{4m^2}}; \quad (127)$$

We follow Wu and Niu [6] in referring to the modes with the plus (minus) sign as phonons (antiphonons). The former correspond to physical excitations in that their normalization is given by (15). The effect of an optical potential is to couple an antiphonon mode with wave vector  $q$  to a phonon mode with wave vector  $q - G$ . The condition that  $E_{-}(q) = E_{+}(q - G)$  implies that the two modes are resonantly coupled and gives the critical wave vector

$$k_c = \frac{1}{G} \sqrt{k_0^2 q^2 + \frac{1}{4} q^4} + \frac{1}{k_0^2 (G - q)^2 + \frac{1}{4} (G - q)^4}; \quad (128)$$

For  $q = 0$ , this gives the critical wave vector in (124). The expression in (128) was shown in [6] to account for the boundary of the dynamically unstable region in the weak potential limit. In fact, it can be shown by means of degenerate perturbation theory (Appendix B) that imposing a weak optical potential indeed gives rise to complex Bogoliubov eigenvalues.

Alternatively, the condition  $E_{-}(q) = E_{+}(q - G)$  can be written as  $E_{+}(q) + E_{+}(q - G) = 0$ . This was interpreted by Machholm et al. as a Landau criterion for the emission of two phonon excitations with zero total energy.

Although this physical interpretation is appealing, it is not clear how it can be used to actually determine the rate at which the excitations are being produced, short of performing the perturbation analysis carried out in Appendix B in terms of phonon and antiphonon modes.

We thus see that the phonon-antiphonon resonance condition, or alternatively the two-phonon Landau criterion, is consistent with the effective mass condition for a dynamical instability in the  $q \rightarrow 0$  limit. A similar statement can be made in the weak coupling limit ( $g \rightarrow 0$ ). Wu and Niu [6] noted from their numerical analysis that one boundary of the dynamically unstable region is given by the condition  $\epsilon_0(q + k) - \epsilon_0(k) = \epsilon_0(k) - \epsilon_0(k - q)$  where  $\epsilon_0(k)$  is the band energy for the optical potential by itself. In the small- $q$  limit, this condition becomes

$$\frac{\partial^2 \epsilon_0(k)}{\partial k^2} = 0;$$

Thus the onset of dynamical instability at  $q = 0$  in the weak coupling limit is again given by the point at which the inverse effective mass goes to zero.

## V. CONCLUSIONS

We have studied the long wavelength phonon excitations in a three dimensional optical lattice. By making use of a systematic expansion of the Bogoliubov equations in terms of the phonon wave vector  $q$ , we obtain the phonon dispersion in the long wavelength limit. Our result (40) for the current-free state defines the sound speed in terms of the effective mass  $m_0$  and variations of the chemical potential with  $n$  and agrees with the result given by Menotti et al. [17]. The effective mass is defined quite generally in terms of the energy per particle,  $\epsilon(n; k)$ , but can also be calculated using the current-free GPHamiltonian in the  $k \rightarrow 0$  limit. We present analytic expressions for the sound speed in the Thomas-Fermi, weak potential, weak coupling and tight-binding limits.

For the current-carrying case, we rederive the dispersion relation obtained by means of a hydrodynamic analysis [7] (see also [13]). Our approach confirms that the dynamics at long wavelengths is defined by the local energy density  $\epsilon(n; \hbar k)$  viewed as a function of the slowly varying local density,  $n(r)$ , and local condensate wave vector,  $\hbar k(r)$ . At long wavelengths, dynamical instabilities arise at the point where the generalized effective mass tensor has a vanishing eigenvalue.

## APPENDIX A: REFLECTION SYMMETRY OF BLOCH FUNCTIONS

In this Appendix we give a proof of the symmetry property (68) used throughout our analysis. We do this for the one-dimensional case for which the wave function is

a solution of

$$\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi(x) + V(x) \psi(x) = E \psi(x); \quad (A1)$$

where the potential is periodic,  $V(x+d) = V(x)$ , and is assumed to have inversion symmetry,  $V(-x) = V(x)$ . In the context of the GP equation,  $V(x) = V_{\text{opt}}(x) + g n_c(x)$  and the inversion property is valid if the condensate density also satisfies  $n_c(-x) = n_c(x)$ . This is ensured if the wave function has the property we wish to prove.

We seek solutions of the Bloch form,  $\psi(x+d) = e^{ikd} \psi(x)$ . Due to the inversion symmetry, the linearly independent solutions of (A1) can be chosen to be even ( $\psi_e$ ) or odd ( $\psi_o$ ) functions of  $x$  and  $\psi(x)$  can be expressed as the linear combination

$$\psi(x) = a \psi_e(x) + b \psi_o(x); \quad (A2)$$

The periodic part of the Bloch function is then

$$w_k(x) = e^{ikx} (a \psi_e(x) + b \psi_o(x)); \quad (A3)$$

The two independent solutions at energy  $E$  are chosen to have the normalization

$$\frac{1}{d} \int_{-d/2}^{d/2} |\psi_{e/o}(x)|^2 dx = 1; \quad (A4)$$

Imposing the Bloch condition, we obtain the relation

$$\frac{\psi_e(0)}{\psi_o(0)} = -\frac{\psi_o'(0)}{\psi_e'(0)} \tan^2 \frac{kd}{2} \quad (A5)$$

where all the functions are evaluated at  $x = d/2$ . Since  $\psi_e$  and  $\psi_o$  are functions of the energy,  $E$ , this equation determines the band energy  $E_k$ . Clearly  $E_{-k} = E_k$ . If  $E_0$  is the band energy at the zone centre ( $k = 0$ ), we must have either  $\psi_e'(d/2; E_0) = 0$  or  $\psi_o(d/2; E_0) = 0$ . The former defines what we shall refer to as an even-parity band, while the latter defines an odd-parity band. The small- $k$  behaviour of  $E$  is thus readily obtained from these properties. For example, for an odd-parity band we have

$$\psi_o(d/2; E) = -\psi_o(E - E_0) +$$

where  $\psi_o(d/2; E) = dE \left. \frac{d\psi_o}{dx} \right|_{x=E_0}$ . We then have

$$E_k = E_0 - \frac{\psi_o(0)}{\psi_o'(0)} \frac{dk}{2}^2 +$$

The coefficient of  $k^2$  defines the effective mass of the band. A similar result applies in the case of the even-parity bands.

Once the energy eigenvalue for a given  $k$  is known, the coefficients  $a$  and  $b$  are related by

$$\frac{b}{a} = i \frac{\psi_e(0)}{\psi_o(0)} \tan \frac{kd}{2}; \quad (A8)$$

For a given band,  $n$ , the ratio  $b/a$  is a continuous function of  $k$ . At  $k = 0$  we choose  $w_{k=0}(x)$  to be real and assume that it is a parity eigenstate. In this situation, we must have either  $b(k=0) = 0$  (even-parity bands) or  $a(k=0) = 0$  (odd-parity bands).

The normalization of  $w_k$  leads to the expressions

$$|a|^2 = \frac{1}{1 + \frac{d^2}{4}}; \quad |b|^2 = \frac{\frac{d^2}{4}}{1 + \frac{d^2}{4}} \quad (A9)$$

where

$$\tan \frac{kd}{2} = \frac{\psi_e(0)}{\psi_o(0)}; \quad (A10)$$

For an even-parity band  $\psi_o(0) \neq 0$ , so that  $a(k) \neq 0$  and  $b(k) \neq 0$ . In this case,

$$w_k(x) = e^{ikx} \left( \frac{\psi_e(x)}{1 + \frac{d^2}{4}} + i \frac{\psi_o(x)}{1 + \frac{d^2}{4}} \right); \quad (A11)$$

Since  $\tan \frac{kd}{2} = \frac{\psi_e(0)}{\psi_o(0)}$ , we see that  $w_k(-x) = w_k(x)$ . On the other hand, for an odd-parity band,  $\psi_e(0) \neq 0$ , and  $b(k) \neq 0$ . As a result, we have

$$w_k(x) = e^{ikx} \left( \frac{i \text{sgn}(k) \psi_e(x)}{1 + \frac{d^2}{4}} + \frac{\psi_o(x)}{1 + \frac{d^2}{4}} \right); \quad (A12)$$

which implies  $w_k(-x) = -w_k(x)$ . We have thus shown that the Bloch functions have the property

$$w_k(-x) = \mp w_k(x) \quad (A13)$$

where the positive (negative) sign corresponds to the even (odd) parity bands. Together with the conjugation property  $w_k(x) = w_k^*(x)$ , we have  $w_k(-x) = \mp w_k(x)$ .

For an even-parity band, the real and imaginary parts of  $w_k$  are

$$\begin{aligned} \text{Re } w_k(x) &= \frac{1}{1 + \frac{d^2}{4}} (\cos(kx) \psi_e(x) + \sin(kx) \psi_o(x)) \\ \text{Im } w_k(x) &= \frac{1}{1 + \frac{d^2}{4}} (-\sin(kx) \psi_e(x) + \cos(kx) \psi_o(x)); \end{aligned}$$

Thus the real part is an even function of  $x$  with the property  $\text{Re } w_k(x) = \text{Re } w_k(-x)$ , while the imaginary part is odd and  $\text{Im } w_k(x) = -\text{Im } w_k(-x)$ . The opposite is true of an odd-parity band. One can show for an arbitrary  $k$  in the lowest band that there is no net change in the phase  $\phi_k(x) = \tan^{-1} \frac{\text{Im } w_k(x)}{\text{Re } w_k(x)}$  as  $x$  varies between  $-d/2$  and  $d/2$ . We make use of this result in Sec. II.

The method described above cannot be used in three dimensions, but perturbation theory allows one to infer the same symmetry property. We write the Schrodinger equation for the Bloch function  $w_k(r)$  as

$$(\hat{h}_0 + V) w_k(r) = E_k w_k(r) \quad (A14)$$

where  $\hat{h}_0 = -\frac{\hbar^2}{2m} \nabla^2 + V(r)$  and  $V(r) = V(r - R)$ . The eigenfunctions of  $\hat{h}_0$ ,  $w_n(r)$ , with

eigenenergies  $E_n$ , are chosen to be parity eigenstates. The state  $w_k$  to first order in  $V$  is

$$w_k = w_n + \sum_{n^0} \frac{P_{n^0 n}}{E_n - E_{n^0}} w_{n^0} \quad (A15)$$

where  $P_{n^0 n}$  is the  $k = 0$  momentum matrix element defined in (70). Since this matrix element only couples states with opposite parity, we see that  $w_k(r) = w_k(r)$ , depending on the parity of the state  $n$ . Thus, the Bloch state exhibits the symmetry property to lowest order in  $k$ . It is evident that this argument can be extended to higher orders in perturbation theory.

#### APPENDIX B: DYNAMIC INSTABILITY IN THE WEAK POTENTIAL LIMIT

As pointed out by Wu and Niu [6], the boundary of the dynamically unstable region in the weak potential limit is given by the condition  $E(q) = E_+(q, G)$  where  $E(q)$  is given by (127). These energies are the eigenvalues of the Bogoliubov equations

$$\hat{B}_0 \begin{pmatrix} u \\ v \end{pmatrix} = E \begin{pmatrix} u \\ v \end{pmatrix} \quad (B1)$$

with

$$\hat{B}_0 = \begin{pmatrix} \frac{\hbar^2}{2m} \frac{d^2}{dz^2} + gn \frac{w_k^{(0)}}{gne^{2ikz}} & gn e^{2ikz} \\ gne^{2ikz} & \frac{\hbar^2}{2m} \frac{d^2}{dz^2} + gn \frac{w_k^{(0)}}{gne^{2ikz}} \end{pmatrix} \quad (B2)$$

The phonon mode of interest is

$$\begin{pmatrix} u_+ \\ v_+ \end{pmatrix} = \begin{pmatrix} a_+ e^{i(q-G+k)z} \\ b_+ e^{i(q-G-k)z} \end{pmatrix} \quad (B3)$$

with  $a_+ = \frac{q}{(\mu_q - E_+)} = 2E_+$  and  $b_+ = \frac{q}{(\mu_q - E_+)} = 2E_+$  where  $\mu_q - E_+ = \frac{\hbar^2}{2m} \frac{d^2}{dz^2} + gn$  and  $E_+ = E_+(q, G) = \frac{\hbar^2}{2m} \frac{d^2}{dz^2} + gn$ . The normalization of the mode is  $a_+^2 = b_+^2 = 1$ . The anti-phonon mode which is coupled to the phonon mode by a weak optical potential,  $V_{opt} = V_0 \cos(Gz)$ , is

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} a e^{i(q+k)z} \\ b e^{i(q-k)z} \end{pmatrix} \quad (B4)$$

with  $a = \frac{q}{(\mu_q - E_-)} = 2E_-$  and  $b = \frac{q}{(\mu_q - E_-)} = 2E_-$ , where  $E_- = E_-(q, G) = \frac{\hbar^2}{2m} \frac{d^2}{dz^2} + gn$ .

We note that in this case the mode has normalization  $a^2 = b^2 = 1$ .

The degeneracy of the phonon and anti-phonon modes ( $E(q) = E_+(q, G) = E_0$ ) suggests that we seek a solution of the Bogoliubov equations

$$\hat{B} \begin{pmatrix} u \\ v \end{pmatrix} = E \begin{pmatrix} u \\ v \end{pmatrix} \quad (B5)$$

in the form

$$\begin{pmatrix} u \\ v \end{pmatrix} = A \begin{pmatrix} u_+ \\ v_+ \end{pmatrix} + B \begin{pmatrix} u_- \\ v_- \end{pmatrix} \quad (B6)$$

Expanding the operator  $\hat{B}$  to first order in the optical potential, we have  $\hat{B} = \hat{B}_0 + \hat{B}_1$  with

$$\hat{B}_1 = \begin{pmatrix} V_{opt} + 2gn(w_1 + w_1^*) & 2gne^{2ikx}w_1 \\ 2gne^{-2ikx}w_1^* & V_{opt} + 2gn(w_1 + w_1^*) \end{pmatrix} \quad (B7)$$

Here we have written the condensate wave function as  $\psi_k(x) = \frac{1}{\sqrt{N}} e^{ikx} (1 + w_1 + w_1^*)$  where the first order correction is  $w_1(x) = \frac{1}{\sqrt{N}} e^{igx} + \frac{1}{\sqrt{N}} e^{-igx}$  with

$$= \frac{\hbar}{2} \frac{q \frac{1}{4} \frac{1}{\mu_q^{(0)} \mu_k^{(0)}}}{\mu_q^{(0)2} + 2gn \mu_q^{(0)} \mu_k^{(0)}} : \quad (B8)$$

Taking the inner product of (B5) with  $(u_+, v_+)$  and  $(u_-, v_-)$ , and noting the different normalizations of the two modes, we obtain the matrix equation

$$\begin{pmatrix} E_0 & E \\ E_0 + E & A \\ & B \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = 0; \quad (B9)$$

where the real coupling parameter is given by

$$= \frac{V_0}{2} + 2gn(a_+ + a_-)(a_+ a_- + b_+ b_-) + 2gn(a_+ b_- + a_- b_+) : \quad (B10)$$

A nontrivial solution to the matrix equation is obtained if

$$E = E_0 \quad ij : \quad (B11)$$

Thus, the line in the  $k$ - $q$  plane defined by (128) lies within the region of dynamical instability when  $V_0$  is finite. As emphasized by Wu and Niu [6], the dynamical instability in the weak potential limit arises from a resonant coupling between phonon and anti-phonon modes.

- (1998).
- [3] O. Morsch, J.H. Müller, M. Cristiani, D. Ciampini and E. Arimondo, Phys. Rev. Lett. 87, 140402 (2001).
  - [4] S. Burger, F.S. Cataliotti, C. Fort, F. Minardi, M. Inguscio, M.L. Chiofalo and M.P. Tosi, Phys. Rev. Lett. 86, 4447 (2001).
  - [5] K. Berg-Sørensen and K. Mølmer, Phys. Rev. A 58, 1480 (1998).
  - [6] B. Wu and Q. Niu, Phys. Rev. A 64, 061603(R) (2001).
  - [7] M. Machholm, C.J. Pethick, and H. Smith, Phys. Rev. A 67, 053613 (2003).
  - [8] J.C. Bronski, L.D. Carr, B. Deconinck, J.N. Kutz and K. Promislow, Phys. Rev. E 63, 036612 (2001).
  - [9] V.V. Konotop and M. Salerno, Phys. Rev. A 65, 021602 (2002).
  - [10] J. Javanainen, Phys. Rev. A 60, 4902 (1999).
  - [11] M.L. Chiofalo, M. Polini and M.P. Tosi, Euro. Phys. J. D 11, 371 (2000).
  - [12] A. Smerzi, A. Trombettoni, P.G. Kevrekidis and A.R. Bishop, Phys. Rev. Lett. 89, 170402 (2002).
  - [13] M. Kramers, L. Pitaevskii, and S. Stringari, Phys. Rev. Lett., 88, 180404 (2002).
  - [14] J. Callaway, Quantum Theory of the Solid State (Academic Press, New York, 1991).
  - [15] H. Pu, L.O. Baksmaty, W. Zhang, N.P. Bigelow and P. Møystre, Phys. Rev. A 67, 043605 (2003).
  - [16] C.M. Bender and S.A. Orszag, Advanced Mathematical Methods for Scientists and Engineers (McGraw-Hill, New York, 1978).
  - [17] C. Menotti, M. Kramers, L. Pitaevskii and S. Stringari, cond-mat/0212299.
  - [18] M. Kramers, C. Menotti, L. Pitaevskii and S. Stringari, cond-mat/0305300.
  - [19] M. Machholm, A. Nicolin, C.J. Pethick and H. Smith, cond-mat/0307183.
  - [20] D. Diakonov, L.M. Jensen, C.J. Pethick and H. Smith, Phys. Rev. A 66, 013604 (2002).
  - [21] D.A.W. Hutchinson, E. Zaremba and A. Griffin, Phys. Rev. Lett. 78, 1842 (1997).
  - [22] E. Zaremba, Phys. Rev. A 57, 518 (1998).
  - [23] A.D. Jackson, G.M. Kavoulakis, and C.J. Pethick, Phys. Rev. A 58, 2417 (1998).
  - [24] A. Trombettoni and A. Smerzi, Phys. Rev. Lett. 86, 2353 (2001).
  - [25] More precisely, we fix the direction of the phonon wave vector and order the expansion according to powers of the wave vector magnitude  $q$ .
  - [26] Alternatively, the hydrodynamic equations can be derived by taking variations of the action  $S = \int d^3r dt f(\mathbf{n}(\mathbf{r},t); \mathbf{h}(\mathbf{r},t)) + \int d\mathbf{r} \mathbf{h} \cdot \nabla \mathbf{n}$  with  $\mathbf{h} = \mathbf{r} \times \mathbf{v}$ .
  - [27] N.W. Ashcroft and N.D. Mermin, Solid State Physics, (Harcourt, 1976), pg. 765.
  - [28] B. Wu and Q. Niu, cond-mat/0306411.
  - [29] The other possibility of phonons with wave vectors  $q$  ( $q > 0$ ) and  $G - q$  gives a critical wave vector larger than (124).