

Magnetization plateau and quantum phase transitions in a spin-orbital model

Zu-Jian Ying^{1,2,3}, Angela Foerster², Xi-Wen Guan⁴, Bin Chen¹, Itzhak Roditi³

1. Hangzhou Teachers College, Hangzhou 310012, China

2. Instituto de Física da UFRGS, Av. Bento Gonçalves, 9500, Porto Alegre, 91501-970, Brasil

3. Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brasil

4. Department of Theoretical Physics, Research School of Physical Sciences and Engineering, and Centre for Mathematics and its Applications, Mathematical Sciences Institute, Australian National University, Canberra ACT 0200, Australia

(March 22, 2024)

A spin-orbital chain with different Lande g factors and one-ion anisotropy is studied in the context of the thermodynamical Bethe ansatz. It is found that there exists a magnetization plateau resulting from the different Lande g factors. Detailed phase diagram in the presence of an external magnetic field is presented both numerically and analytically. For some values of the anisotropy, the four-component system undergoes five consecutive quantum phase transitions when the magnetic field varies. We also study the magnetization in various cases, especially its behaviors in the vicinity of the critical points. For the SU(4) spin-orbital model, explicit analytical expressions for the critical fields are derived, with excellent accuracy compared with numerics.

Orbital degeneracy in electron systems leads to rich and novel magnetic phenomena in many transitional metal oxides [1]. Among them are the orbital ordering and orbital density wave, which have been observed experimentally in a family of manganites [2]. A tractable model to describe 2-fold orbital degenerate system is the SU(4) model [3], which has attracted much attention [3–9]. In the one-dimensional case the model is exactly solvable by Bethe ansatz (BA) [4]. An interesting question is to study the critical behavior of such a system in an external magnetic field, especially when different Lande g factors for spin and orbital sectors are involved. One may expect that the difference of g factors will bring about new physics as a result of the competition of the spin and orbital degrees of freedom. In Ref. [5], the authors studied the magnetic properties of the SU(4) model via BA, without taking different g factors into account, whereas numerical calculation was performed in Ref. [6] for the model with different g factors for up to 200 lattice sites. However, a full picture about the critical fields is still lacking. Another motivation is to see whether or not any magnetization plateau (MP), an interesting magnetic phenomenon, occurs in such a spin-orbital model. As is well known, antiferromagnetic chains with integer spin are gapful [10], whereas for half-integer spin there also exists a gapful phase with a MP in the presence of a large planar anisotropy [11]. Also fractional MP have been observed and can be explained by Shastry-Sutherland lattice [12]. But an MP arising from different Lande g factors has not been addressed yet.

Deviation from the SU(4) symmetry was considered by variation in the interaction parameters of neighbor sites [7,8], while another possible deviation may result from the one-ion interaction. Since many compounds are magnetically anisotropic in which the orbital angular momentum (OAM) may be constrained in some direction due to crystalline field, the angle between spin and OAM determines the spin-orbital coupling (SOC) energy. This kind

of one-ion SOC leads to magnetic anisotropy [13]. Under the influence of molecular field and an external field, the spin is parallel to the OAM, then the SOC energy depends on whether they are in the same directions. In such a case, $s_i^z \tau_i^z$ type of interaction describes well the SOC energy. Here we introduce such an SU(2) × SU(2) SOC interaction into the SU(4) model. A detailed investigation of the phase diagram is undertaken both numerically and analytically in the context of the thermodynamical Bethe ansatz (TBA). We find that the system exhibits an MP resulting from different g factors when the SOC is sufficiently strong. The critical behavior of the magnetization in the vicinities of the critical points is revealed. For certain values of Lande g factor, the model undergoes five consecutive quantum phase transitions when the external magnetic field varies. Further, the explicit analytic expressions for the critical fields for the SU(4) model are derived, with excellent accuracy compared to numerical results.

The model and TBA. Consider an L -site chain with the Hamiltonian

$$H = H_0 + H_z + M; \quad H_0 = \sum_{i,j} P_{ij} s_i^\dagger s_j; \quad H_z = \sum_i s_i^z \tau_i^z; \quad M = \sum_i g_s H_i s_i^z - \sum_i g_t H_i \tau_i^z; \quad (1)$$

where s and τ are spin-1/2 operators for spin and orbital sectors, and g_s and g_t denote, respectively, the Lande g factors. We assume $g_s > g_t$ throughout the paper. H_0 is the SU(4) model with $P_{ij} = (2s_i^\dagger s_j + 1)(2\tau_i^\dagger \tau_j + 1)$ exchanging the four site states $|s_i^\dagger \tau_i^\dagger\rangle$: $1 = |s_i^\dagger \tau_i^\dagger\rangle$, $2 = |s_i^\dagger \tau_i\rangle$, $3 = |s_i \tau_i^\dagger\rangle$, $4 = |s_i \tau_i\rangle$. The symmetry is broken into SU(2) × SU(2) by H_z and further into four U(1)'s by the external magnetic field H . The model can be solved exactly via BA approach. The BA equations are the same as the SU(4) model [4,14] under the periodic boundary conditions, with the energy eigenvalues given by

$$E = \sum_{i=1}^M a_i^{(1)} + \sum_{k=1}^4 E_k N_k; \quad (2)$$

where $a_n^{(1)} = 1/(2^n) n = (2^2 + n^2 = 4)$, and $E_1 = \frac{z=4}{g} H = 2$; $E_2 = \frac{z=4+g}{g} H = 2$, $E_3 = \frac{z=4}{g_+} H = 2$; $E_4 = \frac{z=4+g_+}{g_+} H = 2$, with $g = g_s - g_t$. N_k is the total site number in state k and $M^{(i)}$ ($i = 1; 2; 3$) is the rapidity number. For a certain choice of the basis order, which depends on whether or not the component is energetically favorable, the energy can be rewritten as $E = \sum_{i=1}^M g^{(i)} M^{(i)} + g^{(3)} M^{(3)}$, with the coefficients $g^{(i)}$ depending on the basis order. In the thermodynamical limit, the summation over the rapidity becomes integral. Following Refs. [15] and [16], one may obtain the ground state (GS) equations for the dressed energies $\epsilon^{(i)}$ ($i = 1; 2; 3$),

$$\epsilon^{(i)} = g^{(i)} a_2^{(i)} + a_1^{(i+1)} + a_1^{(i-1)}; \quad (3)$$

where $\epsilon^{(0)} = \epsilon^{(4)} = 0$ and the symbol $\otimes$ denotes the convolution. The dressed energies describe elementary excitations over the Fermi seas, i.e., the GS with all states with negative dressed energies filled. According to an energetics argument, we may divide the external field H into three regions: (I) $0 < H < H_{R1}$, (II) $H_{R1} < H < H_{R2}$, (III) $H_{R2} < H < 1$ with $H_{R1} = \frac{1}{2} \ln 2 (2g_s)$; $H_{R2} = \frac{1}{2} \ln 2 (2g_t)$. For $z > 0$; the corresponding basis order are: (I) $(1, 2; 3; 4)^T$, (II) $(1, 3; 2; 4)^T$, (III) $(3, 1; 2; 4)^T$; for $z < 0$: (I) $(3; 4; 1, 2)^T$, (II) $(3; 1, 4; 2)^T$, (III) the same as $z > 0$. These five basis orders provide a full description of the phase diagram of the system.

Plateau. The competition between the anisotropy parameter z and the magnetic field H results in a novel quantum phase diagram. In the absence of the magnetic field, it is easy to find that the states 3 and 4 are gapful for $z > \frac{c}{z} = 4 \ln 2$. Whereas for $z < \frac{c}{z}$, the components 1 and 2 are gapful. Therefore, the GS is in an $su(2)$ spin-orbital liquid state in strong anisotropy regime in the absence of the field. However, the presence of the magnetic field completely splits all four components energetically. The magnetization $M^z = g_s s^z + g_t z^z$ increases from zero. For large positive z , the field brings the component 3 closer to the GS, while the component 2 gradually gets out of the GS. Certainly, if the field reaches the first critical field where the component 3 has not yet involved in the GS, a quantum phase transition from the spin-orbital liquid phase to a ferromagnetic phase occurs. Thus a magnetization plateau opens with a constant magnetization $M^z = g_- = 2$. Nevertheless, this plateau will disappear when the field is strong enough $H > H_{c2}^P$. In such a case, the component 3 becomes involved in the GS. The critical field H_{c2}^P indicates a quantum phase transition from the ferromagnetic GS into a spin-orbital liquid phase. If the field continues

to increase, the spin and orbital sectors become fully-polarized at the third critical point H_{c3}^P . From the TBA equations (3), we get the exact expressions for the critical fields

$$H_{c1}^P = \frac{4}{g}; \quad H_{c2}^P = \frac{z=2}{g_t} \frac{4}{g_t}; \quad H_{c3}^P = \frac{z=2+4}{g_t}; \quad (4)$$

Notice that the plateau opens only if $z > \frac{P}{z} = 8g_s = g$ and $0 < g_t < g_s$. If the g factors are the same, the plateau disappears because the components 1 and 2 remain degenerate in the field. The critical behavior of the magnetization in the vicinities of the critical points may be summarized as follows

$$HM^z = HM^z_{ic} + k_M H^{\frac{1}{2}}; \quad (5)$$

where $HM^z_{ic} = g_- = 2$ are, respectively, plateau and saturation magnetizations. H is the small deviation from the critical points and $\alpha = 1$ depending on M^z is increasing or decreasing. $k_M = g^{3/2} = g$ near H_{c1}^P and $k_M = g_t^{3/2} = g$ near H_{c2}^P and H_{c3}^P . The coefficients g and g_t in (4) and (5) can be easily understood, since only 1 and 2 exist in the GS before H_{c1}^P is reached, the differences of their field energy and magnetization are $g H$ and g respectively. While for $H_{c2}^P < H < H_{c3}^P$, only 1 and 3 compete in the GS, since 2 already gets out before 3 enters the GS. 1 and 3 differ in field energy by $g_t H$ and in magnetization by g_t . In Fig. 1A we plot the magnetization curves for different values of the parameter z , including a plateau case.

The phase diagram: numerical and analytical. Here we present a detailed analysis of the GS phase diagram both numerically and analytically. In Fig. 2, we plot the phase diagram with respect to the parameters z and H for fixed values of g_s and g_t ($g_s = 2.0$ and $g_t = 1.0$). For convenience, we refer to the GS with i components as i -state GS. Then for the phase transition between 3-state and 2-state GS, the critical fields follow from the Wiener-Hopf method [17], which is valid for large Fermi boundaries. Explicitly, we have

$$\begin{aligned} H_c^{PC+} &= (z - \frac{c}{z}) g_+^{-1} - g_-^2 g_+^{-3} (z - \frac{c}{z})^2; \\ H_c^{QP} &= \frac{\frac{c}{z} + \frac{z}{2}}{2g_s - g_t} + \frac{g_-^2 (z - \frac{K}{z})^2}{(2g_s - g_t)^3}; \\ H_c^{QF} &= \frac{\frac{c}{z} - \frac{z}{2}}{2g_s + g_t} + \frac{[g_t \frac{c}{z} - (g_s + g_t) \frac{z}{2}]^2}{(2g_s + g_t)^3}; \\ H_c^{FC} &= 2g_+^{-2} g_-^2 [1 - \frac{1}{4} g_-^2 g_+^{-2} (z + \frac{c}{z})^2]; \quad (6) \end{aligned}$$

where $1 = 1/(2^2)$ and $\frac{K}{z} = 2g_t H^K = g_t = g_- \frac{c}{z}$ which corresponds to infinite Fermi boundaries. Point Q is determined by $\frac{Q}{z} = 2g_t H^Q$, with $H^Q = 2 \ln 2 - g_s + 4 - 2 \ln 2 g_t^2 = g_s^3$. Near P or F the above analytic results deviate due to small Fermi boundaries. But in this case, an analysis may be carried out in terms of expansion of small Fermi boundaries. This leads us to $H_c^{PK}; H_c^{PC+} =$

$\frac{z}{2g_s} (g_+ = 2)^{\frac{3}{2}} = (\frac{5}{g_s^2}) (\frac{P}{z} z)^{\frac{3}{2}}$. Similarly, near F, we have $H_{C_+}^{FQ}, H_{C_+}^{FC} = \frac{z}{2g_s} (g_+ = 2)^{\frac{3}{2}} = (\frac{5}{g_s^2}) (\frac{F}{z} z)^{\frac{3}{2}}$.

For $j_z j < \frac{C}{z}$, the GS involves all the four components, the magnetic field first brings about a phase transition from a 4-component liquid to a 3-component liquid at the phase boundary $C_+ Q C_-$. Here one of the four components, which is energetically unfavorable, completely gets out of the GS, then the corresponding Fermi sea disappears. So the critical field only involves two Fermi points B_1 and B_2 . At point C_+ , the component 2 is degenerate with the component 1 which is the most energetically favorable, the Fermi point B_1 lies at infinity. Increasing H along $C_+ M N Q$ drives 2 away from 1 , so 2 becomes less energetically favorable in the GS. Therefore the first Fermi sea shrinks, i.e., B_1 decreases from infinity. Beyond M point, both the increase of H and the decrease of z make 3 sink below the 2 which is rising, the energy difference between 3 and 1 begins to dominate over B_1 . As 3 is drawing near 1 , the first Fermi sea becomes broadened again with an increase of B_1 . After point N, 3 sinks beyond 1 to be the lowest state, 1 becomes less favorable in the GS. The first Fermi sea shrinks again, B_1 begins to decrease along NQ from the infinity at N. A similar analysis is applicable to B_2 , which is mainly influenced by the energy difference between the second and third components in energy levels. B_2 rises from zero at C_+ to infinity at M and decreases along MNQ to zero at Q. In the respective sections of $C_+ M N Q$ the critical fields take the form

$$\begin{aligned} H_{C_+ M N Q}^{C_+} &= a_{1;1} g_+ + H_{B_2}^{C_+ M N Q} + w H_{B_1}^{C_+ M N Q} \\ &= H_1^{C_+} + \frac{2}{a_{1;3}} g_+ + w \frac{2}{a_0} (g_+ a_0 - g_t a_{1;3})^2 = g_+^3 \\ &= H_1^{N} + \frac{2}{a_0} [(g_+ a_0 + g_s a_{1;3})^2 + w (g_+ a_0 - g_t a_{1;3})^2] = g_+^3 \\ &= H_1^{NQ} + \frac{2}{a_0} (g_+ a_0 + g_s a_{1;3})^2 = g_+^3 + w \frac{2}{a_{1;3}} g_+; \end{aligned}$$

where $a_{1;1} = a_0 \frac{m}{2} z, a_0 = \frac{P}{2} \frac{3}{2} \ln 3, \frac{2}{a_0} = \frac{3}{16} z$, and $w = 2/3$. In each case, the first term comes from infinite B_1 and B_2 , the second term is correction from finite but large B_2 , and the third term is the leading correction from the larger B_1 . For $C_+ M N$ near point M the B_2 and B_1 terms need to be exchanged since B_2 becomes larger. Along NQ B_1 is always larger than B_2 . The location of $B_1 = B_2$ in $C_+ M N$ may be estimated by $H_{B_1} = H_{B_2}$, which gives $z = 2a_0(2g_s - g_t) = (3g_s)$ for $C_+ M$ and $z = \frac{2}{3}a_0$ for $M N$. This coincides well with numerics, e.g., for $M C_+$ and $g_s = 2.0; g_t = 1.0$, the analytic result is $z = 1.073$ whereas the numerical one is 1.042.

Similarly, for $C_- V Q$, the term resulting from the infinite Fermi boundaries is $H_1 = a_{1;1} g_-$, the correction terms are $H_{B_2}^{VC} = \frac{2}{a_{1;3}} g_-$, $H_{B_1}^{VC} = H_{B_1}^{VQ} = \frac{2}{a_0} [g_+ a_0 + g_t a_{1;3}]^2 = g_-^3$, and $H_{B_2}^{VQ} = \frac{2}{a_0} [g_+ a_0 + g_s a_{1;3}]^2 = g_-^3$, respectively. These expressions are not valid for $g_s = g_t$ due to small Fermi boundaries.

When the system is fully-polarized, only 3 exists in the GS, while the other components are all gapful, with a gap $\Delta = \min E_i - E_3, i = 1, 2; 4g$. This gap is closed if $H < H_f$, with the fully-polarized critical point

$$H_f = \max (z = 2 + 4) g_t^{-1}; 4g_+^{-1}; \quad (7)$$

This expression is exact and valid for all z . When $\frac{F}{z} = 8g_s = g_+$, the strong negative anisotropy makes 1 and 2 too far away from 4 . Before the field brings them close enough to get involved in the GS, the component 4 has been all pumped out by the field from the GS at critical point $H_f = 4g_+$. For all $z = \frac{F}{z}$, the magnetization is the same as shown by curve a in Fig.1A.

The analytic results are compared with the numerics in Fig.2, with very satisfactory accuracy.

Five consecutive H_c phase transitions. The competition of anisotropy z and the external field H also leads to an unusual magnetic phenomenon. For some fixed values of z, g_s and g_t , the system undergoes five consecutive quantum phase transitions when H varies, although the 4-component model usually has at most three consecutive phase transitions. A strong negative anisotropy z makes 4 energetically quite favorable. The field H expels 2 first from the 4-component GS. However, before H overwhelms the influence of z on 4 , further increase of H will make 2 closer to 4 and draw it back into the GS. This process brings about the first two phase transitions. Then H plays a dominant role, it begins to bring out $4, 2$ and 1 from the GS one by one. This results in other three consecutive phase transitions. The variation of the state component numbers in the GS is: $4! 3! 4! 3! 2! 1$. This five H_c 's case exists for all $0 < g_t < g_s$ and becomes more visible when g_t is larger. One case is marked by black dots in Fig.3B for $g_s = 2.0$ and $g_t = 1.8$. Another possible case for five H_c transitions to occur is the GS composed of 3 and 4 . The field brings 1 into the GS first. As the difference of g_t and g_s is getting smaller, 2 has closer energy to 1 . Further increase of H brings 2 into the GS before 4 completely gets out. The component number changes in such a way: $2! 3! 4! 3! 2! 1$. This case occurs when the point Q is below C_- in Fig.2, approximately requiring $1 > g_t = g_s > 1 - 2 \ln 2 g_t^3 = (\frac{2}{3} g_s^3)$. Four consecutive H_c transitions take place for $z = \frac{C}{z}$. In Fig.1B and C, we plot the magnetization curves which display five consecutive H_c phase transitions.

The SU(4) model. If we set the anisotropy parameter z to be zero, the model reduces to the SU(4) model with different Lande g factors in the spin and orbital sectors. In this special case, the above results for the critical fields give rise to

$$\begin{aligned} H_{c1}^{SU(4)} &= a_0 g_+^{-1} + \frac{2}{a_0^2} (2g_s - g_t)^2 g_+^{-3} + w \frac{2}{a_0^2} g_+^{-1}; \\ H_{c2}^{SU(4)} &= \frac{C}{z} (2g_s - g_t)^{-1} + \frac{1}{a_0} (g_t - \frac{C}{z})^2 (2g_s - g_t)^{-3}; \quad (8) \end{aligned}$$

$$H_{c3}^{SU(4)} = 4g_t^{-1}:$$

We compare the above analytic results with TBA numerical ones in the Fig. 3A, which shows an excellent accuracy for most values of g_t . Notice that the deviations occur only for less-physical values of g_t due to small Fermi points. For $g_s = 2.0$ and $g_t = 1.0$, the discrepancies of $H_{c1}^{SU(4)}$ and $H_{c2}^{SU(4)}$ (TBA numerics: 0.3695, 0.9415; analytic: 0.3697, 0.9386 (200 sites results [6] for comparison: 0.31, 0.93)) are, respectively, 0.05% and 0.3%. $H_{c3}^{SU(4)}$ is exact. The magnetization of the SU(4) model is shown in Fig. 1A, which also coincides with the numerical result for 200 sites [6].

We thank Huan-Qiang Zhou, You-Quan Li and M.T. Batchelor for discussions and comments. ZJY thanks FAPERGS and FAPERJ for partial support. AF thanks FAPERGS and CNPq. XWG thanks the Australian Research Council for support. BC thanks NSF (Natural Science Foundation) of Zhejiang, China (RC02068) and NSF of China (10274070). IR thanks PRONEX and CNPq.

-
- [1] Y. Tokuro and N. Nagaosa, *Science* 288, 462 (2000).
 - [2] E. Saitoh et al, *Nature (London)* 410, 180 (2001).
 - [3] Y. Q. Li et al, *Phys. Rev. Lett.* 81, 3527 (1998).
 - [4] Y. Q. Li et al, *Phys. Rev. B* 60, 12781 (1999).
 - [5] Y. Yamashita et al, *Phys. Rev. B* 61, 4012 (2000).
 - [6] S. J. Gu and Y. Q. Li, *Phys. Rev. B* 66, 092404 (2002).
 - [7] F. M.ila et al, *Phys. Rev. Lett.* 82, 3697 (1999).
 - [8] P. A zaria et al, *Phys. Rev. Lett.* 83, 624 (1999).
 - [9] B. Frischmuth et al, *Phys. Rev. Lett.* 82, 835 (1999).
 - [10] F. D. M. Haldane, *Phys. Lett. A* 93, 464 (1983).
 - [11] M. Oshikawa et al, *Phys. Rev. Lett.* 78, 1984 (1997).
 - [12] K. Kodama et al, *Science* 298, 395 (2002); K. Onizuka et al, *J. Phys. Soc. Jpn.* 69, 1016 (2000); G. Misguich et al, *Phys. Rev. Lett.* 87, 097203 (2001); B. S. Shastri and B. Sutherland, *Physica B* 108, 1069 (1981).
 - [13] J. C. Slonczewski, *Phys. Rev.* 110, 1341 (1958).
 - [14] B. Sutherland, *Phys. Rev. B* 12, 3795 (1975).
 - [15] M. Takahashi, *Prog. Theor. Phys.* 46, 401 (1971).
 - [16] C. N. Yang and C. P. Yang, *J. Math. Phys.* 10, 1115 (1969).
 - [17] M. G. Krein *Usp. Mat. Nauk* 13, 3 (1958).




FIG. 1. (A) Typical magnetization behaviors for fixed values of Lande g factors $g_s = 2.0$ and $g_t = 1.0$. The numbers in brackets indicates values of the anisotropy parameter z . The dotted line e denotes the magnetization for the SU(4) model. Curve j exhibits a magnetization plateau. (B) Magnetization corresponding to consecutive H_c quantum phase transitions for $g_s = 2.0$, $g_t = 1.9$, and $z = 3.0$. Here the variation of the state component numbers is 2! 3! 4! 3! 2! 1. The number i labels the state i , e.g., the state components in the phase 123 are $i_1 i_2 i_3$ in which i_1 is the most energetically favorable whereas i_3 is the least energetically favorable. The phase variations between H_{c2} and H_{c3} are 3412! 3142! 3124. The dotted line for comparison is an extension of the phase 34 by assuming the components unchanged. (C) Magnetization for consecutive H_c phase transitions for $g_s = 2.0$ and $g_t = 1.0$, with $z = 1.4$. $H_{c1} = 0.531$ (transition 3142! 214) and $H_{c5} = 3.33$ (31! 3) are relatively far away. The variation of the component numbers in the phase transitions is 4! 3! 4! 3! 2! 1.




FIG. 2. Phase diagram for fixed values of Lande g factors $g_s = 2.0$ and $g_t = 1.0$. Inside $C_+ Q C_-$, the 4-state phases are 1234 (I_+), 1324 (II_+), 3124 (III), 3142 (II_-), and 3412 (I_-). The discrepancy of the analytic curves from the numerical ones is not visible for most regions of $C_+ PQ$ and $C_+ MNQ$ (with typical differences within 1.0%, and 0.1%, respectively). There is less accuracy in regions I_- and II_- for the analytic results due to larger driving terms and smaller Fermi boundaries. Magnetization plateaux and fully-polarized cases are exact.




FIG. 3. (A) Comparison of the analytic and the numerical results for the three critical fields of the SU(4) model with respect to the orbital g factor g_t for a fixed value of the spin g factor $g_s = 2.0$. The numbers label the state components as in Fig 2, (B) Phase diagram in the Δ_z vs magnetic field H plane for different values of g_t (1.8, 1.3, 0.6, 0).