## A re the Tails of Percolation Thresholds G aussians ?

P M C. de O liveira<sup>1;2</sup>, R A. Nobrega<sup>2</sup> and D. Stau er<sup>1;3</sup>.

- Laboratoire de Physique et M ecanique des M ilieux H eterogenes E cole Superieure de Physique et de Chim ie Industrielles
  nue Vauquelin, 75231 Paris Cedex 05, France
- Instituto de F sica Universidade Federal F lum inense av. Litorânea s/n, Boa V iagem, Niteroi, Brasil 24210-340
  Institute for Theoretical Diversion
- Institute for Theoretical Physics Cologne University D-50923 Koln, Euroland

emails: pm col ifu br; rafaellal ifu br; stau erl thp unikoeln de.

PACS numbers: 02.70.-c, 05.10 Ln, 64.60 Ak, 05.70 Jk

A bstract: The probability distribution of percolation thresholds in nite lattices were not believed to follow a norm al Gaussian behaviour. With increasing computer power and more e cient simulational techniques, this belief turned to a stretched exponential behaviour, instead. Here, based on a further improvement of M onte C arb data, we show evidences that this question is not yet answered at all.

In reference [1], the percolation on a N -site square lattice is treated with high numerical accuracy. Indeed, the best known estimate for the critical threshold,  $p_c = 0.59274621$  (13), comes from this work. In order to study this kind of problems, the authors follow a very fruitful M onte C arlo approach which allows one to obtain continuous functions of p, the concentration of occupied sites, namely the canonical-like average

$$R(p) = \sum_{n}^{X} C_{N}^{n} p^{n} (1 p)^{N n} R_{n} ; \qquad (1)$$

of som e quantity R. Here, R<sub>n</sub> is a uniform average over all con gurations with just n occupied sites, i.e. a microcanonical-like average.  $C_N^n$  are simple binom ial factors. By lling the initially empty lattice, site by site at random, and repeating this process m any times, one is able to get the discrete set of microcanonical averages R<sub>n</sub> accumulated into an n-histogram, over the entire




Figure 1: Spanning probability function for a xed lattice size L (solid line). For larger and larger sizes (dotted lines), this function approaches a step. By xing som e value r at the vertical axis, one can nd a sequence of values p<sub>L</sub> (r)

at the horizontal axis approaching the critical threshold  $p_c$ , for increasing lattice sizes.

range, n = 0, 1, 2 ::: N. From this set of numbers, the determ ination of the continuous p-function R (p) is straightforward.

In particular the authors of [1] x attention at the horizontal wrapping probability  $R_L$  (p) around a L L torus, i.e. a square lattice with periodic boundary conditions on both directions. In the therm odynam ic limit, this function approaches a step:  $R_1$  (p) = 0 below the critical threshold  $p_c$ , and  $R_1$  (p) = 1 above  $p_c$ . For nite sizes,  $R_L$  (p) presents a sigm oid aspect similar to gure 1. A good approach to  $p_c$  is to choose some xed value r, and solve the equation  $R_L$  (p) = r, getting the root p shown at the horizontal axis. Here, one can appreciate the advantage of knowing  $R_L$  (p) as a continuous function of p. K exping the same value r and repeating this task for a series of increasing lattice sizes (dotted lines), one gets a series of roots  $p_{L_1}$  (r),  $p_{L_2}$  (r),  $p_{L_3}$  (r) :::which converges to the desired threshold  $p_c$  in the therm odynam ic limit.

The above reasoning is valid no matter which is the xed value for r one chooses. However, for the very particular choice r = 0.521058290, a universal probability exactly known through conform al invariance argu-

m ents [2], the convergence becom es fast, i.e. the root p(L) di ers from  $p_c$  as  $L^{2 l=} = L^{2:75}$ , where = 4=3 is the correlation length critical exponent. The above quoted accurate value for  $p_c$  was obtained in this way. For details, see [1] and references therein.

Reference [3] proposes the mathematical form

$$p_{L}(r) = p_{c} + \frac{1}{L^{1=}}^{h} A_{0}(r) + \frac{A_{1}(r)}{L} + \frac{A_{2}(r)}{L^{2}} + \cdots$$
 (2)

for estimators  $p_L$  obtained from quantities like R (p). The option for the wrapping probabilities around the torus and the convenient choice of P inson's number r = r lead to vanishing values for the two rst term s A<sub>0</sub> (r) = A<sub>1</sub> (r) = 0, a lucky coincidence which accelerates very much the convergence.

The quantity  $R_L$  (p) is obtained, as quoted before, by lling up the initially empty lattice site by site, at random . C lusters of neighbouring occupied sites grow . As soon as the horizontal wrapping along the torus is set, one books the corresponding value of n, the number of occupied sites so far, and stops the process. For that particular sample, the wrapping probability  $R_L$  is a step function, i.e.  $R_L = 0$  below n and  $R_L = 1$  above. The same routine is repeated m any times, in order to have a probability distribution for n. The various step functions are then superimposed to get the m icrocanonical averages  $R_n$  in equation (1), stored in an n-histogram . Finally, the continuous canonical average  $R_L$  (p) can be calculated for any value of p.

Each process of lling-up the lattice (one sam ple) yields a single value n for the statistics, i.e. just one more entry on the n-histogram. In [3], we decided to improve this point, by changing the de nition from wrapping to spanning probability, gure 1. We x two parallel horizontal lines on the L L torus, separated by a distance of L=2, for instance lines i = 1 and i = 1 + L=2. The measured quantity is now the probability of having these two lines connected by the same cluster of neighbouring occupied sites, instead of the wrapping probability along the whole torus. The advantage is that we can measure the same thing for lines i = 2 and i = 2 + L=2, also for lines i = 3 and i = 3 + L=2, and so on. Moreover, also vertical parallel lines can be included into this counting. At the end, from a single sam ple we store L new entries into our n-histogram, instead of just one more entry. Note that this advantage even increases for larger and larger lattices.

W ithin the same computational e ort, our approach allows the test of larger lattices. Because of that, we were able to con m the validity of equation (2) with high precision, by sampling 27 di erent lattice sizes from L = 18 up to L = 1594, a 8-thousand factor in number of sites. On the other hand, our de nition does not allow the chance of vanishing both A<sub>0</sub> (r) and A<sub>1</sub> (r) at once. We can have only A<sub>0</sub> (r) = 0, for a particular universal probability r = 0.984786 (11) num erically determ ined within the same work [3]. Independently, Cardy [4] tried to determ ine it by conform al invariance arguments, however, in looking for con gurations which link two parallel lines, he was forced to disregard con gurations which also wrap along the other direction. As a result of using larger lattices but a slower convergence rate of L<sup>11=</sup> = L<sup>1:75</sup>, we get the same gure  $p_c = 0.59274621$  (33) [3] as in [1], but within a 3 times larger error bar.




Figure 2: Test of equations (3) or (4), for tails on the left of gure 1. The ve continuous curves correspond to L = 1594, 1354, 1126, 958 and 802, from left to right. In each case we sam pled 4 m illion lattice-lling processes, which corresponds to 6, 5, 5, 4, and 3  $10^9$  entries in each n-histogram, respectively. The statistics is in proved by a factor over than 1000, com pared with [1] for equivalent lattice sizes. The dashed lines show the alternative slopes 2 (right) or 4/3 (left). In the authors' opinion, no de nitive conclusion is possible.

The non-G aussian behaviour of the nite-lattice-threshold distribution near the in nite-lattice critical point is already stablished [5]. Here, we pro t from the same simulational data in order to investigate the distribution tails, far from the critical point. Which is the mathematical form of the tail




Figure 3: A lternative test of equations (3) or (4), for the same lattice sizes L = 1594, 1354, 1126, 958 and 802, from left to right.

observed in gure 1, below the root p? One possible answer is a simple Gaussian form [6,7]

$$R_{L}(p) \exp[K(p p_{c})^{2}]$$
: (3)

A nother alternative is a stretched exponential [8, 9, 10]

$$R_{L}(p) = \exp[C(p_{c} p)];$$
 (4)

where the strict equality (for large L and far from  $p_c$ ) is a consequence of the periodic boundary condition [1], which holds for our data. Pro ting from this

strict equality, one can test equation (4) by constructing a plot of  $\ln[\ln(R)]$  against  $\ln(p_c p)$ . This was done in [1], and we repeat the same for our data, in gure 2. Note that our range for  $\ln(p_c p)$  (up to 3:4 for L 10<sup>3</sup>) is larger than in reference [1] (up to 4:6 for the same size). This means that we are testing more deeply the distribution tails, thanks to our trick of sampling L new entries for each run. Even so, the conclusion in favour of either equation (3) or (4) is by no means obvious. Note a further di culty in what concerns equation (3), because the leading multiplicative constant in front of the exponential is not necessarily 1.

A nother, perhaps better way to address the same question is by plotting ln (R) twice, against (p  $p_c$ )<sup>2</sup> and ( $p_c$  p)<sup>4=3</sup>. Figure 3 shows the result for our data. Note that our range for ln (R) (down to 16) doubles the one presented in [1]. The would-be G aussian case (up) presents clear positive curvatures, whereas the would-be stretched exponential (down) presents negative curvatures although not so pronounced. The exponent 4=3 seem s to t better, but one cannot extract a clear conclusion from these data.

Stillm one unde ned is the situation of the other tails on the right of gune 1, above  $p_c$ . In this case (not shown), our accuracy limit for ln (1 R) (down to 16) is reached much closer to  $p_c$  than the case shown in gures 2 and 3, below  $p_c$ .

C oncluding, we present new M onte C arlo data concerning the distribution probability of percolation thresholds on a nite square lattice. W e address the question of the m athem atical form of the distribution tails, equation (3) against (4). Even considering that our statistics is over than 1000 times larger than previous works, no de nitive conclusion can be extracted from our data, in what concerns the asymptotic tail exponent.

A know ledgem ents: This work is partially supported by B razilian agencies FAPERJ and CNPq.

## References

- M E J. Newman and R M. Zi, Phys. Rev. E 64, 016706 (2001); Phys. Rev. Lett. 85, 4104 (2000).
- [2] H.T. Pinson, J. Stat. Phys. 75, 1167 (1994).

- [3] P.M. C. de O liveira, R.A. Nobrega and D. Stau er, Braz. J. Phys. 33, 616 (2003), also in www.lanlgov.cond-m.at/0308525.
- [4] J.Cardy, J.Phys. A 35, L565 (2002).
- [5] This point was not yet realized in D. Stau er and A. Aharony, Introduction to Percolation Theory, 2<sup>nd</sup> edition, Taylor and Francis, London (1992), but was corrected in the 1994 printing.
- [6] M E. Levinshtein, B.I. Shklovskii, M S. Shur and A L. Efros, Zh. Eksp. Teor. Fiz. 69, 386 (1975) [Sov. Phys. JEPT, 42, 197 (1976)].
- [7] F.Wester, Int. J.M od. Phys. C 11, 843 (2000).
- [8] L.Berlyand and J.W ehr, J. Phys. A 28, 7127 (1995).
- [9] U.Haas, Physica A 215, 247 (1995).
- [10] J.P.Hoviand A.Aharony, Phys. Rev. E 53, 235 (1996).