D irected cycles and related structures in random graphs: II | D ynam ic properties

Valm ir C. Barbosa^{a;} Raul Donangelo^b Sergio R. Souza^b

^aProgram a de Engenharia de Sistem as e Computação, COPPE, Universidade Federal do Rio de Janeiro, C.P. 68511, 21941-972 Rio de Janeiro, Brazil

^bInstituto de F sica, Universidade Federal do Rio de Janeiro, C.P. 68528, 21941–972 Rio de Janeiro, Brazil

A bstract

W e study directed random graphs (random graphs whose edges are directed) as they evolve in discrete time by the addition of nodes and edges. For two distinct evolution strategies, one that forces the graph to a condition of near acyclicity at all times and another that allows the appearance of nontrivial directed cycles, we provide analytic and simulation results related to the distributions of degrees. W ithin the latter strategy, in particular, we investigate the appearance and behavior of the strong components that were our subject in the rst part of this study.

Key words: Random networks, Directed random networks, Evolving networks, Strong components PACS: 05.50.+ q, 89.75.k, 89.75 Fb, 89.75 H c

1 Introduction

In the st part of this study [1], we considered random digraphs on a xed set of n nodes with Poisson-distributed in-and out-degrees. For a wide range of values for z (the mean in- or out-degree of a node), we investigated the behavior of the strong components of the digraph, which essentially are maxim all subgraphs whose nodes can all be reached from one another by following

Corresponding author.

Em ailaddresses: valmir@cos.ufrj.br (Valm ir C.Barbosa),

donangel@if.ufrj.br (RaulDonangelo), srsouza@if.ufrj.br (Sergio R. Souza).

the directions of the edges. We contributed analytic and simulation results to the special cases of cycle components and knots. More special cally, we demonstrated that cycle components are concentrated near z = 1 and necessarily small, encompassing a number of nodes proportional to $\ln n$, while knots tend to occur only sparsely as size-one components for very small values of z, then become practically absent as z is increased, then nally occur again as a single knot that encompasses nearly all n nodes for larger values of z. The latter transition is sharp and happens roughly at $z = \ln (2n)$.

In this second part, our aim is to study the behavior of strong components when the digraph is no longer static, but rather evolves in time. This type of study was also pioneered by Erd&s and Renyi [2], who considered undirected graphs on a xed set of nodes that progressively become more and more interconnected by the random addition of edges between pairs of nodes, thus leading to a Poisson distribution of node degrees. Recently, though, in an attempt to model the networks that occur in m any areas of interest, considerable e ort has been directed towards studying evolution scenarios in which nodes and edges may enter and leave the network continually. The reader is referred to the surveys in [3] for details related to several areas. One of the surveys [4] highlights the mathematics of a variety of such models (e.g., [5]), many of which exhibit the power-law (as opposed to Poisson) distribution of degrees that is characteristic of networks that grow by some sort of preferential (as opposed to random) attachment of nodes to each other.

O ur focus is on the study of digraphs that evolve by the addition of new nodes, as well as new edges, in discrete time. Also, although we do touch the issue of preferential attachments brie y at one point, for the most part we follow [1] and concentrate on the case of random connections. Even though some of the models discussed in the literature (cf. [3]) can be claimed to include digraphs at least inherently we believe our emphasis on random connections in the evolution of digraphs covers new ground.

The following notation is common to all the remaining sections. For t 0 an integer, we let D_t denote the digraph at time t and N_t , with $n_t = N_t j$, its node set. D_0 is assumed to have no nodes, so $N_0 = j$ and $n_0 = 0$. Unlike the static case [1], it is now necessary to consider in- and out-degrees separately. For node i, we let d_t^+ (i) denote its in-degree at time t and d_t (i) its out-degree, with expected values z_t^+ (i) and z_t (i), respectively.

The paper is organized as follows. In Section 2 we consider evolution scenarios in which the deployment of edges disallows every nontrivial directed cycle. In Section 3 we modify the deployment rule in order to allow the appearance of arbitrary directed cycles, and with them nontrivial strong components. Conclusions are given in Section 4.

2 N early acyclic evolution

2.1 Analytic results

In the rst evolution scenario that we consider, at every time step t > 0 a new node is added to the digraph, so we have $n_t = t.W$ e assume that nodes are numbered consecutively from 1 as they enter the digraph, so node i is the node added at time t = i.W hen a node enters the digraph, a random number of edges is also added, all of them directed away from the new ly added node. Except for the possibility of self-bops (an edge leading from a node to itself is called a self-bop), the digraph is at all times acyclic (i.e., has no directed cycles spanning m ore than one node) and all its strong components have one single node.

For some xed z, we start by considering the case in which an edge is deployed from node i to each of nodes 1;:::; i independently with probability z=i (provided, of course, that z i; for z > i, it must be assumed that i is connected to all of 1;:::; i). In this case, for su ciently large i and t i, the out-degree d_t (i) is clearly Poisson-distributed with mean z_t (i) = z.

As for the in-degree d_t^+ (i), its expected value z_t^+ (i) is, for su ciently large i and t i, given by

$$z_{t}^{+}(i) = \frac{x^{t}}{u=i} \frac{z}{u} = z(H_{t} - H_{i-1});$$
(1)

where $H_m = {P_{u=1}}^{P_{m}} 1=u$ is the mth harmonic number, given for large m by $H_m = \ln m + ,$ with denoting Euler's constant [6]. Thus, for i 0,

$$z_{t}^{+}$$
 (i) $\ln \frac{t}{i}^{z}$: (2)

In order to discover how d_t^+ (i) is distributed, let $P_t(i;k)$ denote the probability that d_t^+ (i) = k fort i and k 0. If t i + 1 < k, then clearly P(i;k) = 0. O therwise, for k t i and z i;t, we have

$$P_{t}(i;k) = \frac{P_{t}(i;0)}{k!} z_{t}^{+}(i)^{i_{k}}; \qquad (3)$$

as we dem onstrate in Appendix A.P_t (i; 0) can now be approximated from (3)

by sum m ing $P_t(i;k)$ over the appropriate range of k:

1
$$P_{t}(i;0) \sum_{k=0}^{t_{X^{i+1}}} \frac{1}{k!} e^{h_{t}} z^{+}_{t} (i) P_{t}(i;0) e^{z^{+}_{t}(i)};$$
 (4)

hence $P_t(i;0)$ e $z_t^{z_t^+(i)}$ and (3) can be re-written as

$$P_{t}(i;k) = \frac{\sum_{t=1}^{h} (i)^{i_{k}} e^{z_{t}^{+}(i)}}{k!};$$
(5)

which is the Poisson distribution with $m ean z_t^+$ (i).

Denoting the overall in-degree distribution by $P_t(k)$, we see that it can be estim ated by averaging (5) over all nodes, that is,

$$P_{t}(k) = \frac{1}{t} \frac{x^{t}}{\sum_{i=1}^{t} \frac{\left[z \ln (t=i)\right]^{k} e^{-z \ln (t=i)}}{k!}}{k!} = \frac{1}{zk!} \frac{z^{\ln t}}{\sum_{x=0}^{t} x^{k} e^{-(1+1=z)x} dx;$$
(6)

where $x = z \ln (t=i)$ (hence di = $(t=z)e^{x=z}dx$). The integrand in (6) peaks at x = kz=(z + 1) with a nite value, so the integral can be extended to in nity, yielding

$$P_{t}(k) = \frac{1}{zk!} \sum_{x=0}^{Z^{k}} x^{k} e^{(1+1-z)x} dx = \frac{z^{k}}{(z+1)^{k+1}}$$
(7)

Let us now turn to the case in which node i, upon entering the digraph at time t = i, connects out to node j 2 f1;::; ig with probability proportional to how m any incoming edges j already has (plus 1, to ensure nonzero probabilities to start with) while aim ing at the same m ean out-degree z. In this case, an edge is deployed from ito j with probability

$$z \quad \frac{1+d_{t-1}^{+}(j)}{t+\sum_{u=1}^{t}d_{t-1}^{+}(u)}; \qquad (8)$$

with the understanding that $d_{t-1}^{+}(i) = 0$, provided $z[1 + d_{t-1}^{+}(j)] + t + \int_{u=1}^{t} d_{t-1}^{+}(u)$ (when this does not hold, then the deployment is assumed to take place with probability 1). While (8) trivially ensures that the expected value z_t (i) of d_t (i) is equal to z for all t that render the expression in (8) a legitimate probability, the underlying distribution of the out-degree d_t (i) is not so easily deduced, although by the results surveyed in [3] its average at

time to ver all the t nodes already present in the digraph must be proportional to k for some > 0.

The same holds for the average distribution $P_t(k)$ of the in-degrees d_t^+ (i). The expected value z_t^+ (i) of d_t^+ (i), in turn, can be assessed as follows. By (8), the expected probability that node i connects out to node j at time t = i is

$$z \quad \frac{1+z_{t-1}^{+}(j)}{t+\sum_{u=1}^{t} z_{t-1}^{+}(u)} : \qquad (9)$$

Thus, for t i, z_t^{\dagger} (i) obeys the recurrence

$$z_{t}^{+}(i) = z_{t-1}^{+}(i) + z \quad \frac{1 + z_{t-1}^{+}(i)}{t + \frac{P_{t-1}}{u = 1} z_{t-1}^{+}(u)}$$

$$z_{t-1}^{+}(i) + z \quad \frac{1 + z_{t-1}^{+}(i)}{t(z+1)}$$

$$= 1 + \frac{z}{t(z+1)} \quad z_{t-1}^{+}(i) + \frac{z}{t(z+1)};$$
(10)

with z_t^+ (i) = 0 fort < i.

It follows from (10) that

$$z_{t}^{+}(i) = \sum_{x=i}^{Y^{t}} 1 + \frac{z}{x(z+1)}$$
 1: (11)

Fort i 0, this solution can be approximated as

$$z_{t}^{+}$$
 (i) $\frac{t}{i}$ $\frac{z}{z+1}$ 1; (12)

as shown in Appendix B.

2.2 Results of simulations

We have conducted extensive simulations to evaluate the results presented in Section 2.1 on the in-degree distributions when the digraph evolves in a nearly acyclic fashion. The results we present in this section are averages of the quantities of interest over a large number of repetitions.

Figures 1 and 2 refer to the case in which edges are deployed uniform ly, that is, the Poisson-distributed number of edges that outgo from each node as it



Fig. 1. A verage in-degree distribution in the nearly acyclic case for $i = 10^3$ and z = 1;4;8;12 when edges are deployed uniform by $(10^5 \text{ sim ulation runs})$. Solid plots give the analytic prediction of (5).

enters the digraph is directed uniform ly towards the nodes already present. Figure 1 contains plots of $P_t(i;k)$ from both the simulations and the analytic prediction of (5) for di erent values of z and t, always with $i = 10^3$. As the plots indicate, agreem ent is very good throughout.

The plots of Figure 2 all refer to $P_t(k)$, for which our analytic prediction is the one in (7). We show data for $t = 10^4$ and several values of z. Notice that once again we obtain good agreem ent between simulation and analytic prediction, but the two start to separate as k increases. The reason for this is clear: as k increases, the peak of the integrand of (6), occurring at x = kz = (z + 1), shifts continually to the right, which renders the extension of the integral's upper limit from the z lnt of (6) to in nity ever less justi able.

D ata for the case of preferential attachment, in which edges get directed towards nodes already in the digraph with probabilities that are proportional to how m any incoming edges those nodes already have, are given in Figures 3 and 4.Figure 3 is given for $t = 10^4$ and a set of di erent z values, and con m s our expectation that in this case $P_t(k)$ is a power law.

For $i = 10^3$ and a few di erent values of z, F igure 4 contrasts simulation data for z_t^+ (i) with the analytic prediction of (12). Evidently, the two agree very well.



Fig. 2. A verage in-degree distribution in the nearly acyclic case for $t = 10^4$ when edges are deployed uniform by $(10^5 \text{ simulation runs})$. Solid plots give the analytic prediction of (7).



Fig. 3. A verage in-degree distribution in the nearly acyclic case for $t = 10^4$ when edges are deployed proportionally to the nodes' in-degrees (10^5 simulation runs).

3 Evolution allowing for cycles

3.1 Analytic results

Our second evolution scenario allows for the appearance of directed cycles other than self-loops by incorporating a probability, denoted by p_r , to control the replacement of certain edges by certain others. At each time step t > 0, one of the following two actions is selected to take place, recalling that the initial number of nodes is $n_0 = 0$ and that, as before, nodes are numbered consecutively from 1 as they enter the digraph:



Fig. 4. A verage in-degree in the nearly acyclic case for $i = 10^3$ and z = 1;4;8;12 when edges are deployed proportionally to the nodes' in-degrees (10^5 simulation runs). Solid plots give the analytic prediction of (12).

W ith probability p, the number of nodes in the digraph remains the same, thus $n_t = n_{t-1}$. In addition, a node i for which $d_{t-1}^+(i) > 0$ is selected and one of its incoming edges, chosen random ly with a uniform distribution, is replaced by a new edge directed away from i. Letting n_{t-1}^+ be the number of nodes that have nonzero in-degrees at time t = 1, the probability that node i is selected is $1=n_{t-1}^+$, provided its in-degree is nonzero.

W ith probability 1 $_{th}$ a new node, say i, is added to the digraph, so $n_t = n_{t-1} + 1$. Then a new edge is deployed from ito each of the digraph's n_t nodes with probability m infz= n_t ; 1g.

Readily, for $p_r = 0$ this evolution scenario is the one we considered rst in Section 2. For $p_r > 0$, what it does is to induce the appearance of directed cycles, and consequently of nontrivial strong components.

Under these new rules, n_t is no longer determ inistically equal to t, nor is node i necessarily the node added to the digraph at time t = i. Instead, n_t is now a random variable with expected value (1 p)t, so the expected time step at which node i is added to the digraph is t = i = (1 p), even though this can happen as early as time t = i. Once node i enters the digraph and receives its Poisson-distributed number of outgoing edges, the process of edge replacement acts to alter this number of edges random by, so we still expect the out-degree d_t (i) to be Poisson-distributed with mean z_t (i) = z.

A nalyzing the distribution of the in-degree d_t^+ (i), though, is significantly more complicated and is achieved by setting up a system of nite-difference equations to describe its behavior. As in Section 2, $P_t(i;k)$ continues to denote the distribution of in-degrees for node i at time twith t i and k 0. In what follows, we assume that node i is already present in the digraph at time t 1. We do this for the sake of sim plicity, and indicate how to compensate for it when we return to this issue later in this section.

We start with the k > 0 case. W ith probability 1 p, $P_t(i;k)$ is the sum of two probabilities, each corresponding to one of the follow ing mutually exclusive events: (i) the node added to the digraph at time t connects out to node i; (ii) the node added to the digraph at time t does not connect out to node i. W ith probability p_r , three other mutually exclusive events must be considered: (iii) node i is the node selected to have one of its incom ing edges replaced; (iv) node i is not the node selected to have one of its incom ing edges replaced, nor is the replacing edge incom ing to i; (v) node i is not the node selected to have one of its incom ing edges replaced, but the replacing edge is incom ing to i.

Let us assume that, when node i is the node selected to have one of its incoming edges replaced, the probability that the random ly chosen replacing edge forms a self-bop, or connects out to a node towards which an edge from i already exists, is negligible. This is certain to hold as the digraph acquires more nodes, and allow sus to conclude the following. In case of event (i) or (v), the in-degree of node i is increased by 1 from timet 1 to time t, while in cases (ii) and (iv) it remains the same and in case (iii) it is decreased by 1. For k > 0, $P_t(i;k)$ is then such that

where we have used the fact that $n_t = n_{t-1}$ in the case of edge replacement at time t. For $n_t n_{t-1}^+$ 1, and approximating n_t by its expected value (1 p)t, (13) becomes

$$P_{t}(\mathbf{i};\mathbf{k}) = P_{t-1}(\mathbf{i};\mathbf{k}) = \frac{p_{r}}{n_{t-1}^{+}} P_{t-1}(\mathbf{i};\mathbf{k}+1)$$

$$(1 \quad p)_{t} + \frac{p_{r}}{n_{t-1}^{+}} + \frac{p_{r}}{(1 \quad p)t} P_{t-1}(\mathbf{i};\mathbf{k})$$

$$+ (1 \quad p)_{t} + \frac{p_{r}}{(1 \quad p)t} P_{t-1}(\mathbf{i};\mathbf{k}-1); \quad (14)$$

with $t = m \inf fz = (1 p)t; 1g$.

For the case of k = 0, the only meaningful possibilities are events (ii) { (iv), hence

$$P_{t}(\mathbf{i};0) = P_{t-1}(\mathbf{i};0) = \frac{p_{r}}{n_{t-1}^{+}} P_{t-1}(\mathbf{i};1)$$

$$(1 = p_{r})_{t} + \frac{p_{r}}{(1 = p_{r})t} P_{t-1}(\mathbf{i};0): \quad (15)$$

Note in addition that, for t > 0, it is also possible to approximate n_t^+ by its expected value, that is,

$$n_{t}^{+}$$
 n_{t} $P_{t}(j;0)$ (1 p)t $P_{t}(j;0);$ (16)
 $j=1$ $j=1$

where the sum m ations give the expected number of nodes that have zero indegree at time t. The system of nite-di erence equations given for k 0 by (14){(16) can then be solved numerically for all t 1 and all i t. The greatest possible in-degree for xed i and t ist i+ 1, corresponding to the case in which node i is added to the digraph at time step i and acquires a new incoming edge at every subsequent step, so k 2 f0;:::;t i+ 1g. All boundary conditions can be set to zero, and the value of n_0^+ is imm aterial, so long as it is set to some nonzero constant.

In order to compensate for the fact that the nite-di erence equations being solved for time t and node i are based on the assumption that node i is in the digraph at time t 1, and also to ensure that a nontrivial solution is obtained, we may heuristically correct the equations for all t 1, all i t, and every appropriate k, by adding adequate probability terms to relect the appearance of edges. If t(i) is the probability that node i is added to the digraph at time t, then we add $t_{1k} t(i)$ minfz=i;1g to (14), where t_{1k} is the K ronecker delta function for k = 1, and $t_{1k}(i)$ (1 minfz=i;1g) to (15).

As for t (i), it is given by

$$t_{t}(i) = (1 p) \frac{t}{i} \frac{1}{1} (1 p)^{i} p_{r}^{t} \frac{i}{r};$$
 (17)

and thus adm its an approximation based on the Poisson distribution, that is,

 $_{t}$ (i) (1 p)P ((1 p)(t 1); i 1); (18)

where we recall that P(x;k) denotes the Poisson distribution with m ean x (cf. [1]).



Fig. 5. A verage out-degree distribution under directed-edge replacements for $i = 10^3$, z = 1;4;8;12, and $p_r = 0.25;0:75$ (10^5 simulation runs). Solid plots show the Poisson distribution with mean given by the corresponding value of z.

3.2 Results of simulations

We start with a presentation of simulation data intended to be confronted with our predictions in Section 3.1 regarding the distributions of in- and outdegrees when, during its evolution, the digraph is allowed to have directed cycles other than self-loops by the action of edge replacements. These data are presented in Figures 5 and 6.

Figure 5 contains plots for $p_r = 0.25$ and $p_r = 0.75$, along with a few values of z and t. Each plot shows simulation data on the distribution of the out-degree d_t (i) of node $i = 10^3$. In the gure, we use Q_t (i;k) to denote the probability that d_t (i) = k, and also show the corresponding Poisson distribution, P (z;k). As we mentioned earlier, our expectation is for Poisson-distributed out-degrees, because the number of outgoing edges a node is given upon entering the digraph is thus distributed, and from there onward all that the edge-replacement mechanism does is to random by alter the initial deployment of edges. We see in Figure 5 that this is indeed the case to a very good degree of agreement between Q_t (i;k) and the Poisson distribution with mean z.

D at a for the in-degree distribution $P_t(i;k)$ are shown in Figure 6 for $i = 10^3$ and two values of $p_r (p_r = 0.25 \text{ and } p_r = 0.75)$. A llplots are shown as functions



Fig. 6. A verage in-degree distribution under directed-edge replacements for $i = 10^3$, z = 1;4;8;12, and $p_r = 0.25;0.75$ (10^5 simulation nuns). Panels placed side by side share the same value of z. Solid plots give the solution of the system of - nite-di erence equations in (14) { (16).

oft for a few di erent values of z and k. For each combination of z and k values, two plots are given, one for simulation data and another to depict the solution of the system of nite-di erence equations in (14){(16) that for k 0 describes the behavior of P_t (i;k) for xed p_r and z. It is clear from the gure that agreem ent is very good between the two plots in all cases, with the very few exceptions of some of the higher values of k, but such discrepancies are the result of insu cient statistics for those particular values of k, despite the 10^5 simulation runs. A greem ent is so good, in fact, that solving the system of equations becomes largely preferable to simulating the system a su ciently large num ber of times, since the latter is unavoidably slower by several orders of m agnitude.

A by-product of our simulations has been the empirical noting that the expected value of n_t^+ is [z=(z+1)](1 p)t, where we recall that (1 p)t is the expected value of n_t^- . This fact is very useful, since it decouples the calculation of n_t^+ from that of the in-degree distributions of all nodes (cf. (16)). Consequently, it becomes possible to solve the equations in (14) and (15) for only a selected set of nodes of interest.

We now present a series of four gures, Figures 7{10, where we show data related to the behavior of the digraph's strong components, which under the policy of edge replacement exist in nontrivial form. All the plots given in



Fig. 7. A verage number of strong components (NSC), of cycle components (NCC), and of knots (NK) as a function of z for $p_r = 0.25; 0.75$ ($10^5 \text{ sim ulation runs}$).

these gures are shown as functions of z for $p_r = 0.25$, $p_r = 0.75$, and a few relevant t values. The data presented henceforth are then to be contrasted with the expectation we have from the static case studied in [1], since the dynam ic case is expected to become qualitatively equivalent to the static case for large enough t, so long as $p_r > 0$ (the larger the value of p_r , the sconer the equivalence can be observed as t grow s). The intuition behind this statem ent is that, when edges are continually replaced, the passage of enough time steps is expected to add such a degree of random ization to the positioning of the edges that it becomes probabilistically indistinguishable from any positioning that could result if edges were deployed totally random ly to begin with, as in the static case.

Figure 7 is devoted to showing how the number of strong components behaved during the simulations we conducted, and also in particular the number of cycle components and of knots. In the static case, the number of strong components tends to one quickly right past z = 1 as z is increased and the giant strong component appears. In the dynam ic case we expect a similar effect as both z and t are increased, that is, we expect the number of strong components to be dram atically reduced. This is what the gure demonstrates, particularly for the higher value of p_r .

Regarding the number of cycle components, in the static case it peaks at z = 1 and decreases rapidly to either side, eventually approaching zero. The

data in F igure 7 tend to support the expected peak at z = 1, but even for the higher value of p_r a nonzero number of cycle components, albeit small, seem s to be sustained as both z and t grow. We will see shortly that this is due to the presence of cycle components that are self-bops, and presume that it can be explained by demonstrating that such components have a non-negligible probability of appearing for small values of t, being on the other hand less likely to be picked for replacement as t increases.

In the static case, knots are expected to comprise one single node and to occur sparsely for very small values of z, then to be altogether absent as z is increased but still kept below $\ln(2n)$, and then to occur as a single, all-encompassing knot for $z > \ln(2n)$. In the dynamic case, we expect a similar behavior, including the e ect around the threshold $z = \ln(2n_t)$ $\ln(2(1 p)t)$. Taking $p_r = 0.75$ and t = 8000, for example, yields z = 8.3, which agrees well with what is shown in Figure 7. But the absence of knots for the intermediate values of z seems too brief when contrasted with the expectation created by the static case. Returning to the corresponding data in [1], speci cally Figure 6, we see that our expectation was built on one single simulation run, which may have been the cause for the seem ingly larger intervals of z inside which knots were totally absent.

These observations are complemented by the data shown in Figures 8(10, where minimum and maximum sizes are shown for the strong components in general, the cycle components, and the knots, respectively, observed during the simulations. As shown in Figure 8, the minimum size of a strong component is one, particularly as t grows, while the maximum approaches n_t (1 p)t as z grows away from one, relecting the expected appearance of the giant strong component component comprising all n_t nodes.

M inimum and maximum sizes observed for the cycle components are as shown in Figure 9. A side for a small vicinity near z = 1, where sizes larger than one occur, cycle components have size one for all values of t. This lends support to our earlier suspicion that the cycle components still persisting as both z and t grow are in fact size-one components.

The case of knots is illustrated by the data in Figure 10. It shows that the single knot that appears suddenly roughly around $z = \ln (2(1 p)t)$ is indeed all-encom passing, as its size tends to n_t (1 p)t as z grows, thus con m ing our expectations from the static case.



Fig. 8. A verage m inimum and maximum sizes of strong components (SC) as a function of z for $p_r = 0.25$;0:75 (10⁵ simulation runs).



Fig. 9.A verage m in im um and m axim um sizes of cycle components (CC) as a function of z for p_r = 0.25;0:75 (10 5 simulation runs).



Fig. 10. Average minimum and maximum sizes of knots as a function of z for $p_r = 0.25; 0.75$ (10⁵ simulation runs).

4 Conclusions

We have in this paper considered random digraphs that grow in discrete time by the continual addition of new nodes and edges. Within this context, we started with a study of digraphs that remain nearly acyclic at all times as nodes are given a Poisson-distributed number of outgoing edges upon entering the digraph. For twom odes of attachment to the nodes already in the digraph, uniform and preferential, we contributed analytic and simulation results that describe the digraph's degree distributions.

In the second part of the paper, we allowed edges to be random ly replaced during the evolution of the digraph, aim ing at allow ing nontrivial strong com – ponents to appear. In this case, too, we contributed analytic and simulation results, with emphasis on the introduction of a system of nite-di erence equations for the computation of a node's in-degree distribution at all times. We nalized by returning to the strong components that were our main subject in [1] and investigated their appearance and behavior along the evolution of the digraph.

W emention that it is possible to extend our study of the evolution of a digraph and its strong components to, in principle, all kinds of connection rules and evolution strategies, particularly those that have special signi cance within a certain application area. The ones we studied were ultim ately targeted at the study of strong components, and for this reason the absence or presence of directed cycles, as well as the conditions for their appearance during the evolution, have been crucially in portant. O ther guiding principles will certainly exist as the focus is shifted by some other motivation.

A Derivation of (3)

Note set that, for each subset of fi;:::;tg comprising k nodes, $P_t(i;k) expresses the probability that an edge exists directed from each of those k nodes to node i, but not from any of the remaining t i+1 k nodes of fi;::;tg. Let C denote the set of all partitions (C;C) of fi;::;tg such that C contains exactly k nodes. Then we have$

$$P_{t}(\mathbf{i};\mathbf{k}) = \begin{array}{c} X & Y & \frac{z}{x} & Y & 1 & \frac{z}{x} \\ & & & \\ & & & \\ & & & \\ & &$$

Since $P_t(i;0) = {}^{Q} t_{x=i}(1 z=x)$, we can write (A.1) as

$$P_{t}(\mathbf{i};\mathbf{k}) = P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j X X Y 1 + \frac{z}{x z}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j t \mathbf{i} + 1 j X Y 1 + \frac{z}{x z}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j t \mathbf{i} + 1 j X Y 1 + \frac{z}{x z}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j \frac{t \mathbf{i} + 1}{t \mathbf{i} + 1} k^{l} X Y 1 + \frac{z}{x z}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j \frac{t \mathbf{i} + 1}{t \mathbf{i} + 1} j \frac{k^{l} X Y}{j \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j}} 1 + \frac{z}{x z}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j \frac{t \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j}}{t \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j} \mathbf{j}}$$

$$= P_{t}(\mathbf{i};0) \sum_{j=0}^{X^{k}} (1)^{k} j \frac{t \mathbf{j} \mathbf{j} \mathbf{j}}{t \mathbf{j} \mathbf{j} \mathbf{j}}$$

$$= P_{t}(\mathbf{j};0) \sum_{j=0}^{X^{k}} (1)^{k} j \frac{t \mathbf{j} \mathbf{j} \mathbf{j}}{t \mathbf{j} \mathbf{j} \mathbf{j}}$$

Fork t i, in (A 2) we can use the approximations

$$\frac{\substack{k \ i+1}{k}}{\substack{t \ i+1 \ i}} \frac{j!(t \ i+1)^{k}}{k!(t \ i+1)^{j}}$$
(A.3)

which yield

$$P_{t}(\mathbf{i};\mathbf{k}) = \frac{P_{t}(\mathbf{i};\mathbf{0})}{\mathbf{k}!} \frac{\mathbf{x}^{k}}{\mathbf{j}=0} \mathbf{x}^{k} (t+\mathbf{i} + \mathbf{i})^{k} \mathbf{j} + \mathbf{i} + \mathbf{i} + \mathbf{k} \frac{\mathbf{t} \mathbf{z}}{\mathbf{i} \mathbf{z} \mathbf{1}} \mathbf{z}^{\mathbf{z} \mathbf{j}}$$
$$= \frac{P_{t}(\mathbf{i};\mathbf{0})}{\mathbf{k}!} \mathbf{k} \frac{\mathbf{t} \mathbf{z}}{\mathbf{i} \mathbf{z} \mathbf{1}} \mathbf{z}^{\mathbf{z} \mathbf{k}}$$
(A.5)

and, for z i;t,

$$P_{t}(i;k) = \frac{P_{t}(i;0)}{k!} \ln \frac{t}{i} = \frac{P_{t}(i;0)}{k!} r_{t}^{+}(i)^{i_{k}}; \qquad (A.6)$$

by (2).

B Derivation of (12)

From (11), we have

$$z_{t}^{+}(i) = \frac{Y^{t}}{\sum_{x=i}^{x=i}} 1 + \frac{z}{x(z+1)}$$

$$= \frac{{}^{t}X^{i+1}}{\sum_{j=0}^{z} \frac{z}{z+1}} \frac{z}{\sum_{i=x_{1} < j < x}^{j} \frac{x}{x_{1}}} \frac{1}{\sum_{j=x_{1} < j < x}^{z} \frac{1}{x_{1}}} \frac{1}{\sum_{j=x_{1} < j < x}^{z} \frac{1}{x_{1}}}$$
(B.1)

Fort i 0, in (B.1) the innerm ost sum mation can be approximated by

$$\frac{1}{j!} \frac{x^{t}}{x=i} \frac{1}{x} \frac{1}{j!} \ln \frac{t}{i} \frac{j}{1}; \quad (B 2)$$

thus yielding, for i 0,

$$z_{t}^{+}(i) \qquad \frac{{}^{t}X^{i+1}}{j!} \frac{1}{j!} \ln \frac{t}{i} \frac{z}{z+1}^{\frac{z}{z+1}} \qquad 1 \qquad \frac{t}{i} \frac{z}{z+1} \qquad 1: \qquad (B.3)$$

and

A cknow ledgem ents

The authors acknow ledge partial support from CNPq, CAPES, the PRONEX initiative of Brazil's MCT under contracts 41.96.0857.00 and 41.96.0886.00, and a FAPERJ BBP grant.

References

- [1] V.C.Barbosa, R.D onangelo, S.R.Souza, D irected cycles and related structures in random graphs: I Static properties, Physica A 321 (2003) 381{397.
- [2] P. Erdøs, A. Renyi, On the evolution of random graphs, Publications of the M athem atical Institute of the Hungarian A cademy of Sciences 5 (1960) 17{61.
- [3] S.Bornholdt, H.G.Schuster (Eds.), Handbook of Graphs and Networks, W iley-VCH, W einheim, Germany, 2003.
- [4] B. Bollobas, O. M. Riordan, M athem atical results on scale-free random graphs, in: Bornholdt and Schuster [3], pp.1{34.
- [5]C.Cooper, A.Frieze, A generalmodel of web graphs, Random Structures and A lgorithm s 22 (2003) 311{335.
- [6] R.L.G raham, D.E.K nuth, O.Patashnik, Concrete M athematics, 2nd Edition, Addison-Wesley, Boston, MA, 1994.