

$$T' = T, S' = \gamma S, \mathcal{P}' = \gamma^2 \mathcal{P}, \mathcal{V}' = \mathcal{V}/\gamma, \text{ and } U' = \gamma U$$

Ryszard Zygadło

Marian Smoluchowski Institute of Physics, Jagiellonian University, Reymonta 4, PL-30059 Kraków, Poland

(Dated: October 5, 2018)

It is shown that the Lorentz invariant temperature $T' = T$ and entropy $S' = \gamma S$, pressure $\mathcal{P}' = \gamma^2 \mathcal{P}$, volume $\mathcal{V}' = \mathcal{V}/\gamma$, and energy $U' = \gamma U$, where $\gamma = (1 - V^2/c^2)^{-1/2}$, are the correct transformational properties of the thermodynamical quantities of the relativistic ideal gas.

The old problem of transformational properties, see Ref. [1] for details, of the thermodynamical quantities, which enter to the basic equation,

$$dU' = T' dS' - \mathcal{P}' d\mathcal{V}', \quad (1)$$

in the moving (along x -axis with the constant velocity V) reference frame, is considered for the ideal relativistic gas. The basic thermodynamical definitions are used and new results are obtained.

The description (1) – *thermodynamics* – requires the following:

- (i) all three terms of Eq. (1) transform according to the same rule;
- (ii) the volume of the box transforms like $\mathcal{V}' = \mathcal{V}/\gamma$, where $\gamma = (1 - V^2/c^2)^{-1/2}$;
- (iii) the energy transforms like

$$U' = \sum_i \langle \varepsilon'_i \rangle = \sum_i \langle \gamma [\varepsilon_i - (p_x)_i V] \rangle = \sum_i \gamma \langle \varepsilon_i \rangle = \gamma U, \quad (2)$$

where $\langle (p_x)_i \rangle = 0$ in the common resting reference frame of a system (as a whole) and thermal bath;

(iv) the entropy is known (as well as the other thermodynamical quantities) *explicitly* in the resting reference frame from Gibbs–Boltzmann equilibrium distribution, particularly

$$dS = \phi_1(T) dT + \phi_2(\mathcal{V}) d\mathcal{V}, \quad (3)$$

where $\phi_1 [= T c_V]$ and ϕ_2 is a (known) *nonhomogeneous* and *homogeneous* (of the order -1 , which is going to appear *crucial*) function of the single argument, respectively.

The rule for the pressure,

$$\mathcal{P}' = \gamma^2 \mathcal{P}, \quad (4)$$

follows as a *necessary condition* for the selfconsistence of thermodynamics. The same result (4) is independently obtained from the mechanical consideration. In fact, considering the point particles (i.e., ideal gas) we can neglect the collisions between particles. Then each particle undergoes the successive collisions with the certain wall of the box ($L \times L \times L = \mathcal{V}$) with certain constant period $\tau_u = 2L\varepsilon/p_u$, $u = x, y, z$, respectively in each direction. In the moving reference frame all τ_u change according to the Lorentz dilatation $\tau'_u = \tau_u/\gamma$. The momentum transfer associated with the single collision in y and z direction is the same $2p'_y = 2p_y$,

$2p'_z = 2p_z$ in the moving reference frame and the areas change according to $\sigma'_y = \sigma_y/\gamma$, $\sigma'_z = \sigma_z/\gamma$. The energy–momentum transfer in x -direction $(0, 2p_x, 0, 0)$ corresponds to $(-2V\gamma p_x, 2\gamma p_x, 0, 0)$ in the moving reference frame and $\sigma'_x = \sigma_x$. Consequently the each ratio $2p_u/\tau_u\sigma_u$ change by the factor γ^2 and thus $\mathcal{P}' = \gamma^2 \mathcal{P}$.

Comparing

$$dU' = T' dS' - \mathcal{P}' d\mathcal{V}' = \gamma(T dS - \mathcal{P} d\mathcal{V}) = \gamma dU \quad (5)$$

and using the pressure transformation formula (4) one obtains

$$(T'/\gamma T) dS'(T', \mathcal{V}/\gamma) = dS(T, \mathcal{V}). \quad (6)$$

Consequently, see Eq. (3), $(T'/\gamma T) \partial S'/\partial T' \equiv (T'/\gamma T) \psi_1(T')$ cannot depend on \mathcal{V}' . Because $0 = \partial \psi_1/\partial \mathcal{V}' = \partial^2 S'/\partial \mathcal{V}' \partial T'$ one obtains $\partial S'/\partial \mathcal{V}' \equiv \psi_2(\mathcal{V}')$ depending only on \mathcal{V}' . The equality

$$(T'/\gamma^2 T) \psi_2(\mathcal{V}/\gamma) = \phi_2(\mathcal{V}) \quad (7)$$

can only be satisfied (*identically* with respect to \mathcal{V}) if

$$T' = \alpha(V) T \quad (8)$$

and both ψ_2 and ϕ_2 are *homogeneous* functions (of the same order) of the argument. Supposing for a moment that $\phi_2 \propto \mathcal{V}^k$ is a homogeneous of the certain order k [for the relativistic ideal gas $\phi_2(\mathcal{V}) = Nk_B/\mathcal{V}$, so $k = -1$] one obtains

$$\psi_2 = (\gamma^{k+2}/\alpha) \phi_2. \quad (9)$$

The resulting by the use of Eq. (8) second equality

$$(\alpha^2/\gamma) \psi_1(\alpha T) = \phi_1(T), \quad (10)$$

because ϕ_1 for the relativistic ideal gas *explicitly* appears to be *nonhomogeneous* function, can only be satisfied *identically*

$$\alpha(V) \equiv 1, \quad (11)$$

and

$$\psi_1 = \gamma \phi_1. \quad (12)$$

Thus the Lorentz invariant temperature is obtained as a *necessary condition* for the selfconsistence of thermodynamics.

Simultaneously, the entropy transforms like the energy,

$$dS' = \gamma\phi_1 dT' + \gamma^{k+2}\phi_2 d\mathcal{V}' \quad (13)$$

and $k = -1$, $\phi_2(\mathcal{V}') = Nk_B/\mathcal{V}'$, are obtained as a *necessary and sufficient* condition for the selfconsistence of thermodynamics.

Consequently the Gibbs–Boltzmann *thermodynamics* of the ideal relativistic gas is *consistent*, and

$$T' = T, S' = \gamma S, \mathcal{P}' = \gamma^2 \mathcal{P}, U' = \gamma U \quad (14)$$

are the resulting Lorentz transformation formulas.

The Eqs. (14) for the ideal gas system are obtained rigorously, just from the basic definition (1). They significantly differs from the previous ones, where entropy and pressure have been considered to be invariant, from so-called Einstein–Planck formulas [2],

$$T' = T/\gamma, S' = S, \mathcal{P}' = \mathcal{P}, U' = U/\gamma \quad (15)$$

and from the Ott formulas [3]

$$U' = \gamma U, T' = \gamma T, S' = S. \quad (16)$$

The different previous results are collected in Ref. [1]. There is a number of reasons, for which those results should be treated with care:

I) using the same *formalism* different groups have obtained the different results and no consense has been reached;

II) the transformational properties are obtained under the assumption that the relations between thermodynamical functions and the canonical partition function remain unchanged in moving reference frame;

III) the approach is tested on the ideal gas model, however the incorrectly transformed partition function is used.

There is the difference between transformational properties of the probability density distributions of the ideal and of the interacting system [4,5]. For the ideal gas the effect of Lorentz transformation on space variables is counted *only* globally $\mathcal{V}' = \mathcal{V}/\gamma$ and thus the momentum distribution change on non-covariant manner

$$\varepsilon W(\vec{p}) = \varepsilon' W'(\vec{p}'), \quad (17)$$

in order to conserve the probability measure. For interacting particles there is no variable separation and the joint probability density distribution in phase-space is invariant

$$f'(\vec{x}'_i, \vec{p}'_i) = f(\vec{x}_i, \vec{p}_i). \quad (18)$$

There is no contradiction between (17) and (18). The special relativity requires the full dynamical description of the system to be covariant. For the projected kinetic, which is involved with the particular choice of variables going to be reduced the Lorentz covariance is not already expected. It is the reason that the first (non-covariant

and true) rule can be obtained from the general one. Both, (17) and resulting from the improper use of (18) $f'(\vec{p}') = f(\vec{p})$ cannot be simultaneously true. It is the basic error of the previous consideration, which manifests if the ideal gas is studied [1].

The particular *explicit* expressions are presented for 1D case ($c = 1$ and $N = 1$ is used). One has

$$W'(p') = \frac{1 + Vp'/\sqrt{p'^2 + m^2}}{2m\sqrt{1 - V^2}K_1(m\beta)} \exp\left[-\frac{\sqrt{p'^2 + m^2} + Vp'}{\beta^{-1}\sqrt{1 - V^2}}\right], \quad (19)$$

$$\langle \varepsilon' \rangle = \int dp' W'(p') \sqrt{p'^2 + m^2} = \frac{-mK_1'(m\beta)}{\sqrt{1 - V^2}K_1(m\beta)}, \quad (20)$$

$$\langle p' \rangle = \frac{mVK_1'(m\beta)}{\sqrt{1 - V^2}K_1(m\beta)}, \quad (21)$$

$$Z'_1(\mathcal{V}', \beta) = h^{-1}\sqrt{1 - V^2}\mathcal{V}'2mK_1(m\beta). \quad (22)$$

and, in the resting reference frame ($V = 0$),

$$\mathcal{P} = \langle p^2/(\mathcal{V}\sqrt{p^2 + m^2}) \rangle = 1/\beta\mathcal{V} = k_B T/\mathcal{V}. \quad (23)$$

$$\begin{aligned} S &= -k_B \langle \log[hW(p)/\mathcal{V}] \rangle \\ &= k_B \{ \beta \langle \varepsilon \rangle + \log[2m\mathcal{V}K_1(m\beta)/h] \}, \end{aligned} \quad (24)$$

where the Planck constant is introduced for dimensional purposes and K_1 is a modified Bessel function.

From Eqs. (20–24) the general conclusion follows that the *functional* relations of the ordinary thermodynamics (in the resting reference frame) are not invariant. For instance, see (23), $\gamma^2 \mathcal{P} = \mathcal{P}' \neq k_B T'/\mathcal{V}' = \gamma \mathcal{P}$. Similarly, $-\partial \ln(Z'_1)/\partial \beta = \langle \gamma[\varepsilon' + p'V] \rangle = \langle \varepsilon \rangle$ do not correspond to the mean energy (20) in the moving reference frame.

The important results of the paper are the following. It has been shown that the previous transformational formulas were incorrect. The most popular Einstein–Planck relations (15) are consistent with the condition (i), however they adopt incorrect energy transformation rule. The Eqs. (2) cannot be questioned, because except the trivial statement that the average is a linear operation, these are in fact the definitions of the successive terms. The previously used invariant properties of the entropy and the pressure are not true. The pressure transformation formula (4), mentioned in Ref. [6], has been proven also by mechanical consideration. It has been shown that the temperature is invariant, as was first suggested by Landsberg [7]. The covariance of Eq. (1) for ideal systems do not require particular condition from $\phi_1(T)$, so the relativistic thermodynamics (1) of ideal gas is possible also for the quantum statistics. The black body radiation thermodynamics is not of the form (3), but otherwise has the property that *all* functions are *homogeneous*. This explains why the consideration of the gas

of photons cannot lead to unambiguous conclusions. In fact, because $U = c\mathcal{V}T^4$, $\mathcal{P} = cT^4/3$, and $S = 4c\mathcal{V}T^3/3$, then the assumption of arbitrary transformational rules of the form

$$\begin{aligned} \mathcal{V}' &= \mathcal{V}/\gamma, \quad T' = \gamma^\tau T, \\ S'(\mathcal{V}', T') &= \gamma^\sigma S(\mathcal{V}, T) = \gamma^{\sigma+1-3\tau} 4c\mathcal{V}'T'^3/3, \\ \mathcal{P}' &= \gamma^\pi \mathcal{P} = \gamma^{\pi-4\tau} cT'^4/3, \\ U' &= \gamma^u U = \gamma^{u+1-4\tau} c\mathcal{V}'T'^4, \end{aligned} \quad (25)$$

results with only two equations for the four variables

$$u + 1 = \pi, \quad u = \sigma + \tau, \quad (26)$$

in order to retain the covariance of Eq. (1). Two arbitrary assumptions, $\pi = 0$ and $\sigma = 0$, leading to Eqs. (15) were incorrect. The proper choice, $u = 1$ results with $\pi = 2$, however τ and $\sigma = 1 - \tau$ remain undetermined. The consideration of ideal gas results with $\tau = 0$ and $\sigma = 1$.

-
- [1] R. Balescu, *Physica* **40**, 309 (1968).
 [2] A. Einstein, *Jarhb. Radioakt. Elektron.* **4**, 441 (1907); M. Planck, *Ann. Phys. (Leipzig)* **26**, 1 (1908).
 [3] H. Ott, *Z. Phys.* **175**, 70 (1963).
 [4] L. Landau and E. Lifshitz, *The Classical Theory of Fields* (Pergamon, Oxford, 1971).
 [5] S.R. de Groot, W.A. van Leeuwen, and Ch.G. van Weert, *Relativistic Kinetic Theory* (North-Holland, Amsterdam, 1980).
 [6] W. G. Sutcliffe, *Nuovo Cimento B* **39**, 683 (1965).
 [7] P.T. Landsberg, *Nature* **212**, 597 (1966); **214**, 903 (1967); P.T. Landsberg and K.H. Jones, *Nuovo Cimento B* **52**, 28 (1967).