

Maximum entropy principle for Renyi's and Tsallis' entropies

A.G.Bashkirov

Institute Dynamics of Geospheres, RAS,
Leninskii prosp. 38 (bldg.1), 119334, Moscow, Russia

Abstract

The equilibrium distributions of probabilities providing maximality of Renyi and Tsallis entropies are rederived. New S-forms of them are found which are normalised with corresponding entropies in contrast to the usual Z-forms normalised with partition functions.

PACS: 05.10.Gg, 05.20.Gg, 05.40.-a

1 Introduction

According to the well-known maximum entropy principle (MEP) developed by Jaynes [1] for a Boltzmann-Gibbs statistics the distribution of probabilities $\mathbf{p} = \{p_i\}$ provides maximality of information entropy with an additional conditions which consist in preassigning the average value $U = \langle H_i \rangle_p = \sum_i H_i p_i$ of the random quantity H_i and the requirement of normalisation of \mathbf{p} .

Then, the distribution $\{p_i\}$ is determined from the extremum of the functional

$$L_G(\mathbf{p}) = \sum_i p_i \ln p_i - \alpha_0 \sum_i H_i p_i - \alpha_1 \sum_i p_i; \quad (1)$$

where α_0 and α_1 are Lagrange multipliers. Its extremum is ensured by the Gibbs canonical distribution, in which $\alpha_0 = 1/k_B T_0$ and $\alpha_1 + 1$ is the free energy.

Really, equating its functional derivative to zero we get

$$\frac{\partial L_G(\mathbf{p})}{\partial p_i} = \ln p_i - \alpha_0 H_i - \alpha_1 - 1 = 0; \quad (2)$$

E-mail address: abas@idg.chph.ras.ru

or

$$p_i = e^{-\beta_0 - \beta_1 - \beta_0 H_i} \quad (3)$$

With account of normalization condition $\sum_i p_i = 1$ we get

$$e^{-\beta_0 - \beta_1} = Z_G \sum_i e^{-\beta_0 H_i} \quad (4)$$

and

$$p_i = Z_G^{-1} e^{-\beta_0 H_i} \quad (5)$$

On the other side, multiplying equation (2) by p_i and summing up over i , with account of normalization condition $\sum_i p_i = 1$ we get

$$\beta_0 + \beta_1 = S_B = k_B \beta_0 U \quad (6)$$

and

$$p_i = e^{-S_B = k_B \beta_0 U - \beta_0 H_i} \quad (7)$$

When $\beta_0 = 1/k_B T_0$ both equations (5) and (7) are equivalent due to definitions of a free energy $F = U - T_0 S$ and $F = -k_B T_0 \ln Z_G$. The first of them, eq. (5), may be called a Z-form and the second one, eq. (7), a S-form of the Gibbs' distribution.

2 MEP for Renyi entropy

If the Renyi entropy $S_R = \frac{k_B}{1-q} \ln \sum_i p_i^q$ is used instead of the Boltzmann entropy the equilibrium distribution must provide maximum of the functional

$$L_R(p) = \frac{1}{1-q} \ln \sum_i p_i^q - \sum_i \lambda_i H_i p_i - \sum_i \mu_i p_i; \quad (8)$$

where λ_i and μ_i are Lagrange multipliers. Note that, in the $q \rightarrow 1$ limit $L_R(p)$ passes to $L_B(p)$.

We equate a functional derivative of $L_R(p)$ to zero, then

$$\frac{\partial L_R(p)}{\partial p_i} = \frac{q}{1-q} \frac{p_i^{q-1}}{\sum_j p_j^q} - H_i - \mu_i = 0; \quad (9)$$

Multiplying this equation by p_i and summing up over i , with account of normalisation condition $\sum_i p_i = 1$ we get $\beta = \frac{q}{1-q} U$. Then, it follows from equation (9) that

$$p_i = e^{-\beta H_i} p_j^q (1 - \frac{q}{q-1} H_i)^{\frac{1}{q-1}}; \quad H_i = H_j \quad U: \quad (10)$$

Using once more the condition $\sum_i p_i = 1$ we get

$$p_j^q = \sum_i (1 - \frac{q}{q-1} H_i)^{\frac{1}{q-1}} p_j^q$$

and, finally,

$$p_i = p_i^{(RZ)} = \frac{1}{Z_R} (1 - \frac{q}{q-1} H_i)^{\frac{1}{q-1}} \quad (11)$$

$$Z_R = \sum_i (1 - \frac{q}{q-1} H_i)^{\frac{1}{q-1}}$$

This is the Renyi distribution in the Z-form. At $q \rightarrow 1$ the distribution $p_i^{(R)} g$ becomes the Z-form of Gibbs canonical distribution in which the constant $\beta = 1/k_B T_0$ is the reciprocal of the temperature.

To obtain the S-form of the Renyi distribution we can find that for the Renyi distribution (11) $S_R = -\ln Z_R$ or rewrite the sum $\sum_j p_j^q$ as $\sum_j p_j^q = \exp[(1-q)S_R/k_B]$. Then we get

$$p_i^{(RS)} = e^{-S_R/k_B} (1 - \frac{q}{q-1} H_i)^{\frac{1}{q-1}}: \quad (12)$$

At $q \rightarrow 1$ this distribution becomes the S-form of Gibbs canonical distribution.

3 MEP for Tsallis entropy

When the Tsallis entropy was used firstly instead of the Boltzmann entropy the equilibrium distribution was derived by Tsallis [2] in the form

$$p_i^T = \frac{(1 + (1-q)H_i)^{\frac{1}{q-1}}}{\sum_i (1 + (1-q)H_i)^{\frac{1}{q-1}}}; \quad (13)$$

This distribution was named later as the 1-st version of thermostatics. It was noticed there that the parameter β was not a Lagrange multiplier because of the starting functional was forced to be taken as

$$L^T(\mathbf{p}) = \frac{1}{1-q} \sum_i p_i^q - (\beta - 1) \sum_i H_i p_i + \sum_i p_i \quad (14)$$

Here, the question arises about forms of Lagrange multipliers $(\beta - 1)$ and $(+)$, but the main problem is that the functional $L^T(\mathbf{p})$ does not pass to the functional (9) when $q \rightarrow 1$ as the second term in (13) vanishes.

It seems reasonable to suppose that just this difficulty caused to introduce the 3-rd version of nonextensive thermodynamics with the escort distribution $P_i = p_i^q = \sum_j p_j^q$. The consistency of the transition to the escort distribution is partly justified by the condition conservation of a preassigned average value of the energy $U = \sum_i H_i P_i$, however other average values are to be calculated with the use of the same escort distribution also, that contradicts to the main principles of probability description.

For the generality sake, the resulted distribution of the Tsallis' 3-rd version is reproduced here as

$$p_i^{T3} = \frac{1}{Z_{T3}} \exp_{q-1} \left((1-q)(H_i - U) \right) = \frac{1}{Z_{T3}} (p_j^{T3})^{qA} \quad ; \quad (15)$$

$$Z_{T3} = \sum_i \exp_{q-1} \left((1-q)(H_i - U) \right) = \sum_j (p_j^{T3})^{qA} \quad (16)$$

where β is the true Lagrange multiplier in the corresponding variational functional of the 3-rd version.

Below, it will be shown that careful solution of the variational problem for the 1-st version of thermostatics gives rise to a probability distribution which doesn't seem to be less acceptable then p_i^{T3} .

Really, if we take the functional

$$L_T(\mathbf{p}) = \frac{1}{1-q} \left(\sum_i p_i^q - 1 \right) - \beta \sum_i H_i p_i + \sum_i p_i \quad (17)$$

and equate its functional derivative to zero we get

$$\frac{\partial L_T(\mathbf{p})}{\partial p_i} = \frac{q}{1-q} p_i^{q-1} - \beta H_i + 1 = 0 \quad (18)$$

Multiplying this equation by p_i and summing up over i , with account of normalisation condition $\sum_i p_i = 1$ we get $\sum_i p_i^q = \frac{q}{1-q} \sum_i p_i^q U_i$. Then, it follows from equation (18) that

$$\begin{aligned} p_i &= \exp^x \left(\sum_j p_j^q - \frac{q-1}{q} H_i^A \right)^{\frac{1}{q-1}} \\ &= \exp^x \left(\sum_j p_j^{qA} - \frac{q-1}{q} H_i^A \right)^{\frac{1}{q-1}} \end{aligned} \quad (19)$$

Using once more the condition $\sum_i p_i = 1$ we get

$$p_i = p_i^{TZ} = Z_{T1}^{-1} \exp^x \left(\sum_j p_j^{qA} - \frac{q-1}{q} H_i^A \right)^{\frac{1}{q-1}}; \quad (20)$$

$$Z_{T1} = \sum_i \exp^x \left(\sum_j p_j^{qA} - \frac{q-1}{q} H_i^A \right)^{\frac{1}{q-1}} \quad (21)$$

This is my modification of the 1-st version of the Tsallis distribution in the Z-form. It differs from the 3-rd version, eq. (15), by the signs before differences $q-1$ and doesn't invoke the escort distribution. In this respect it seems to be much more attractive than p^{T3} . At $q \rightarrow 1$ the distribution p^{TZ} becomes the Z-form of the Gibbs canonical distribution.

It should be noted that both p^{T3} and p^{TZ} are explicitly self-referential in contrast to p^{RS} .

To obtain the S-form of the modified 1-st version of the Tsallis distribution we rewrite the sum $\sum_j p_j^q$ in the upper line of eq. (19) in term of the Tsallis entropy. Then we get finally

$$p_i = p_i^{TS} = \frac{1}{(q-1)} \left(\frac{S_T}{k_B} - \frac{U}{q} + \frac{H_i}{q} \right)^{\frac{1}{q-1}}; \quad (22)$$

At $q \rightarrow 1$ this distribution becomes the S-form of the Gibbs canonical distribution.

4 Conclusion

In this article, I derived and rederived some useful equations for equilibrium probability distributions on the base of MEP for the Renyi and Tsallis en-

tropies. I hope that they can abandon the use of escort distributions for calculations of average values in favour of original distributions.

Besides, the new S-forms of the Renyi and Tsallis distributions are proposed. They are normalised with the use of corresponding entropies but not partition functions. These forms may appear to be useful in different applications and in construction of new, non-Gibbsian, thermostatics.

Acknowledgements

It is pleasure to thank A. Vityazev for fruitful discussions and supporting this work, and I acknowledge warm, friendly and illuminating discussions with S. Abe, C. Beck, A. Wang and G. Wilk held at NEXT2003 on Sardgenia.

References

- [1] Jaynes E.T. Phys.Rev. **106**, 620; **107**, 171 (1957)
- [2] Tsallis C. J.Stat.Phys. **52**, 479 (1988)