

Energy Gaps and Roton Structure above the $\ell = 1=2$ Laughlin State of a Rotating Dilute Bose-Einstein Condensate

Tatsuya Nakajima¹ and Masahito Ueda²

¹Physics Department, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan

²Department of Physics, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan, and CREST, Japan Science and Technology Corporation (JST), Saitama 332-0012, Japan

(Dated: 11 March 2003; published as PRL 91 (2003) 140401)

Exact diagonalization study of a rotating dilute Bose-Einstein condensate reveals that as the first vortex enters the system the degeneracy of the low-energy yrast spectrum is lifted and a large energy gap emerges. As more vortices enter with faster rotation, the energy gap decreases towards zero, but eventually the spectrum exhibits a rotonlike structure above the $\ell = 1=2$ Laughlin state without having a phonon branch despite the short-range nature of the interaction.

PACS numbers: 03.75.Kk, 05.30.Jp, 67.40.Db

Rotating dilute Bose-Einstein condensates (BECs) have attracted considerable interest in recent years [1, 2, 3, 4]. When the single-particle energy-level spacing is much larger than the interaction energy per particle, the system becomes highly degenerate with increasing the angular momentum (AM) L . Interactions then play a pivotal role in lifting the degeneracy and determining the many-body ground state [5], in close analogy with the physics of the fractional quantum Hall effect [6, 7]. Motelson [8], on the other hand, pointed out the relevance of the yrast line, which traces the lowest-lying states of the system as a function of the AM and has served as a key concept in nuclear physics [9], to a rotating dilute BEC. In view of recent development of Feshbach techniques [10] and experimental achievements of fast rotating BECs [1, 2, 3], such an extreme but highly interdisciplinary arena seems to be within experimental reach.

Bertsch and Papenbrock numerically found that the yrast line depends linearly on L for $L \leq N$, where N is the number of bosons [11], and showed that high-lying states above the yrast line are dominated by single-particle excitations [12]. Of particular interest is the yrast spectrum for low-lying states because it gives us insights about how the many-body wave function responds as vortices enter the system. Motelson pointed out that for $L \leq N$ low-lying states are quasi-degenerate and dominated by collective multipolar excitations [8]. An exact diagonalization study [13] and subsequent analytical studies [14, 15, 16] have demonstrated the existence of the quasi-degenerate yrast spectrum that is dominated by interactions between octupole modes. The purpose of this Letter is to report the results of our exact diagonalization study on faster rotating BECs. Our primary findings are two distinct kinds of energy gaps: one above the single-vortex state and the other associated with vortex-antivortex pair excitations above the $\ell = 1=2$ Laughlin state. We also find that higher excited states with $L < N$ form linear energy bands.

Consider a weakly interacting N -boson system trapped

in an axisymmetric parabolic confining potential. The potential is assumed to be isotropic in the radial direction but in the axial direction it is assumed to be so tightly confined that all particles occupy the lowest-energy state in that direction. The problem is thus essentially two-dimensional with projected AM, L , on the symmetry axis being conserved. The many-body Hamiltonian of this system can be written as $H = \sum_i h_i + V$, where $h_i = \frac{\hbar^2}{2M} \nabla_{\mathbf{r}_i}^2 + M \frac{1}{2} \omega_z^2 r_i^2$ denotes the single-particle Hamiltonian for the i -th particle and $V = (4\hbar g/M!) \sum_{i < j} (r_i - r_j)$ describes the contact interaction between bosons having mass M . Here g characterizes the strength of interaction between bosons, $g = \hbar! = a_s = (2\hbar/M!)^{1/2}$, $!$ and $!_z$ are trap frequencies in the radial and axial directions, respectively, and a_s is the s -wave scattering length. Throughout this Letter we shall assume weak repulsive interactions such that $\hbar! > gN > 0$.

The single-particle state is characterized by the radial quantum number n and the AM quantum number m , with the corresponding eigenenergy given by $\hbar!(2n + \frac{1}{2}m + 1)$. For a given AM $L > 0$, it is energetically favorable to assume $n = 0$ and let all particles have nonnegative AM, i.e., $m \geq 0$. In this lowest-Landau-level approximation, the noninteracting part of the Hamiltonian contributes a constant term $\hbar!(L + N)$, and we shall henceforth ignore this constant part and focus on the interaction Hamiltonian. It is convenient to take as a basis set of states single-particle states described by $\phi_m(z) = (z^m / m!) \exp(-\frac{1}{2}|z|^2)$, where $z = x + iy$ and the lengths are measured in units of $(\hbar/M!)^{1/2}$. Expanding the field operator as $\hat{\psi}(z) = \sum_m \hat{c}_m \phi_m(z)$, the second-quantized form of the contact interaction Hamiltonian is written as $\hat{V} = g \sum_{m_1, m_2, m_3, m_4} V_{m_1 m_2 m_3 m_4} \hat{c}_{m_1}^\dagger \hat{c}_{m_2}^\dagger \hat{c}_{m_3} \hat{c}_{m_4}$, where the matrix elements $V_{m_1 m_2 m_3 m_4}$ are given by $V_{m_1 m_2 m_3 m_4} = \frac{1}{m_1 + m_2, m_3 + m_4} \frac{(m_1 + m_2)!}{(m_1! m_2! m_3! m_4!)}$. Our task thus reduces to finding the energy spectrum of \hat{V} under the restrictions of a fixed particle number $\sum_m \hat{c}_m^\dagger \hat{c}_m = N$ and a fixed AM $\sum_m m \hat{c}_m^\dagger \hat{c}_m = L$.

Figure 1 shows the yrast spectrum for $0 \leq L \leq 35$ with $N = 25$, where the energy is measured in units of gN from the yrast line $E = gN$ ($N - 1 \leq L \leq 2$) [11]. For $L \leq N$, there are quasi-degenerate excited states arising from pairwise repulsive interaction between octupole modes [13]. However, the excitation energies are very small, of the order of g . This quasi-degeneracy may be regarded as a precursor for spontaneous symmetry breaking of the axisymmetry associated with the entrance of the first vortex. Remarkably, the quasi-degeneracy for $L \leq N$ is lifted with increasing L and that a large energy gap of the order of gN appears as the first vortex enters the system. The energy gap implies that the many-body wave function responds to external rotation in such a manner that a single vortex state becomes stabilized.

The emergence of the energy gap can be inferred from the fact that excitations associated with the single-particle state m ($m \geq 3$) become increasingly massive, as the AM per particle L/N approaches unity. As shown in Table I, the excitation energy increases by an amount of the order of gN when the condensate develops from the nonvortex state at $L = 0$ to the single-vortex state




FIG. 1: The yrast spectrum for $0 \leq L \leq 35$ with $N = 25$. As the angular momentum per particle L/N approaches unity, excitations associated with single-particle states m ($m \geq 3$) become increasingly more massive. The amount of increase in the excitation energy is given in Table I. Among rotational bands stabilized by these energy gaps, three bands are shown by linking the collective excitations, $Q_1 \mathbf{j} = N \mathbf{i}$ (solid curve), $Q_2 \mathbf{j} = N \mathbf{i} \mathbf{i}$ (dashed curve), and $Q_3 \mathbf{j} = N \mathbf{i} \mathbf{i} \mathbf{i}$ (dotted curve), for $m = 3, 4, 5, 6, 7, 8$. Those states that include excitations of the center-of-mass motion are omitted except for the states, $Q_1 \mathbf{j} = 0 \mathbf{i}$ and $Q_1 \mathbf{j} = N \mathbf{i}$ (shown by squares). The energy is measured from the yrast line $E = gN$ ($N - 1 \leq L \leq 2$) in units of gN .

at $L = N$, and an increase in excitation energy is larger for smaller m . In particular, excitations of the $m = 3$ mode, which are nearly gapless for $L \leq N - 1$, become increasingly massive as the vortex enters the system. The energy gap continues to increase till $L = N + m - 1$ for each m . We also note that the energy bands (shown in Fig. 1 as dash-dotted lines) for $m = 4, 5, 6, 7, 8$ are almost linear. When $L \leq N - 1$, a regular structure of rotational bands is stabilized by the energy gaps associated with the single-particle states m . In Figure 1, we show three rotational bands by linking the collective excitations $Q_1 \mathbf{j} = N \mathbf{i}$, $Q_2 \mathbf{j} = N \mathbf{i} \mathbf{i}$, and $Q_3 \mathbf{j} = N \mathbf{i} \mathbf{i} \mathbf{i}$ (Eq. 7) with solid, dashed, and dotted curves, respectively. Here $Q = (1/N!) \sum_{p=1}^N z_p^N$ describes the collective multipolar excitation of order Q , and $\mathbf{j} = N \mathbf{i}$, $\mathbf{j} = N \mathbf{i} \mathbf{i}$, and $\mathbf{j} = N \mathbf{i} \mathbf{i} \mathbf{i}$ denotes the yrast states for $L = N$, $L = N - 1$, and $L = N - 2$, respectively.

The regularity of rotational bands for $L \leq N - 1$ is more clearly seen in Fig. 2, where the energy is measured from the line $E = E_{L=N-5} gN$ ($L \leq N - 5$) [17]. As in Fig. 1, the collective excitations, $Q_1 \mathbf{j} = N \mathbf{i}$, $Q_2 \mathbf{j} = N \mathbf{i} \mathbf{i}$, and $Q_3 \mathbf{j} = N \mathbf{i} \mathbf{i} \mathbf{i}$ (Eq. 7), are linked by solid, dashed, and dotted curves, respectively. Those states that contain excitations of the center-of-mass motion are not shown except for the states $Q_1 \mathbf{j} = N \mathbf{i}$, $Q_2 \mathbf{j} = N \mathbf{i} \mathbf{i}$, and $Q_3 \mathbf{j} = N \mathbf{i} \mathbf{i} \mathbf{i}$ (shown by squares). Other low-lying excitations can be understood as (multiple) excitations of these collective modes.

When $L \leq N - 1$, the system can respond in such a manner that the interaction energy becomes zero. In fact, the $Q = 1/2$ Laughlin state [18], $\Psi = \prod_{i < j} (z_i - z_j)^2$, appears at $L = N - 1$ as the first zero-interaction-energy state; the contact interaction does not affect this state because the minimum of the relative AM between particles is two for this state. In such a high AM regime, we find a remarkable energy gap to appear above the $Q = 1/2$ Laughlin state as shown in Fig. 3.

To investigate long-wavelength collective excitations while keeping the average density unaltered (i.e., neutral excitations), it is convenient to use the spherical geometry [19] rather than the disk geometry which is used thus far. The two geometries are related with each other by the stereographic mapping. The z -component of the total AM in the spherical geometry is given by $L_z = L - NS$, where $2S + 1$ is the number of single-particle states on the sphere and $2S$ is chosen to be $2(N - 1)$ in order to

m	3	4	5	6	7	8
$E = gN$	1	7/8	3/4	21/32	19/32	71/128

TABLE I: The increase in energy of the excitation associated with the single-particle state m ($m \geq 3$) when the system develops from the nonvortex condensate with $L = 0$ into the single-vortex condensate with $L = N$.




FIG. 2: Regular rotational bands above the single-vortex condensate are shown, where the energy is measured from the line $E = E_{L=N} - 5gN (L/N) = 32$ in units of gN . Among the rotational bands, three bands are shown by linking the collective excitations, $Q_{\mathbf{j}} = N\mathbf{i}$ (solid curve), $Q_{\mathbf{j}} = N-1\mathbf{i}$ (dashed curve), and $Q_{\mathbf{j}} = N-2\mathbf{i}$ (dotted curve), for $\ell = 7$. The low-lying excitations can be understood as (multiple) excitations of these collective modes. Except for the states $Q_{\mathbf{j}} = N\mathbf{i}$, $Q_{\mathbf{j}} = N-1\mathbf{i}$, and $Q_{\mathbf{j}} = N-2\mathbf{i}$ (shown by squares), those states that include excitations of the center-of-mass motion are omitted.

study the $\ell = 1/2$ Laughlin state.

Figure 3 shows the excitation spectrum calculated in the spherical geometry for $0 \leq L \leq 56$ with $N = 8$. The $\ell = 1/2$ Laughlin state appears at $L = N(N-1) = 56$ (i.e., $L_z = 0$) as the zero-interaction-energy state (i.e., the lowest energy state) for the contact repulsive interaction $(4g/R^2) \sum_{i < j} (\mathbf{r}_i - \mathbf{r}_j)^2$, where \mathbf{r}_i is a unit vector that specifies the location of the i th boson on the sphere and the radius of the sphere is given by $R = \sqrt{S/2}$. The total AM, L_{tot} , of this state is zero and the eigenstates with $L_{\text{tot}} = 0$ are plotted for $L = NS$. The eigenstates with non-zero total AM are plotted only for $L = NS + L_{\text{tot}}$ without showing their $(2L_{\text{tot}} + 1)$ -fold degeneracy.

The inset of Fig. 3 shows the excitation spectrum above the Laughlin state as a function of L_{tot} in the spherical system. A rotonlike structure manifests itself as the lowest-energy excitation branch. We have confirmed that the excitation energy remains finite (1.229 ± 0.038 in units of g) in the thermodynamic limit [20]. We have also confirmed that the feature of the rotonlike minimum becomes more pronounced if we use long-range interactions such as the Coulomb interaction, $\sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$. The two-roton continuum is also seen to exist above the single-roton branch.

To understand the physics underlying these gap-




FIG. 3: The excitation spectrum for $0 \leq L \leq 56$ with $N = 8$, calculated in the spherical geometry. As the zero-interaction-energy state (i.e., the lowest energy state), the $\ell = 1/2$ Laughlin state appears at $L = N(N-1) = 56$. In the inset, the excitation spectrum above the Laughlin state is shown against the total angular momentum in the spherical system. The lowest-energy branch exhibits a rotonlike structure without having a phonon branch despite the short-range nature of the interaction.

ful excitations, let us recall that it is possible to produce quasi-particle excitations upon the Laughlin state [18]. One of them is the quasi-hole (vortex) excitation, $Q_{\mathbf{p}} [e^{i\mathbf{p} \cdot \mathbf{r}}] \sum_{i < j} (\mathbf{r}_i - \mathbf{r}_j)^2$, and another one is the 'quasi-boson' (anti-vortex) excitation, $Q_{\mathbf{p}} [e^{i\mathbf{p} \cdot \mathbf{r}}] \sum_{i < j} (\mathbf{r}_i - \mathbf{r}_j)^2$, where \mathbf{r}_0 denotes the location of the vortex or that of the anti-vortex. Then the lowest-lying neutral excitations are considered to be the bound states of a vortex-antivortex pair.

By investigating a system with $2S = 2N - 3$ or $2S = 2N - 1$, we find that in the thermodynamic limit the quasi-boson creation energy remains finite (1.248 ± 0.007 in units of g) [20], while the quasi-hole creation energy is zero for repulsive contact interaction. Since each of quasi-boson and quasi-hole excitations has AM $N=2$, the branch composed of a quasi-boson-quasi-hole pair is expected to end at $L_{\text{tot}} = N$. This is indeed the case in the lowest-lying excitation branch as shown in Fig. 4, where the branches for system size $N = 4; 5; 6; 7$ and 8 are plotted against 'wave number' $k = L_{\text{tot}}/R$.

In Fig. 4 it is seen that the minimum of the neutral-excitation energy (indicated by closed symbols) is realized at $k_{\text{min}} \sim 3(M/\hbar)^{1/2}$. For $k > k_{\text{min}}$, the excitation energy increases with increasing k and approaches the quasi-boson creation energy (indicated on the right side of the figure by a line for each N). As shown in the




FIG. 4: The lowest-lying rotonlike excitation branch is shown against wave number $k = L_{\text{tot}}/R$ for $N = 4; 5; 6; 7$ and 8 . The minimum of the neutral-excitation energy is realized at $k_{\text{min}} \propto (M/\hbar)^{1/2}$ (indicated by closed symbols), and each branch ends at $L_{\text{tot}} = N$ as expected for a quasiboson pair excitation. For $k < k_{\text{min}}$, the excitation energy increases with increasing k and approaches the quasiboson creation energy (indicated on the right side of the figure by a line for each N). In the inset, the quasiboson creation energy and the minimum value of the neutral-excitation energy are shown against $1/(N-1)$ by open and closed circles, respectively.

inset of Fig. 4, the difference between the quasiboson creation energy (indicated by open circles) and the minimum value of the neutral-excitation energy (indicated by closed circles) decreases as the system size increases and may vanish in the thermodynamic limit. In the presence of this finite energy gap, however, this rotonlike minimum can survive even for very large values of N by introducing, e.g., laser-induced dipole-dipole interactions between bosons [21].

It should be noted that the lowest-lying spectrum has no phonon branch, despite the short-range nature of the interaction [22]. A similar roton spectrum has been reported in the fractional quantum Hall system [23], where absence of the phonon branch has been attributed to the Higgs mechanism caused by the long-range nature of the Coulomb interaction. However, the rotonlike structure with no phonon branch found in this Letter for the contact interaction does not fit into this category. In the composite-boson picture, the $\nu = 1/2$ Laughlin state is nothing but a condensate of composite bosons, each of which is composed of a boson and two flux quanta attached to it. In the composite-particle picture for the quantum Hall system, it has been discussed that interactions between composite particles are different from those between original particles [24]. It might be conceived that the finite gap in the long-wavelength limit

is related with momentum-dependent long-range interactions introduced on the composite-boson transformation (without being induced by a laser [21]). However, the understanding of the gap formation found in this Letter remains to be clarified and merits further investigation.

T.N. and M.J. acknowledge support by Grant-in-Aids for Scientific Research (Grant No.14740181 and No.15340129) by the Ministry of Education, Culture, Sports, Science and Technology of Japan, respectively. M.J. also acknowledges support by the Toray Science Foundation and by the Yamada Science Foundation.

-
- [1] P.C. Haljan et al, Phys. Rev. Lett. 87, 210403 (2001); P. Engels et al, *ibid.* 89, 100403 (2002).
 - [2] J.R. Abo-Shaeer et al, Science 292, 476 (2001).
 - [3] P. Rosenbusch et al, Phys. Rev. Lett. 88, 250403 (2002).
 - [4] E. Hodby et al, Phys. Rev. Lett. 88, 010405 (2002).
 - [5] N.K.W. Ilkin et al, Phys. Rev. Lett. 80, 2265 (1998).
 - [6] N.K.W. Ilkin and J.M.F. Gunn, Phys. Rev. Lett. 84, 6 (2000).
 - [7] T.-L. Ho, Phys. Rev. Lett. 87, 060403 (2001).
 - [8] B. Mottelson, Phys. Rev. Lett. 83, 2695 (1999).
 - [9] I. Hamamoto, in *Treatise on Heavy-Ion Science*, edited by D.A. Bromley (Plenum, New York, 1985), vol.3, p.313; I. Hamamoto and B. Mottelson, Nucl. Phys. A 507, 65 (1990).
 - [10] S. Inouye et al, Nature 392, 151 (1998); S.L. Comish et al, Phys. Rev. Lett. 85, 1795 (2000).
 - [11] G.F. Bertsch and T. Papenbrock, Phys. Rev. Lett. 83, 5412 (1999).
 - [12] T. Papenbrock and G.F. Bertsch, Phys. Rev. A 63, 023616 (2001).
 - [13] T. Nakajima and M. Ueda, Phys. Rev. A 63, 043610 (2001).
 - [14] M. Ueda and T. Nakajima, Phys. Rev. A 64, 063609 (2001).
 - [15] G.M. Kavoulakis et al, Phys. Rev. A 63, 055602 (2001).
 - [16] V. Bardek et al, Phys. Rev. A 64, 015603 (2001); V. Bardek and S.M. Eljanac, *ibid.* 65, 013602 (2002).
 - [17] G.M. Kavoulakis et al, Phys. Rev. A 62, 063605 (2000).
 - [18] R.B. Laughlin, Phys. Rev. Lett. 50, 1395 (1983).
 - [19] F.D.M. Haldane, Phys. Rev. Lett. 51, 605 (1983); G. Fano et al, Phys. Rev. B 34, 2670 (1986).
 - [20] The best fit result for the neutral excitation gap is $1.229 + 2.394/(N-1) + 0.2476/(N-1)^2$ in units of g , while the one for the quasiboson creation energy is $1.248 + 2.367/(N-3) + 0.2279/(N-3)^2$.
 - [21] D.H.J. O'Dell et al, Phys. Rev. Lett. 90, 110402 (2003); L. Santos et al, *ibid.* 90, 250403 (2003).
 - [22] In an unpublished work by N. Regnault and Th. Jolicoeur (cond-mat/0212477), a related subject on the finiteness of neutral-excitation energy gap is discussed.
 - [23] S.M. Girvin et al, Phys. Rev. Lett. 54, 581 (1985).
 - [24] G. Dev and J.K. Jain, Phys. Rev. Lett. 69, 2843 (1992); T. Nakajima and H. Aoki, *ibid.* 73, 3568 (1994). On the theory of composite particle as a dipole, see R. Shankar and G. Murthy, *ibid.* 79, 4437 (1997) and references therein.