

Dissipationless shock waves in repulsive Bose-Einstein condensates

A. M. Kamchatnov¹, A. Gammal², and R. A. Kraenkel³

¹Institute of Spectroscopy, Russian Academy of Sciences,
Troitsk 142190, Moscow Region, Russia

²Instituto de Física, Universidade de São Paulo,
05315-970, C. P. 66318 São Paulo, Brazil

³Instituto de Física Teórica, Universidade Estadual Paulista-UNESP
Rua Pamplona 145, 01405-900 São Paulo, Brazil

April 14, 2024

Abstract

We consider formation of dissipationless shock waves in Bose-Einstein condensates with repulsive interaction between atoms. It is shown that big enough initial inhomogeneity of density leads to wave breaking phenomenon followed by generation of a train of dark solitons. Analytical theory is confirmed by numerical simulations.

Experiments on free expansion of Bose-Einstein condensate (BEC) have shown [1] that evolution of large and smooth distributions of BEC is described very well by hydrodynamic approximation [2] where dispersion and dissipation effects are neglected. At the same time, it is well known from classical compressible gas dynamics (see, e.g. [3]) that typical initial distributions of density and velocity can lead to wave breaking phenomenon when formal solution of hydrodynamical equations becomes multiple-valued. It means that near the wave breaking point one cannot neglect dispersion and/or dissipation effects which prevent formation of a multiple-valued region of a solution. If the dissipation effects dominate the dispersion ones, then the multiple-valued region is replaced by the classical shock, i.e. narrow layer with strong dissipation within, which separates smooth regions with different values of density, fluid velocity and other physical parameters. This situation was studied in classical gas dynamics and found many practical applications. If, however, the dispersion effects dominate dissipation ones, then the region of strong oscillations is generated in vicinity of wave breaking point [4, 5]. Observation of dark solitons in BEC [6, 7, 8] shows that the main role in dynamics of BEC is played by dispersion and nonlinear effects taken into account by standard Gross-Pitaevskii (GP) equation [9], and dissipation effects are relatively small and can be considered as perturbation. Hence, there are initial distributions of BEC which can lead to formation of dissipationless shock waves. Here we shall consider such possibility.

The starting point of our consideration is the fact that the sound velocity in BEC is proportional to the square root from its density (see, e.g. [9] and references therein). Thus,

if we create inhomogeneous BEC with high density hump (with density n_1) in the center of lower density distribution (with density n_0), and after that release this central part of BEC, then the high density hump will tend to expand with velocity $\sqrt{2} \frac{v_1}{v_0}$ greater than the sound velocity $\frac{v_1}{v_0}$ of propagation of disturbance in lower density BEC. As a result, wave breaking and formation of dissipationless shock wave can occur in this case. Note that initial distribution of this type was realized in the recent experiment [10] where generation of oscillations was also observed. In this experiment the lower density disk-shaped BEC was confined by magnetic trap and density distribution had standard Thomas-Fermi (TF) parabolic form. Additional potential was applied to BEC in the central part of above TF profile which lead to narrow parabolic hump in the density distribution. After the central potential was switched off, the hump started to expand against wide lower density TF profile leading to generation of oscillations in the transition region between high and low density condensates. The theory of dissipationless shock waves in media described by one-dimensional (1D) nonlinear Schrödinger (NLS) equation was developed in [11, 12]. Since the GP equation in some cases can be reduced to the 1D NLS equation, this theory can be applied to description of dissipationless shock waves in BEC. Here we suggest such description and confirm it by numerical simulations.

We consider BEC confined in a "pancake" trap with the axial frequency ω_z much greater than the transverse one $\omega_x = \omega_y = \omega_\perp$. Let the density of atoms in the central part of BEC be of order of magnitude n_0 and satisfy the condition $n_0 a_s a_z^2 \gg 1$, where $a_s > 0$ is the s-wave scattering length and $a_z = (\hbar/m a_z)^{1/2}$ is the amplitude of quantum oscillations in the axial trap. Then the condensate wavefunction can be factorized as $\psi = \psi(z) \phi(x; y)$, where $\psi(z) = \frac{1}{\sqrt{2} a_z^{1/2}} \exp(-z^2/2a_z^2)$ is the ground state wavefunction of axial motion, and $\phi(x; y; t)$ satisfies the reduced 2D GP equation

$$i\hbar \partial_t \phi = -\frac{\hbar^2}{2m} (\partial_{xx} + \partial_{yy}) \phi + V(x; y) \phi + g_{2D} |\phi|^2 \phi; \quad (1)$$

where $V(x; y)$ is the potential of a transverse trap, $g_{2D} = \frac{\hbar^2}{2m} \frac{1}{2} a_s^{-2} a_s = (\hbar a_z)^{-1}$ is the effective nonlinear interaction constant, and ϕ is normalized to the number of atoms, $\int |\phi|^2 dx dy = N$. The initial distribution of density is determined by the potential $V(x; y)$ and consists of wide background with a hump in its center. We assume that the background width is much greater than the hump's width, so that at the initial stage of evolution we can consider an expansion of the central part against the constant background. In a similar way, at the initial stage of evolution, when the radius of the central part does not change considerably, we can neglect the curvature of axially symmetrical distribution and consider its 1D cross section. It means that we can neglect the dependence of ϕ on y coordinate and consider ϕ as a function of x and t only. As a result, we arrive at 1D NLS equation with inhomogeneous initial distribution of density. To simplify the notation, we introduce dimensionless variables $t^0 = \omega_z t/2$, $x^0 = x/a_z$; $u = (\frac{\hbar^2}{2m} a_s a_z^{-1} n_0)^{1/2}$. Then the initial stage of evolution of the wavefunction profile in the x axis cross section is governed by the NLS equation,

$$iu_t + u_{xx} - 2ju^2u = 0; \quad (2)$$

where the primes in t^0 and x^0 are omitted for convenience of the notation.

Evolution of smooth pulses before the wave breaking point can be described in the hydrodynamic approximation which can be achieved by substitution

$$u(x;t) = \frac{p}{(x;t)} \exp \left(i \int_{x^0}^x v(x^0;t) dx^0 \right) \quad (3)$$

and separation of the real and imaginary parts. As a result we obtain the system

$$\frac{1}{2} \dot{t} + (v)_x = 0; \quad \frac{1}{2} \dot{v}_t + vv_x + \frac{1}{2} \ddot{x} = 0; \quad (4)$$

where we have neglected the so-called "quantum pressure" term with higher space derivatives what is correct until the density distribution has smooth enough profile. To get simple qualitative picture of the wave breaking of BEC density distribution, let us consider idealized case with a box-like hump in the initial distribution,

$$\rho(x;0) = \begin{cases} \rho_0; & |x| \leq a; \\ 0; & |x| > a; \end{cases} \quad (5)$$

Although this distribution has a parameter with dimension of length $|width\ of\ the\ hump|$ it does not play any role for $t \leq a/(2^{p-1})$, i.e. until two waves propagating inward the hump with sound velocity $c_1 = 2^{p-1}$ meet at $x = 0$. Hence, for this initial period of evolution the solution for $x > 0$ can only depend on the self-similar variable $\xi = (x - a)/t$, centered at $x = a$, and for $x < 0$ on $\xi = (x + a)/t$ centered at $x = -a$. Since the picture is symmetrical, it is enough to consider only a half of the solution corresponding to $x > 0$. It is easy to find that the density is given by the formulae

$$\rho(x;t) = \begin{cases} \rho_0 & \text{for } 0 < x < a - 2^{p-1}t; \\ 2^{p-1} \left(\frac{x-a}{t} \right)^2 = 9 & \text{for } a - 2^{p-1}t < x < a + 2^{p-1}t; \\ 2^{p-1} + \left(\frac{x-a}{t} \right)^2 = 9 & \text{for } a + 2^{p-1}t < x < 2a + 2^{p-1}t; \\ 0 & \text{for } x > a + 2^{p-1}t; \end{cases} \quad (6)$$

and similar formulae can be written for the velocity field $v(x;t)$. This solution describes the wave breaking phenomenon which takes place if $\rho_1 > 4\rho_0$. The density profile shown in Fig. 1 clearly illustrates the origin of the multiple-valued region which should be replaced by the oscillatory shock wave when the dispersion effects are taken into account.

For more realistic initial pulses the density profile is a smooth function without cusp points. In vicinity of the wave breaking point the solution can be approximated by a cubic function for one Riemann invariant $\psi_+ = v/2 + \frac{p}{2}$ of the system (4) and by constant value for another one $\psi_- = v/2 - \frac{p}{2}$ (see [11, 12]). After Galileo and scaling transformations the hydrodynamic solution can be written in the form

$$x - (3\psi_+ + \psi_-)t = \frac{3}{4}; \quad \psi_- = \text{const}; \quad (7)$$

and again for $t > 0$ it has a multiple-valued region of ψ_+ . It means that dispersion effects have to be taken into account which lead to formation of dissipationless shock wave after wave breaking point.

In framework of Whitham theory of modulations [13, 5] one can obtain an approximate solution of the NLS equation (2) in analytic form where the dissipationless shock wave is

presented as a modulated periodic nonlinear wave solution of the NLS equation. The density is expressed in terms of Jacobi elliptic function

$$\rho(x,t) = \rho_0(x,t) \left[1 + \frac{1}{4} (\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + \frac{1}{4} (\alpha_1 \alpha_2 + \alpha_3 \alpha_4) \operatorname{sn}^2 \left(\frac{x}{L} \sqrt{\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2}; m \right) \right] \quad (8)$$

where

$$x = x_0 + (\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)t; \quad m = \frac{(\alpha_1^2 + \alpha_2^2)(\alpha_3^2 + \alpha_4^2)}{(\alpha_1^2 + \alpha_3^2)(\alpha_2^2 + \alpha_4^2)}; \quad (9)$$

and parameters $\alpha_i, i = 1; 2; 3; 4$ change slowly along the dissipationless shock. Their dependence on x and t is determined implicitly by the solution

$$x - v_i(t) = w_i(\eta); \quad i = 1; 2; 3; \quad \alpha_4 = \bar{\alpha}_4 = \text{const} \quad (10)$$

of Whitham equations, where Whitham velocities v_i and w_i are given by quite complicated expressions in terms of elliptic integrals (see [11, 12]):

$$w_i = \frac{8}{35} w_i^{(3)} + \frac{4}{5} \bar{w}_i^{(2)} - \frac{1}{35} v_i^2 + \frac{1}{35} \bar{v}_i^3; \quad i = 1; 2; 3; \quad (11)$$

$$w_i^{(k)} = W^{(k)} + (v_i - s_i) \partial_i W^{(k)}; \quad (12)$$

$$W^{(1)} = V = s_1; \quad W^{(2)} = \frac{3}{8} s_1^2 - \frac{1}{2} s_2; \quad W^{(3)} = \frac{5}{16} s_1^3 - \frac{3}{4} s_1 s_2 + \frac{1}{2} s_3; \quad (13)$$

$$v_i(\eta) = 1 - \frac{L}{\partial_i L} \partial_i V; \quad \partial_i = \partial / \partial \eta; \quad i = 1; 2; 3; 4; \quad (14)$$

where

$$L = \frac{2K(m)}{(\alpha_1^2 + \alpha_3^2)(\alpha_2^2 + \alpha_4^2)} \quad (15)$$

is a wavelength, $K(m)$ is the complete elliptic integral of the first kind, and s_1, s_2, s_3 are determined by the expressions

$$s_1 = \sum_i X_i; \quad s_2 = \sum_{i < j} X_{ij}; \quad s_3 = \sum_{i < j < k} X_{ijk}; \quad (16)$$

Equations (11) can be solved with respect to $\alpha_i, i = 1; 2; 3$ giving them as functions of x and t . Subsequent substitution of these functions $\alpha_i(x,t), i = 1; 2; 3$ into (8) yields the modulated periodic wave which represents the dissipationless shock wave. The resulting profile of density in dissipationless shock wave is shown in Fig. 2. At one its edge it consists of the train of dark solitons, and at another edge describes small amplitude oscillations propagating with local sound velocity into unperturbed region described by smooth solution of hydrodynamical equations. The modulated periodic wave replaces the multiple-valued

region shown by dashed line which was obtained in hydrodynamic approximation after the wave breaking point. This multiple-valued region is analogous to that in Fig. 1 with account of change of variables due to Galileo and scaling transformations.

To check the described above picture of formation of dissipationless shock wave and to extend it to real 2D situation, we have solved numerically the 2D generalization of Eq. (2),

$$iu_t + u_{xx} + u_{yy} - 2ju^2u = 0; \quad (17)$$

with the initial condition

$$(u; 0) = \begin{cases} u_0 + (u_1 - u_0)(1 - r^2/a^2); & |r| \leq a; \\ 0; & |r| > a; \end{cases} \quad (18)$$

(where $r = (x^2 + y^2)^{1/2}$) similar to one studied experimentally [10]. Plot of two-dimensional initial density distribution is shown in Fig. 3 and density distribution after time evolution $t = 2$ is shown in Fig. 4. As we see, the parabolic hump expands with formation of dissipationless shock wave in the transition layer between high density region to low density one. To see more clearly the evolution of the hump, its cross section profiles are shown in Fig. 5 at different values of time t . Slowly propagating dark solitons are clearly seen as well as small amplitude sound waves propagating into undisturbed low density region. Dissipationless shock wave generated at the right side of the profile coincides qualitatively with results of analytic theory shown in Fig. 2. One may assume that oscillations in BEC density profile observed in experiment [10] have the same origin.

Further evolution of the hump will ultimately lead to its spreading over large area with small amplitude oscillations, that is the shock wave does not persist permanently. Rather, it is a transient phenomenon caused by different values of characteristic velocities in high density hump and low density background. Slowly propagating dark solitons are generated in the transient layer between these two regions in order to reconcile two different values of velocities of disturbance propagation. This mechanism of dark soliton generation is quite general and can manifest itself in various geometries and initial BEC distributions¹.

In conclusion, we have studied theoretically and numerically the process of formation of dissipationless shock waves in the density distribution of BEC. We believe that oscillations in BEC density profile observed in recent experiment [10] can be explained by this theory.

This work was supported by FAPESP (Brazil) and CNPq (Brazil). A.M.K. thanks also RFBR (grant 01-01-00696) for partial support.

References

- [1] U. Ernst, A. Marte, F. Schreck, J. Schuster, and G. Rempe, *Europhys. Lett.* 41, 1 (1998).
- [2] Y. Castin and R. Dum, *Phys. Rev. Lett.* 77, 5315 (1996).
- [3] L.D. Landau and E.M. Lifshitz, *Hydrodynamics*, 4th ed. (Nauka, Moscow, 1988).

¹Formation of dark soliton trains in 1D geometry was considered in recent preprint [14]

- [4] A.V. Gurevich and L.P. Pitaevskii, Zh. Exp. Teor. Fiz. 65, 590 (1973) [Sov. Phys. JETP, 38, 291 (1973)].
- [5] A.M. Kamchatnov, Nonlinear Periodic Waves and Their Modulations (World Scientific, Singapore, 2000).
- [6] S. Burger, K. Bongs, S. Dettmer, W. Ertmer, K. Sengstock, A. Sanpera, G.V. Shlyapnikov, and M. Lewenstein, Phys. Rev. Lett. 83, 5198 (1999).
- [7] J. Denschlag, J.E. Sinsarian, D.L. Feder, Ch.W. Clark, L.A. Collins, J. Cubizolles, L. Deng, E.W. Hagley, K. Helmerson, W.P. Reinhardt, S.L. Rolston, B.J. Schneider, and W.D. Phillips, Science, 287, 97 (2000).
- [8] B.P. Anderson, P.C. Haljan, C.A. Regal, D.L. Feder, L.A. Collins, C.W. Clark, and E.A. Comell, Phys. Rev. Lett. 86, 2926 (2001).
- [9] L.P. Pitaevskii and S. Stringari, Bose-Einstein Condensation (Clarendon Press, Oxford, 2003).
- [10] E.A. Comell, talk at NATO Advanced Workshop "Nonlinear Waves: Classical and Quantum Aspects", Lisbon, 2003.
- [11] A.M. Kamchatnov, in Nonlinearity and Disorder: Theory and Applications, eds. F. Abdullaev, O. Bang and P.O. Sorensen, Kluwer, Boston, 2001, p. 255.
- [12] A.M. Kamchatnov, R.A. Kraenkel, and B.A. Umrov, Phys. Rev. E 66, 036609 (2002).
- [13] G.B. Whitham, Linear and Nonlinear Waves, Wiley, N.Y., 1974.
- [14] B. Damski, preprint cond-mat/0309421.

Figures captions

Fig. 1. Wave breaking of a box-like initial density distribution (shown by dashed line) in hydrodynamic approximation. The plot corresponds to the solution (6) with $a = 10$; $\rho_0 = 1$; $\rho_1 = 10$; $t = 1$. Point A propagates inward the box with local sound velocity $2\sqrt{\rho_1}$, point C propagates outward along the background density with local sound velocity $2\sqrt{\rho_0}$, and point B corresponds to intersection of two simple wave solutions with profiles AB and CB.

Fig. 2. Formation of dissipationless shock wave after wave breaking point according to Whitham modulation theory applied to 1D NLS equation. Dashed line corresponds to multiple-valued region arising in hydrodynamic approximation given by Eq. (7) (it is analogous to the region between the points B and C in Fig. 1), and solid line represents modulated periodic wave given by Eqs. (8) and (10). Both profiles are calculated for $\bar{\rho} = 10$ at time $t = 1$.

Fig. 3. Two-dimensional initial distribution of BEC density with paraboloid hump on constant background given by Eq. (18) with $a = 10$; $\rho_0 = 1$; $\rho_1 = 10$.

Fig. 4. Two-dimensional density distribution of BEC after time $t = 2$ of evolution from initial paraboloid density on constant background according to numerical solution of 2D NLS equation.

Fig. 5. Cross sections of density profile at different evolution time according to numerical solution of 2D NLS equation (17) with initial condition (18).

This figure "fig1.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/cond-mat/0310457v1>

This figure "fig2.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/cond-mat/0310457v1>

This figure "fig3.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/cond-mat/0310457v1>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/cond-mat/0310457v1>

This figure "fig5.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/cond-mat/0310457v1>