

Dynamics of laser-induced lattice formation in Bose-Einstein condensates

M. Kalinski¹, I.E. Mazets^{2,3}, G. Kurizki², B.A. Malomed⁴, K. Vogel¹ and W.P. Schleich¹¹ Abteilung für Quantenphysik, Universität Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany,² Department of Chemical Physics, Weizmann Institute of Science, Rehovot 76100, Israel,³ Io e Physico-Technical Institute, St.Petersburg 194021, Russia,⁴ Department of Interdisciplinary Studies, Faculty of Engineering, Tel-Aviv University, Tel-Aviv 69978, Israel

We present variational and numerical solutions of the generalized Gross-Pitaevskii equation for an atomic Bose-Einstein condensate with time-dependent, laser-induced dipole-dipole interactions (LDDI). The formation of a spatially modulated ground state due to LDDI in an elongated (cigar-shaped) BEC is investigated. We find that modulated densities can emerge if the laser intensity exceeds a critical value, which depends on the linear density of atoms in the BEC.

PACS number: 03.75.Kk

I. INTRODUCTION

The confinement of atomic Bose-Einstein condensates (BECs) in optical traps [1,2] and lattices [3] is commonly provided by a spatial inhomogeneity of the laser intensity, which gives rise to gradient (or dipole) forces acting on single atoms. Completely different dynamical effects are generated in BECs by interatomic laser-induced dipole-dipole interactions (LDDI) [4,5]. These effects take place even if the atoms are illuminated by a plane electromagnetic wave. One such effect, predicted by means of a 1D variational ansatz, is that LDDI caused by an off-resonant circularly polarized plane wave traveling along a cigar-shaped BEC results in a spatially periodic density modulation [5]. In the present paper we discuss how such density modulations can emerge from a uniform BEC when the laser intensity reaches sufficiently large values.

A. Supersolids versus BECs in optical lattices

It is tempting to refer to such a density-modulated BEC as a supersolid. Indeed, Ref. [5] used this term to emphasize the presence of both spatial modulation and phase coherence. Moreover, there exists also close connection to a BEC in an optical lattice [3]. For this reason we dedicate this subsection to a comparison of our density-modulated BEC to supersolids and BECs in optical lattice.

Superfluid systems with spatially modulated densities have a long history. The original question, however, was not whether a homogeneous BEC can be spatially modulated. Instead, the discussion started from the question whether a solid composed of bosonic atoms can be superfluid. First, Penrose and Onsager [6] demonstrated that Bose-Einstein condensation is impossible in an ideal crystal. However, later it became clear [7,8] that if there were vacancies present in a crystal then Bose-Einstein condensation would be possible. The work of Chester [9] and

Leggett [10] led to a deeper understanding of Bose-condensed solids. However, the authors of Refs. [6-10] concentrated on crystals where the occupation number of a lattice site never exceeds 1. Thus the estimation for the ratio between the superfluid density and the normal density in such a system was rather pessimistic ($\sim 10^{-4}$) [10]. In contrast, the situation for liquid ^4He is more favorable. For example, Pitaevskii [11] predicted the formation of a standing wave in helium (below the critical temperature) owing to a supercritical velocity in a capillary tube. The systems studied in the references mentioned above are quite different from a Bose-condensed dilute gas.

However, the studies of Poméau and Rica [12] and Goral et. al [13] are closer to our work. Poméau and Rica analyzed what could happen if the roton minimum in the liquid helium spectrum reaches the horizontal axis and even if the homogeneous system would become unstable at the corresponding wavelength. Their finding is that as soon as the uniform ground state becomes unstable, another ground state, spatially modulated, comes to existence, instead. Poméau and Rica call such a density-modulated state "supersolid", an expression already used by Cheng [8].

Goral et. al discussed quantum phases of dipolar bosons in an optical lattice. Their system consists of a dilute atomic gas where the atoms interact not only via the short-range s-wave scattering, but also { due to their large magnetic dipole moments } via long-range dipole-dipole interaction. This dipole-dipole interaction is somewhat similar to the LDDI, except that the magnetic dipole-dipole interaction is purely static, varies as the inverse distance cubed, and does not exhibit oscillatory (retardation) features, which are essential in considerations of the LDDI [4,5]. They found that in an optical lattice such a system can have various phases, including a supersolid phase. We conclude our brief discussion of supersolids by mentioning that recently a supersolid phase of ^4He may have been observed [14].

Another superfluid system that displays spatially modulated densities is a BEC in an optical lattice. Here up to several hundred atoms can occupy one lattice site. As first demonstrated by Anderson and Kasevich [15], macroscopic quantum interference can be observed in such systems. As we show in the present paper, in our system many atoms contribute to a maximum in the density. Therefore, we expect our system to have similar coherence properties as a BEC in an optical lattice.

Due to its analogy to BECs in optical lattices we may call the effect considered in the present paper a spontaneous formation of an optical lattice in a laser-illuminated BEC. Such a lattice is self-sustained, since it is formed by interfering incident and coherently backward scattered light waves.

B. Formulation of the problem

While Ref. [5] has suggested the intriguing possibility of "spontaneous laser-induced periodicity" in a BEC, it has left open basic questions related to their observability: (a) Is the suggested 1D variational steady-state solution suitable for a cigar-shaped BEC, or should we expect radial density modulation as well? (b) Is a spatially modulated ground state (the signature of spontaneous laser-induced periodicity) obtainable from a uniform initial ground state by gradually increasing the laser intensity, even though it must compete with evaporation via Rayleigh scattering and possible non-adiabatic effects? (c) Is this density modulation distinguishable from that obtained by the superradiant response of a BEC [16]?

In this work we aim at a realistic investigation of laser-induced formation of spatially modulated densities in a BEC that would provide clear answers to the questions raised above. This is accomplished through an exhaustive numerical investigation that fully accounts, for the first time, for the 3D geometry of the cigar-shaped sample and the time-dependent dynamics of the formation process. Our numerical results show that in the proposed setting spontaneous laser-induced periodicity is experimentally feasible. They are supplemented by a simple variational approximation, which is shown to explain the essential features of the numerical results and their stability. We also compare and contrast the differences of the present LDDI effects to superradiant light scattering by a BEC [16].

C. Outline of the paper

Our paper is organized as follows: In Sec. II we introduce a Gross-Pitaevskii equation which takes into account the laser-induced dipole-dipole interaction and discuss the interplay between the long-range dipole-dipole

interaction and the short-range s-wave scattering. Section III is devoted to a variational approach to the problem. We assume a cigar-shaped BEC with a tight radial confinement and arrive at a one-dimensional equation. We use variational calculations to obtain approximate stationary solutions of this equation. In Sec. IV we solve the full Gross-Pitaevskii equation numerically and find time-dependent as well as stationary solutions. Based on these results we discuss a possible experiment in Sec. V. Furthermore, we comment on the depletion of the condensate in such an experiment and mention other strategies to obtain spatially modulated densities. We also compare density-modulated BECs formed by LDDI to coherent atom recoil lasing. We conclude by summarizing our results in Sec. VI.

II. GROSS-PITAEVSKII EQUATION

The time evolution of the BEC order parameter is governed by the 3D Gross-Pitaevskii equation

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r};t) = \frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r};t) + V_{\text{trap}}(\mathbf{r}) \psi(\mathbf{r};t) + \frac{4\hbar^2 a_s}{Zm} |\psi(\mathbf{r};t)|^2 \psi(\mathbf{r};t) + \int d^3\mathbf{r}' |\psi(\mathbf{r}';t)|^2 V_{\text{dd}}(\mathbf{r} - \mathbf{r}') \psi(\mathbf{r};t); \quad (1)$$

Here

$$V_{\text{trap}} = \frac{m}{2} \frac{\hbar^2}{m^2} (x^2 + y^2) + \frac{1}{2} \frac{\hbar^2}{m^2} z^2 \quad (2)$$

denotes the external trapping potential and m and a_s are the atomic mass and the s-wave scattering length due to short-range interatomic forces, respectively. The last term in Eq. (1), which comes in addition to the terms of the usual Gross-Pitaevskii equation [17,18], accounts for the long-range LDDI. Note that $\psi(\mathbf{r};t)$ is normalized to the number of atoms.

For a circularly-polarized laser wave propagating along the z axis, the kernel of this effective interaction reads [5]

$$V_{\text{dd}}(\mathbf{r}) = U_0 \left[3 \cos^2 \theta - 1 \right] \frac{\cos qr + qr \sin qr}{(qr)^3} \cos^2 \theta + 1 \frac{\cos qr}{qr} \cos(qr \cos \theta); \quad (3)$$

Here, the direction of the interatomic separation vector \mathbf{r} relative to the z axis is determined by the polar and azimuthal angles θ and ϕ , respectively, and q denotes the laser wave number. The characteristic energy U_0 corresponding to the interaction of two atoms separated by $qr = 1$ is proportional to the laser intensity. On the other hand, the rate Γ_{Ray} of the incoherent (Rayleigh) photon scattering per atom is also proportional to the laser intensity which provides us with the connection $U_0 = \frac{3}{4} \hbar \Gamma_{\text{Ray}}$.

From the outset, we must address two questions: (i) Can we use in Eq. (1) the bare potential $V_{dd}(r)$ of Eq. (1)? (ii) To what extent does the interplay between the long-range potential $V_{dd}(r)$ and the short-range interaction potential change the value of a_s in Eq. (1)?

To answer the first question we recall that according to Belyaev's theory of effective interactions [19], the bare potential should be replaced by an effective potential, such that the scattering amplitude produced by the effective potential in the first Born approximation coincides with the exact scattering amplitude generated by the bare potential. We have performed such a calculation for the isotropic part $V_i(r)$ of the LDDIP potential, which has the asymptotic forms $V_i(r) \sim \frac{1}{2} (qr)^{-2} \sin(2qr)$ for $qr \ll 1$, and $V_i(r) \sim (22=15) \frac{1}{2} (qr)^{-1}$ for $qr \gg 1$ [20]. We have found, that, for this isotropic part, the Born scattering amplitude is always close to the exact one. We conjecture that the same is true for the entire (anisotropic) potential $V_{dd}(r)$.

To address the second question we first recall [21] that the part of V_{dd} being proportional to $(3 \cos^2 \theta - 1)r^{-3}$ has no significant influence on a_s as long as we are far from the shape resonance. In contrast, Ref. [22] shows that the addition of the Coulomb interaction to a short-range potential, may dramatically change the s-wave scattering properties. Since for short distances the isotropic part of V_{dd} has Coulomb character, this contribution may have some impact on the s-wave scattering length.

To clarify this issue, we have numerically calculated the phase of the scattered wave, combining the isotropic part of V_{dd} and the short-range potential. The latter has been taken into account by the boundary condition $(r)^0 = (r)^1; 0 = 1 = a_s$, for the solution (r) of the radial Schrödinger equation describing the scattering of two atoms. We have not found any dramatic change of a_s throughout the relevant range of relative momenta of the colliding atoms. This result is due to the fact that the gravity-like part of $V_{dd}(r)$ has a cutoff at $r \sim q^{-1}$, which is much shorter than the corresponding Bohr's radius [4] for experimentally feasible laser intensities. Thus, we may use, with reasonable accuracy, the same value of a_s as without the laser field.

III. VARIATIONAL CALCULATIONS

Before we solve Eq. (1) numerically we first want to gain a qualitative understanding of the dynamics of the system described by Eq. (1). For this purpose we have developed a quasi-1D variational approximation which we discuss in this section. This approximation already predicts spontaneous laser-induced periodicity, that is, a spatially modulated ground state, when the laser intensity exceeds a threshold value.

We start from the fact that Eq. (1) can be derived from the Lagrangian $L = \int d^3r L$ with

$$L = \frac{i\hbar}{2} \left[\psi(r;t) \frac{\partial}{\partial t} \psi(r;t) - \psi(r;t) \frac{\partial}{\partial t} \psi(r;t) \right] - \frac{\hbar^2}{2m} |\nabla \psi(r;t)|^2 - V_{\text{trap}}(r) |\psi(r;t)|^2 - \frac{2\hbar^2 a_s}{2m} |\psi(r;t)|^4 - \frac{1}{2} \int d^3r^0 |\psi(r^0;t)|^2 V_{dd}(r - r^0) |\psi(r;t)|^2 : \quad (4)$$

We assume a cigar-shaped condensate where the radial confinement is so tight that, regardless of the atomic density, only the ground state of the transverse motion is occupied. This assumption allows us to approximate [23] the transverse distribution of the order parameter by

$$\psi(r;t) = \left(\frac{P}{\pi} \right)^{1/2} \exp[-(x^2 + y^2)/(2 \frac{P}{\pi})] \psi(z;t) \quad (5)$$

with $P = \frac{P}{h^2 m \omega_z}$. This approximation reduces the full Gross-Pitaevskii equation to a one-dimensional equation for the longitudinal part $\psi(z;t)$.

Since for

$$|\psi|^2 \sim q^{-1} > \frac{P}{x^2 + y^2} \quad (6)$$

the dipole-dipole potential takes the form

$$V_{dd}(r) \sim 2 \frac{1}{q} \cos^2(qz) \quad (7)$$

we expect a spatial structure with the period $\lambda = q$ to appear in the ground state. Furthermore, we assume for our variational approximation that the trapping potential $V_{\text{trap}}(r)$ does not confine the BEC in z-direction, that is we assume $\omega_z = 0$. We therefore pursue the 1D variational ansatz

$$\psi(z;t) = a(t) \cos(2qz) + c(t); \quad (8)$$

where $a(t)$ and $c(t)$ are complex variables.

To stabilize the BEC against long-wavelength perturbations [5], we must also require that the short-range interaction, characterized by a positive scattering length, $a_s > 0$, is stronger than the net laser-induced attraction of two atoms separated by $r < q^{-1}$. Moreover, we argue that this short-range repulsion term does not affect the dynamics at the wavelength $\lambda = q$, since, for moderate BEC densities, $\lambda = q$ is smaller than the BEC healing length, which is determined by the BEC density and a_s . Only under these conditions the variational ansatz Eq. (8) describes the dynamics of the condensate appropriately.

We now derive equations of motion for the coefficients $a(t)$ and $c(t)$. For this purpose we substitute Eq. (5) with Eq. (8) into the Lagrangian density Eq. (4) and perform the integration. Here we consider a "cigar" of size ℓ in the z -direction and neglect interaction terms that do not contain the large logarithmic factor $2 \log[\ell/(2\lambda)]$, which diverges as $\ell \rightarrow 1$. The effective Lagrangian density

$$\frac{2L}{\hbar} = i(c \dot{c} - \dot{c}c) + \frac{i}{2}(\dot{a} \dot{a} - \dot{a}a) - \frac{1}{2}g^2 a^2 + g(c \dot{c} + \frac{1}{2}\dot{a}a)^2 + \frac{1}{4}(c \dot{a} + \dot{a}c)^2; \quad (9)$$

contains the recoil frequency $\omega_q = 2\hbar q^2/m$ associated with the momentum transfer $2\hbar q$, and $g = (\hbar q)^{-1}U_0$ is the effective dipole-dipole coupling constant. The Euler-Lagrange equations corresponding to Eq. (9) read

$$\dot{a} = i g a + i(g=2)(3\dot{c}^2 a + \dot{a} \dot{c}^2 + c^2 \dot{a}); \quad (10)$$

$$\dot{c} = i(g=4)(4\dot{c}^2 c + 3\dot{a} \dot{c}^2 + a^2 \dot{c}) \quad (11)$$

and thus the effective Hamiltonian density

$$\frac{2H}{\hbar} = \frac{1}{2}g^2 \dot{a}^2 + \frac{1}{4}g^2 \dot{c}^2 + 6\dot{a} \dot{c}^2 + 4\dot{c}^4 + (ac)^2 + (a \dot{c})^2 \quad (12)$$

is a constant of the motion. Another dynamical invariant of Eqs. (10) and (11) is the linear density of atoms

$$N = \dot{c}^2 + \dot{a}^2 = 2; \quad (13)$$

Hence, we are left with three independent real variables, which can be chosen as

$$n = \frac{1}{2}(\dot{a}^2 - \dot{c}^2); \quad (14)$$

$$s = \frac{1}{2} \text{Re}(a \dot{c}); \quad (15)$$

$$p = \frac{1}{2} \text{Im}(a \dot{c}); \quad (16)$$

We then arrive at the set of equations

$$\dot{n} = (g=2)sp; \quad (17)$$

$$\dot{s} = -qp; \quad (18)$$

$$\dot{p} = s - q + \frac{1}{2}gn \quad (19)$$

which is similar to those governing the evolution of the Bloch vector [24]. In particular, s is the analog of the Rabi frequency. A general solution of these equations can be found in terms of elliptic functions.

We now focus on stationary solutions, that is x points of Eqs. (17)–(19) and discuss their stability. We recognize that the choice $s = p = 0$ is the simplest stationary solution. The definitions of s and p imply that either a or c are zero. In case $a = 0$ and $c \neq 0$ we find from Eq. (8) a spatially uniform solution. From the definition of n and the dynamical invariant N we arrive at $n_u = -2N$. The case $c = 0$ and $a \neq 0$ corresponds to a purely sinusoidal $\psi(z)$ leading to $n_s = 2N$. These two x points exist for all values of the parameters.

The third x point results from the choice $p = 0$ and $s \neq 0$. This condition implies $q + gn = 2 = 0$, which leads to $n_{ss} = -2q/g$. Moreover, $p = 0$ and $s \neq 0$ demands $a \neq 0$ and $c \neq 0$ and therefore corresponds to a spatial modulation on top of a constant background. From the definitions of s and p we find

$$s_{ss} = \frac{q}{2N^2} (q/g)^2; \quad (20)$$

Since s has to be real, this x point exists provided $gN > q$, that is the product of g determined by the laser intensity and the atom density N exceeds a threshold value, determined by the recoil frequency q .

We proceed with a standard stability analysis and find the perturbation eigenfrequencies

$$\omega_u = \frac{q}{q - q + gN}; \quad (21)$$

$$\omega_s = \frac{q}{q - q + gN = 2}; \quad (22)$$

$$\omega_{ss} = \frac{q}{(gN)^2 - \frac{q^2}{g}}; \quad (23)$$

From the expressions for ω_{ss} and ω_u we recognize that the spatially modulated solution is stable throughout its range of existence, whereas the uniform solution is unstable in the same region. The uniform solution is stable if $gN < q$, that is when the modulated solution does not exist. The sinusoidal solution is always stable. However, its energy determined by the Hamiltonian (12) is larger than the one corresponding to the other stable solution. Hence, the modulated solution represents the spatially modulated ground state of the quasi-1D BEC illuminated by laser light. If the condition $gN > q$ is not met, then the ground state corresponds to the uniform solution. The latter becomes unstable with the increase of the laser intensity and/or the BEC density beyond the bifurcation point $gN = q$.

IV. NUMERICAL RESULTS

We now turn to the numerical solution of Eq. (1), using the second-order split-operator method. The integral convolution of the kernel V_{dd} with the matter density $j j^2$ in the last term of Eq. (1) is calculated using the Fast Fourier Transform (FFT) technique [25].

To circumvent the difficulties with the singularity of V_{dd} at $r = 0$ we introduce a small cut-off radius $r_c = 10^{-3} q^{-1}$. We calculate the time evolution of the condensate starting from the ground state of the BEC in an elongated ($\omega_z \ll \omega_\perp$) trap and switch on the laser intensity. In order to maximize the overlap of the initial (cigar-shaped) and final (spatially modulated) ground state, we try to keep the size of the condensate fixed when we increase the intensity of the laser field. In our numerical calculations we change the trapping potential by decreasing ω_z , so as to compensate for the laser-induced compression (electrostriction) of the condensate, see upper left plot in Fig. 1. Another way is to ramp up the scattering length by approaching a Feshbach resonance at a rate similar to the laser switch-on.




FIG. 1. Spontaneous laser-induced periodicity due to LIDDI. A BEC is illuminated by a circularly polarized laser beam oriented along the z -axis. The upper left plot shows the time dependence of the laser intensity and the longitudinal trap frequency. The time evolution of the density is shown in (a)–(e) for various times marked in the upper left plot. We have used scaled quantities where time is measured in periods of the transverse motion and lengths are measured in units of half a laser wavelength. The dimensionless dipole-dipole interaction strength is defined by $\gamma = 2^2 N U_0 / (\hbar \omega_q)$. The other parameters are $2^2 \omega_z = \omega_q = 5$ and $N q a_s = 25$. Here N is the number of particles and ω_q denotes the recoil frequency introduced in Eq. (9). Note that we have used different scalings for the radial and the longitudinal direction.

Figure 1 clearly shows spontaneous laser-induced periodicity: After we have increased the laser intensity and decreased the longitudinal frequency sufficiently enough, a spatially modulated density emerges.

Moreover, we perform also imaginary time calculations to find the quasi-stationary density distribution for two different values of the laser intensity. Our numerical results show that these density distributions are spatially modulated only if the laser intensity exceeds a certain critical threshold, see Fig. 2. For the larger laser intensity we find a quasi-stationary distribution which has a similar shape as the final state in Fig. 1e. It is noteworthy

that no oscillatory structure is numerically observed in the radial direction.

We find qualitative agreement between the variational approach and the numerical results. Our variational approach predicts spontaneous laser-induced periodicity for $gN = q > 1$, whereas from our numerical data we find a transition to a spatially modulated density at $gN = q \approx 2.5$. The agreement is not too bad considering the fact that there are only few density peaks in the numerically obtained density distribution.




FIG. 2. Stationary solutions of Eq. (1) for a scaled laser intensity of $\gamma = 234.55$ (a) and $\gamma = 268.06$ (b). The other parameters are the final values of Fig. 1. Only if the laser intensity is large enough the density is spatially modulated. Again, we have used different scalings for the radial and the longitudinal direction.

For numerical reasons we had to limit the length of the cigar-shaped BEC to few laser wavelengths. We have not found any spatially modulated density, neither in the longitudinal nor in the transverse direction, if the condensate is significantly wider than approximately half a wavelength. Our numerical results therefore do not allow us to answer the question whether we have to confine the radial direction to approximately half a laser wavelength in order to observe spontaneous laser-induced periodicity, or whether a sufficiently large aspect ratio can already lead to spontaneous laser-induced periodicity.

V. DISCUSSION

A possible experimental realization could involve BECs with a linear density of approximately 630 atoms/ μm . For ^{87}Rb the laser detuning may be chosen to be 2–6.5 GHz, the resonant wavelength being 0.795 μm . A pronounced spatially modulated density should then appear at the laser intensity $I_c = 38 \text{ mW}/\text{cm}^2$, that corresponds to $\gamma_{\text{Ray}} = 133 \text{ s}^{-1}$. Our variational approach [Eq. (20)] gives $I_c = 34 \text{ mW}/\text{cm}^2$, where $2 \log[\gamma] = (2 \gamma)$.

We may assume exponential growth of the laser intensity at the rate of 10^3 s^{-1} and simultaneous decrease of the longitudinal frequency ω_z . Under these conditions,

the collective oscillations in the BEC, which are excited due to non-adiabaticity of switching of the intensity, are found to be small. The total depletion of the BEC due to incoherent photon scattering is around 50 percent. Thus one can see that spontaneous laser-induced periodicity is possible, although accompanied by considerable losses of the number of condensed atoms.

A. Quantum depletion

We are aware of the problem of the quantum depletion of the condensate. Indeed, in a recent paper [26] the quantum depletion has been calculated in the Popov approximation for a case of a laser-radiated BEC in a geometry alternative to that of our present work. In contrast to the present geometry, where a circularly polarized plane wave is traveling along a cigar-shaped BEC, Ref. [26] considers a situation in which the wave vector and the polarization vector of a linearly polarized laser beam are perpendicular and parallel to the major axis of a cigar-shaped BEC, respectively. As the laser-induced roton-like minimum in the BEC elementary excitation spectrum tends to touch the horizontal axis, the quantum depletion grows rapidly, the peak being centered at the roton-minimum position in the momentum space. However, in the present paper we consider, unlike Ref. [26], not a stationary but a time-dependent regime. The time of formation of the distribution of the above-condensate atoms in a system where both the short-range (s-wave scattering) and long-range (LDDI) interactions are present is an important question on its own. However, we assume that the increase of the laser intensity in our case is rapid enough to prevent the total depletion of the condensate before the spontaneous laser-induced periodicity, that is, a transition from a uniform BEC to a spatially modulated BEC, can be observed.

Nevertheless, a certain depletion is unavoidable. Thus, as soon as we exceed the intensity threshold, the above-condensate atoms act as a seed perturbation, which will grow exponentially at the wave vectors where the spatially uniform solution becomes unstable. Therefore, the instability growth is concurrent to the emergence of the coherent spatial structure. Although the magnitude of the seed perturbation has to be calculated beyond the mean-field (Gross-Pitaevskii equation) approach, the instability increment can be derived from the linearized Gross-Pitaevskii equation [18]. In numerical simulations of unstable BECs the role of seed perturbation is played by numerical noise [27]. Stable numerical algorithms avoid amplification of numerical rounding errors provided that the numerical solution converges to the exact one, which is stable. However, if one reaches an instability range, the numerical errors begin to grow due to the properties of the equation being solved, and not due to an unstable numerical algorithm. Since our numerical solutions (see Fig. 1) display a stable spatial structure, we

may conclude that our transition from one stable branch (the uniform one) to another (the spatially modulated one) is fast enough to prevent an appreciable condensate depletion and the development of a dynamical instability.

B. Alternative approach

A better strategy to obtain similar spatially modulated densities is to first prepare the density-modulated ground state in an optical lattice formed by a far-detuned standing wave. We then switch off the optical lattice and simultaneously turn on a running wave which creates the LDDI. It is now this running wave that keeps the BEC in the initially prepared density-modulated ground state. Although it is technically simpler to create a spatially modulated self-sustained structure in a BEC starting from a BEC confined in an external optical lattice, the way studied in the present paper has its advantage. It clearly shows the spontaneous emergence of spatially modulated structures when the laser intensity is gradually increased above a threshold value.

C. Laser-induced lattice formation versus coherent atom recoil lasing

We emphasize the key difference between the formation of spatially modulated densities due to LDDI in a laser-illuminated BEC and superradiant light scattering by a BEC: The latter, related to coherent atom recoil lasing (CARL) [16], is essentially non-stationary, resulting in the generation of phonons in a BEC. In contrast, the density modulations discussed in the present paper are quasi-stationary. Accordingly, under the conditions of slow switch-on of the laser, we may expect a spatially-oscillatory ground state without triggering superradiance. On the other hand, CARL has no threshold in terms of the laser-beam intensity, unlike the effects discussed above. Hence, a weak, abruptly switched-on laser pulse would yield only superradiance.

VI. CONCLUSIONS

To conclude, our 3D numerical calculations have substantiated the simple 1D variational ansatz, predicting the gradual formation of an axial oscillatory ground-state density modulation in a laser-illuminated cigar-shaped BEC. Its origin can be unambiguously traced to LDDI. Although the depletion of the condensate due to incoherent photon scattering and due to instabilities at the transition from a uniform condensate to a spatially modulated condensate is considerable, we believe that the creation of spatial modulations in a BEC is, as discussed in the present paper, experimentally feasible. We emphasize the similarities between the spatial modulations due

to LDDI and the modulations of a BEC in an optical lattice. We may call the formation of a spatially modulated density a spontaneous formation of a self-sustained optical lattice which is formed by interfering incident and coherently backward scattered light waves.

ACKNOWLEDGMENTS

The support of the German-Israeli Foundation, the EC (the QUACSRN) and Minerva is acknowledged. IEM also thanks the Russian foundations (RFBR 02{02{17686, UR.01.01.040, E02{32{287).

-
- [1] A. Stenger, S. Inouye, D.M. Stamper-Kum, H.-J. Miesner, A.P. Chikkatur, and W. Ketterle, *Nature* 396, 345 (1999).
 - [2] M.D. Barrett, J.A. Sauer, and M.S. Chapman, *Phys. Rev. Lett.* 87, 010404 (2001).
 - [3] See, e.g., papers cited by B.Goss Levi, *Phys. Today* 54 (8), 15 (2001); 55 (3), 18 (2002).
 - [4] D.O'Dell, S.Giovanazzi, G.Kurizki, and V.M.Akulin, *Phys. Rev. Lett.* 84, 5687 (2000); S.Giovanazzi, D.O'Dell, and G.Kurizki, *Phys. Rev. A* 63, 031603(R) (2001); S.Giovanazzi, G.Kurizki, I.E.Mazets, and S.Stringari, *Europhys. Lett.* 56, 1 (2001); G.Kurizki et al., *Springer Lectures in Physics* 602, 382 (Springer, 2002).
 - [5] S.Giovanazzi, D.O'Dell, and G.Kurizki, *Phys. Rev. Lett.* 88, 130402 (2002).
 - [6] O.Penrose and L.Onsager, *Phys. Rev.* 104, 576 (1956).
 - [7] R.T.Whitlock and P.R.Zilsel, *Phys. Rev.* 131, 2409 (1963).
 - [8] Y.-Ch.Cheng, *Phys. Rev. B* 14, 1946 (1976).
 - [9] G.V.Chester, *Phys. Rev. A* 2, 256 (1970).
 - [10] A.J.Leggett, *Phys. Rev. Lett.* 25, 1543 (1970).
 - [11] L.P.Pitaevskii, *JETP Lett.* 39, 511 (1984).
 - [12] Y.Pomeau and S.Rica, *Phys. Rev. Lett.* 72, 2426 (1994).
 - [13] K.Goral, L.Santos, and M.Lewenstein, *Phys. Rev. Lett.* 88, 170406 (2002).
 - [14] E.Kim and M.H.W.Chan, *Nature* 427, 225 (2004).
 - [15] B.P.Anderson and M.A.Kasevich, *Science* 282, 1686 (1999).
 - [16] S. Inouye, A.P. Chikkatur, D.M. Stamper-Kum, J.Stenger, D.E.Pritchard, and W. Ketterle, *Science* 285 571 (1999); D.Schneble, Y.Torii, M.Boyd, E.W.Streed, D.E.Pritchard, and W. Ketterle, *Science* 300, 475 (2003); M.G.Moore and P.Meystre, *Phys. Rev. Lett.* 83, 5202 (1999); N.Piovella, R.Bonifacio, B.W.J.McNeil, and G.R.M.Robb, *Opt. Commun.* 187, 165 (2001).
 - [17] L.P. Pitaevskii, *Sov. Phys. JETP* 13, 451 (1961); E.P.Gross, *Nuovo Cimento* 20, 454 (1961).
 - [18] F.Dalfó, S.Giorgini, L.P. Pitaevskii, and S.Stringari, *Rev. Mod. Phys.* 71, 463 (1999).
 - [19] S.T.Belyaev, *Zh.Eksp.Teor.Fiz.* 34, 417, (1958); *ibid.*, 433.
 - [20] The potential in this limit accounts for the "artificial gravity" that may give rise to spherically-symmetric self-bound BEC states [4].
 - [21] S.Yi and L.You, *Phys. Rev. A* 61, 041604(R) (2000).
 - [22] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics*, (Oxford, Butterworth-Heinemann, 1997).
 - [23] L.Salasnich, A.Parola and L.Reatto, *Phys. Rev. A* 65, 043614 (2002).
 - [24] S.Stenholm, *Foundations of Laser Spectroscopy* (Wiley, NY, 1984).
 - [25] K.Goral and L.Santos, *Phys. Rev. A* 66, 023613 (2002).
 - [26] I.E.Mazets, D.H.J.O'Dell, G.Kurizki, N.Davidson, and W.P.Schleich, "Depletion of a Bose-condensate by laser-induced dipole-dipole interactions". *ArXiv: cond-mat/0310672* (*J.Phys.B*, in press).
 - [27] P.A.Ruprecht, M.J.Holland, K.Burnett, and M.Edwards, *Phys. Rev. A* 51, 4704 (1995).