

# FERROELECTRIC PHASE TRANSITIONS IN A LATTICE PSEUDO-JAHN-TELLER MODEL

Dennis P. Clougherty

Department of Physics

University of Vermont

Burlington, VT 05403-0125

dpc@physics.uvm.edu

**Keywords:** ferroelectric, antiferroelectric, pseudo-Jahn-Teller

**Abstract** A lattice generalization of the  $(b_{1u} \ b_{2g}) \ b_{3u}$  pseudo-Jahn-Teller model is analyzed within mean-field theory. Ferroelectric or antiferroelectric ground states are found for sufficiently large pseudo-Jahn-Teller coupling.

## Introduction

Inspired by the family of fullerene-based polymeric structures synthesized [1, 2], a model describing vibronic effects in a linear chain of molecules is introduced and analyzed within mean-field theory. The zero-temperature phase diagrams at carrier concentrations  $n = 1$  and  $2$  are presented for this model. Only a local  $(b_{1u} \ b_{2g}) \ b_{3u}$  pseudo-Jahn-Teller interaction is considered here; spin and charge fluctuation effects are ignored.

The local symmetry of the molecule is taken to be  $D_{2h}$  for simplicity; the set of  $b$ -representations transform as components of a vector. The electronic structure of the molecule is taken to have a  $b_{1u}$  level as the HOMO and a  $b_{2g}$  level as the LUMO. A distortion of  $b_{3u}$  symmetry will vibronically couple these levels. For a molecule with some ionic bonding character, such a  $b_{3u}$  distortion produces an electric dipole moment in proportion to the displacement [3]. It is the ordering of these local dipole moments that is the concern of this work.

## 1. THE HAMILTONIAN

The Hamiltonian is given by

$$H = H_0 + H_{\text{PJT}} + H_{\text{elast}} \quad (1.1)$$

where

$$H_0 = \sum_n (c_{1n}^\dagger c_{1n+1} + c_{2n}^\dagger c_{2n+1} + \epsilon) + \sum_n c_{2n}^\dagger c_{2n} \quad (1.2)$$

where  $n$  runs over molecule sites,  $c_{1n}^\dagger$  ( $c_{1n}$ ) creates (annihilates) an electron in orbital  $b_{1u}$  at site  $n$ ,  $c_{2n}^\dagger$  ( $c_{2n}$ ) creates (annihilates) an electron in orbital  $b_{2g}$  at site  $n$ , and  $\epsilon$  is the energy spacing between the  $b_{1u}$  and  $b_{2g}$  orbitals. The hopping matrix is taken to be diagonal in orbital type and spin, and the spin indices are suppressed for clarity.

The pseudo-Jahn-Teller Hamiltonian on the  $n$ th molecular site is

$$H_{\text{PJT}} = g \sum_n (c_{1n}^\dagger c_{2n} Q_n + \text{h.c.}) \quad (1.3)$$

where  $g$  is the pseudo-Jahn-Teller coupling strength, and  $Q_n$  is a displacement of  $b_{3u}$  symmetry on the  $n$ th molecule. The magnitude of the induced local electric dipole is proportional to  $Q_n$ . Hence, a ferrodistortive state is equated with a ferroelectric one.

$H_{\text{elast}}$  is the elastic energy for the molecular displacements.

$$H_{\text{elast}} = \frac{1}{2} \sum_n Q_n^2 \quad (1.4)$$

The Hamiltonian can be written in  $k$ -space as

$$H = \sum_{jk} (c_{jk}^\dagger c_{jk} + \frac{g}{N} \sum_{k,q} (c_{1k+q}^\dagger c_{2k} Q_q + \text{h.c.})) + \frac{1}{2} \sum_q Q_q^2 \quad (1.5)$$

where  $g(k) = 2t \cos k$  and  $u(k) = 2t \cos k$ .

## 2. MEAN FIELD THEORY

With a single mode of distortion given by a non-vanishing  $Q_q$ ,  $H$  can be diagonalized with a canonical transformation so that

$$H = \sum_k (E_{k+} n_{k+} + E_{k-} n_{k-}) + \frac{1}{2} Q_q^2 \quad (1.6)$$

where

$$E_k = \frac{1}{2} (u(k) + g(k+q)) \pm \frac{1}{2} \sqrt{(g(k+q) - u(k))^2 + \frac{4g^2}{N} Q_q^2} \quad (1.7)$$

and  $n_k$  are the usual occupation number operators of the vibronic quasiparticles.

The ground state energy of a chain with  $n$  carriers ( $0 \leq n \leq 2$ ) per molecule on average is obtained by fractionally filling the lowest energy vibronic band. In the continuum limit,

$$E(Q_q) = \frac{N}{2\pi} \int_{-\pi}^{\pi} dk E_k + \frac{1}{2} Q_q^2 \quad (1.8)$$

Minimizing  $E(Q_q)$  with respect to the distortion gives the mean-field equation to determine the magnitude of  $Q_q$ .

$$\frac{1}{2g^2} = \frac{1}{2\pi} \int_{-\pi}^{\pi} dk \frac{1}{\frac{4g^2}{N} Q_q^2 + (\epsilon + 4t \sin q/2 \sin(k + q/2))^2} \quad (1.9)$$

For sufficiently small coupling, this equation can not be satisfied. There is a  $q$ -dependent critical coupling strength  $g_c$  given by

$$g_c^2(q) = \frac{1}{2\pi} \int_{-\pi}^{\pi} dk \frac{1}{1 + u \sin q/2 \sin(k + q/2)} \quad (1.10)$$

where  $u = \frac{4t}{N}$ . For values of  $g$  above  $g_c(q)$ , a mean-field distortion of wavenumber  $q$  sets in; for values of  $g$  below  $g_c(q)$ , there is no such distortion.

### 3. THE PHASE DIAGRAM

For the case of an average occupancy of one carrier per molecule, the ground state is found to have a  $q = 0$  distortion for  $g > g_0 (= \frac{1}{2})$ , consequently breaking inversion symmetry of the chain. The phase diagram is pictured in Fig. 1.

For the case of two carriers per molecule, the ground state has a  $q = \pi$  distortion when

$$g^2 > \frac{g_0^2}{2} \frac{1}{1 - u^2} \quad (1.11)$$

The results are summarized in Fig. 2.

### Acknowledgments

This work was supported by the National Science Foundation through a grant for the Institute for Theoretical Atomic and Molecular Physics at Harvard University and the Smithsonian Astrophysical Observatory.




Figure 1 Zero-temperature phase diagram for  $n = 1$ .




Figure 2 Zero-temperature phase diagram for  $n = 2$ .

## References

- [1] Pekker, S. et al., Science 256, 1077 (1994).
- [2] Stephens P.W. et al., Nature 370, 636 (1994).
- [3] Clougherty D. P. and Anderson F. G., Phys. Rev. Lett. 80, 3735 (1998).