

Quantum-classical transition of the escape rate of uniaxial antiferromagnetic particles in an arbitrarily directed field

Bin Zhou^{1,2,3}, Rubao Tao¹, and Shun-Qing Shen²

¹Department of Physics, Fudan University, Shanghai 200433, China

²Department of Physics, The University of Hong Kong, Hong Kong, China

³Department of Physics, Hubei University, Wuhan 430062, China

(Dated: April 14, 2024)

Quantum-classical escape rate transition has been studied for uniaxial antiferromagnetic particles with an arbitrarily directed magnetic field. In the case that the transverse and longitudinal fields coexist, we calculate the phase boundary line between first- and second-order transitions, from which phase diagrams can be obtained. It is shown that the effects of the applied longitudinal magnetic field on quantum-classical transition vary greatly for different relative magnitudes of the non-compensation.

PACS numbers: 75.50.Xx, 75.45.+j, 03.65.Sq

Escape from a stable or metastable state at high temperatures is governed by a classical thermal activation rate. At low temperatures close to zero, quantum tunneling becomes relevant. When these two escape rates are equal there exists a crossover temperature T_0 at which a transition between classical and quantum regimes occurs. The study of the quantum-classical transition is an interesting subject with a long history.[1] One of the main issues in this subject is to determine whether the transition is first- or second-order. The transition was recognized as a smooth second-order one in the quantum mechanical models of Abeck [1] and the cosmological models of Linde.[2] However, it was shown [3] that the smooth transition is not generic. Chudnovsky has suggested that the order of transition is determined by the behavior of the Euclidean time oscillation period $\langle E \rangle$, where E is the energy near the bottom of the Euclidean potential, which corresponds to the top of the potential barrier.[3] The non-monotonic behavior of the oscillation period as a function of energy, i.e., the existence of a minimum in the $\langle E \rangle$ curve, was proposed as a condition for the first order transition in quantum-mechanical tunneling.[3] Later, a sufficient criterion for the first-order phase transition was obtained by carrying out the nonlinear perturbation near the sphaleron solution.[4]

Since the first- and second-order transitions between the quantum and classical behaviors of the escape rates in spin systems were introduced by Chudnovsky and Garanin,[5, 6] the topic has attracted considerable attention.[7, 8, 9, 10, 11, 12, 13, 14, 15] Most theoretical studies have been focused on the ferromagnetic particles. However, most ferromagnetic systems are actually ferrimagnetic particles. For instance, both $Mn_{12}Ac$ and Fe_8 are characterized by a large spin ground state which originates from incomplete compensation of antiferromagnetically coupled spins.[16] The strong exchange interaction should be taken into consideration. In Ref.[17] Kim treated the phase transition in ferrimagnetic or antiferromagnetic particles for two general forms of the magnetic anisotropy energy. Very recently, the quantum-

classical transition in antiferromagnetic particles with biaxial symmetry in the presence of an applied magnetic field along the medium axis or along the easy axis was investigated.[18, 19] Note that recent work of Chudnovsky and Garanin postulates dislocations as the main source of spin tunneling in Mn_{12} crystals.[20] Their theory shows that when the external magnetic field is applied along the c axis of the crystal, local rotations of the magnetic anisotropy axis due to dislocations result in the effective local transverse magnetic field. Experimental evidence of the effects of dislocations on tunneling has been also reported.[21, 22, 23, 24] Therefore in the study of the quantum-classical transition of Mn_{12} , the case of coexistence of the transverse and longitudinal magnetic field is worth investigating. Considering the molecular cluster Mn_{12} actually is ferrimagnetic, the exchange interaction should be also taken into account. In this paper we aim to investigate the quantum-classical transition of the escape rate of uniaxial antiferromagnetic particles in an arbitrarily directed field, i.e., the coexistence of the transverse and longitudinal magnetic field. It is shown that the effects of the applied longitudinal magnetic field on the quantum-classical transition vary greatly for different relative magnitudes of the non-compensation.

We consider a small uniaxial antiferromagnetic particle with two magnetic sublattices whose magnetizations, m_1 and m_2 , are coupled by the strong exchange interaction $m_1 m_2 = \chi$, where χ is the perpendicular susceptibility. The system of interest has a non-compensation of sublattice with m ($= m_1 - m_2 > 0$), and easy-axis anisotropy along the z-axis. In the presence of an arbitrarily directed magnetic field, i.e., the coexistence of a transverse magnetic field H_x along the x-axis and a longitudinal one H_z along the z-axis, the Euclidean action is written as[25]

$$S_E(\theta; \phi) = V \int_0^{\beta} d\tau \left[\frac{m_1 + m_2}{2} \dot{\theta}^2 - \frac{m}{2} \dot{\phi}^2 - \cos \theta \right. \\ \left. + \frac{e_2}{2} [(-\cos \theta + i H_x \sin \theta)^2 + (-\sin \theta + i H_x \cos \theta \cos \phi - i H_z \sin \theta)^2] \right]$$

$$+ K_k \sin^2 \theta - m H_x \sin \theta \cos \theta \\ m H_z \cos \theta - g; \quad (1)$$

where V is the volume of the particle, γ the gyromagnetic ratio, $e_\gamma = \gamma / (m_2 - m_1)$ and K_k the longitudinal anisotropy. The polar coordinate θ and the azimuthal coordinate ϕ , which are the angular components of m_1 in the spherical coordinate system, can determine the direction of the Neel vector. Dot over a symbol denotes a derivative with respect to the Euclidean time t .

The classical trajectory corresponding to the Euclidean action (1) is determined by the equations

$$\frac{d}{dt} \left(\sin^2 \theta \right) + x \left(\frac{d}{dt} \sin^2 \theta - 2 \frac{d}{dt} \cos \theta \sin^2 \theta \right) \\ + \frac{1}{2} \frac{d^2}{dt^2} \sin^2 \theta = \frac{V}{K_k} \\ = 0; \quad (2)$$

$$\frac{d}{dt} \left(\sin^2 \theta \right) + x \left(\frac{d}{dt} \sin^2 \theta + 2 \frac{d}{dt} \cos \theta \sin^2 \theta \right) \\ - \sin^2 \theta = \frac{V}{K_k} \\ = 0; \quad (3)$$

where $n = m = (K_k)^{-1}$, $x = e_\gamma = (K_k)^{-1}$, $b_{x(z)} = H_{x(z)}$, $V = \partial V / \partial \theta$ and $\dot{V} = \partial V / \partial \dot{\theta}$. The inverted potential is

$$V(\theta; \phi) = K_k \left(\sin^2 \theta + 2 h_x \sin \theta \cos \theta + 2 h_z \cos \theta \right. \\ \left. + \frac{x b_x^2}{2} \sin^2 \theta + \frac{x b_x^2}{2} \cos^2 \theta - \cos^2 \theta \right. \\ \left. + \frac{x b_z^2}{2} \sin^2 \theta - x b_x b_z \sin \theta \cos \theta \cos \phi \right); \quad (4)$$

where $h_{x(z)} = H_{x(z)} / H_c$ and $H_c = 2 K_k m$. In Fig.1 the effective potential $V(\theta; \phi = 0)$ is drawn. The minima of the potential correspond to the equilibrium orientation of the Neel vector. The metastability condition that $dV(\theta; 0) / d\theta = 0$ and $d^2 V(\theta; 0) / d\theta^2 = 0$ determines the critical parameters at which the barrier vanishes.

In the high-temperature regime the sphaleron solution of Eqs.(2) and (3) is $(\theta_0; \phi_0 = 0)$. θ_0 is the position of the top of the potential barrier $V(\theta_0; 0)$, and is determined by $dV(\theta; 0) / d\theta = 0$ and $d^2 V(\theta; 0) / d\theta^2 > 0$.

θ_0 has a cumbersome analytical form and its numerical result will be applied below to determine phase boundary lines. Furthermore, its behavior is illustrated in Fig.2 for given h_x and y . Above, $y = x n^2 (= \gamma^2 K_k m^2)$ and the parameter y indicates the relative magnitude of the non-compensation. For large non-compensation ($y \gg 1$; i.e., $m \gg \gamma^2 K_k$) and for small non-compensation ($y \ll 1$; i.e., $m \ll \gamma^2 K_k$), the system becomes ferromagnetic and nearly compensated antiferromagnetic, respectively.[17] Note that $\theta_0 = \pi/2$ for $h_z = 0$. The crossover behavior of the escape rate of this model from quantum tunneling to thermal activation can be obtained from the deviation of the period of the periodic instanton from that of the sphaleron. To this end we expand

FIG.1: The effective potential $V(\theta; \phi = 0)$:

FIG.2: θ_0 versus h_z for some given h_x and y :

$V(\theta; \phi)$ about the sphaleron configurations θ_0 and ϕ_0 , i.e., $\theta = \theta_0 + \delta\theta$ and $\phi = \phi_0 + \delta\phi$, where $\phi_0 = 0$. Substituting them into Eqs.(2) and (3) one yields the following power series equations of the fluctuations $\delta\theta$ and

$$G_1(\delta\theta; \delta\phi) + G_2(\delta\theta; \delta\phi) + G_3(\delta\theta; \delta\phi) + \dots = 0; \quad (5)$$

where G_1, G_2, G_3, \dots denote terms which contain linear, quadratic, cubic and higher powers of the small fluctuations, respectively:

$$G_1(\delta\theta; \delta\phi) = \frac{1}{2} \left(\frac{d^2 V}{d\theta^2} \delta\theta^2 + 2 \frac{d^2 V}{d\theta d\phi} \delta\theta \delta\phi + \frac{d^2 V}{d\phi^2} \delta\phi^2 \right);$$

$$G_2(\delta\theta; \delta\phi) = \frac{1}{6} \left(\frac{d^3 V}{d\theta^3} \delta\theta^3 + 3 \frac{d^3 V}{d\theta^2 d\phi} \delta\theta^2 \delta\phi + 3 \frac{d^3 V}{d\theta d\phi^2} \delta\theta \delta\phi^2 + \frac{d^3 V}{d\phi^3} \delta\phi^3 \right);$$

$$G_3(\delta\theta; \delta\phi) = \frac{1}{24} \left(\frac{d^4 V}{d\theta^4} \delta\theta^4 + 6 \frac{d^4 V}{d\theta^3 d\phi} \delta\theta^3 \delta\phi + 12 \frac{d^4 V}{d\theta^2 d\phi^2} \delta\theta^2 \delta\phi^2 + 6 \frac{d^4 V}{d\theta d\phi^3} \delta\theta \delta\phi^3 + \frac{d^4 V}{d\phi^4} \delta\phi^4 \right);$$

$$+ 2ix(b_z \sin 2\phi_0 - b_x \cos 2\phi_0)^2 - \\ + ix b_x \sin^2 \phi_0^2 -;$$

$$G_1(\phi; \omega) = i \sin \phi_0 - ix(b_z \sin 2\phi_0 + 2b_x \sin^2 \phi_0) - \\ + x \sin^2 \phi_0 + B_1 \phi;$$

$$G_2(\phi; \omega) = i \cos \phi_0 - 2ix(b_z \sin 2\phi_0 + b_x \cos 2\phi_0) - \\ + x \sin 2\phi_0 (-\phi + \phi_0) + B_2 \phi;$$

$$G_3(\phi; \omega) = \frac{1}{2} i \sin \phi_0^2 - ix b_x \sin^2 \phi_0^2 - \\ + 2ix(b_z \sin 2\phi_0 - b_x \cos 2\phi_0)^2 - \\ + x \cos 2\phi_0 (2\phi - \phi_0^2) + B_3 \phi^2 + B_4 \phi^3;$$

where

$$A_1 = \frac{V}{K_k}; A_2 = \frac{V}{2K_k}; A_3 = \frac{V}{2K_k}; \\ A_4 = \frac{V}{2K_k}; A_5 = \frac{V}{6K_k}; \quad (6)$$

$$B_1 = \frac{V}{K_k}; B_2 = \frac{V}{K_k}; B_3 = \frac{V}{2K_k}; B_4 = \frac{V}{6K_k}; \quad (7)$$

It is introduced that $V = [\partial^2 V = \partial^2] = \phi_0; = \phi_0, V = [\partial^2 V = \partial^2] = \phi_0; = \phi_0$, and so on.

Denoting $\phi(\omega) = \phi(\omega; \phi_0)$, we have $\phi(\omega + \sim) = \phi(\omega)$ at finite temperature and write it as the Fourier series $\phi(\omega) = \sum_{n=1}^{\infty} \phi_n \exp[i!_n]$, where $!_n = 2\pi n = \sim$. Since simple analysis shows that ϕ is real and imaginary, to the lowest order we write them in the form $\phi = a_1 \cos(!)$ and $\phi = ia_1 \sin(!)$. Here a serves as a perturbation parameter. Substituting them into Eq.(5) and neglecting terms of order higher than a , we obtain the relation

$$\frac{1}{\omega} = \frac{x!^2 + A_1}{! [\sin 2x(b_z \cos \phi_0 + b_x \sin \phi_0)] \sin \phi_0} \\ = \frac{! [\sin 2x(b_z \cos \phi_0 + b_x \sin \phi_0)] \sin \phi_0}{x!^2 \sin^2 \phi_0 - B_1} \quad (8)$$

and the oscillation frequency

$$!^2 = \frac{1}{2x^2} f(A_1 - B_1 \csc^2 \phi_0) x \\ + [\sin 2x(b_z \cos \phi_0 + b_x \sin \phi_0)]^2 g \\ \frac{1}{2x^2} (4A_1 B_1 x^2 \csc^2 \phi_0 + f(A_1 - B_1 \csc^2 \phi_0) x \\ + [\sin 2x(b_z \cos \phi_0 + b_x \sin \phi_0)]^2 g^2)^{1/2}; \quad (9)$$

Next, let us write $\phi = a_1 \cos(!) + \phi_2$, and $\phi = ia_1 \sin(!) + i_2$, where ϕ_2 and i_2 are of the order of a^2 . Inserting them into Eq.(5), we arrive at $! = !_+$ and

$$\phi_2 = a^2 p_0 + a^2 p_2 \cos(2!); \quad i_2 = a^2 q_2 \sin(2!); \quad (10)$$

where the analytic forms of coefficients p_0, p_2 and q_2 are cumbersome, which are listed in the Appendix.

This implies that there is no shift in the oscillation frequency. In order to find the change of the oscillation period, we proceed to the third order of perturbation theory by writing $\phi = a_1 \cos(!) + \phi_2 + \phi_3$, and $\phi = ia_1 \sin(!) + i_2 + i_3$, where ϕ_3 and i_3 are of the order of a^3 . Substituting them again into Eq.(5), and retaining only the terms up to $O(a^3)$, we have

$$n^4 y^2 (!^2 - !_+^2) (!^2 - !_+^2) = a^2 \frac{1}{4 \sin^2 \phi_0} g(h_x; h_z; y); \quad (11)$$

where

$$g(h_x; h_z; y) = g_1(h_x; h_z; y) + g_2(h_x; h_z; y); \quad (12)$$

The forms of $g_1(h_x; h_z; y)$ and $g_2(h_x; h_z; y)$ are

$$g_1(h_x; h_z; y) = (A_1 + yw^2) f_3 B_4^2 - 6 h_x w \sin^2 \phi_0 \\ B_3 - 2B_2(2p_0 - p_2) + 3yw^2 \cos 2\phi_0 \\ + 2yw^2 \sin 2\phi_0 (2p_0 + p_2) + \frac{1}{2} [-2B_2 q_2 \\ + 2w \cos \phi_0 (2p_0 + p_2) - 4h_x yw \cos 2\phi_0 \\ - \frac{1}{2} w \sin \phi_0 - 8h_z yw (2p_0 + p_2) \cos 2\phi_0 \\ + 4yw \sin 2\phi_0 (h_z - 4h_x p_0 - 2h_x p_2) \\ + 4q_2 yw^2 \sin 2\phi_0] g; \quad (13)$$

$$g_2(h_x; h_z; y) = (B_1 - yw^2 \sin^2 \phi_0) f_2^3 h_x yw \sin^2 \phi_0 \\ - 2(A_3 + 3yw^2 \cos 2\phi_0) \\ + [4A_2 q_2 - 2(2p_0 + p_2) w \cos \phi_0 \\ + \frac{3}{2} w \sin \phi_0 - 4q_1 yw^2 \sin 2\phi_0 \\ + 4yw \cos 2\phi_0 (3h_x + 4h_z p_0 + 2h_z p_2) \\ - 4yw \sin 2\phi_0 (3h_z - 4h_x p_0 \\ - 2h_x p_2)] + 4A_4 (2p_0 + p_2) \\ + 4q_2 w (4h_z y \cos 2\phi_0 + 4h_x y \sin 2\phi_0 \\ \cos \phi_0) + 3A_5 g; \quad (14)$$

where $w = n!_+$ and $\phi_2 = \phi_1 = 1$. Again, $y = x/n^2 (= \sim K_k m^2)$ and the parameter y indicates the relative magnitude of the non-compensation. Also, p_0, p_2 and q_2 are obtained by replacing ϕ_1 by ϕ_1 and dropping ϕ_1^2 in p_0, p_2 and q_2 , respectively. It can be shown that for $h_z = 0$ Eq.(12) is reduced to the case corresponding to uniaxial antiferromagnetic particle with a transverse magnetic field only has been investigated in Ref.[17].

As shown by Chudnovsky[3], if the oscillation period is not a monotonic function of a , where a is a function of E in the absence of dissipation, the system exhibits a first-order transition. Thus, the period $(= 2\pi/!)$ in Eq.(11) should be less than $\phi_+ (= 2\pi/!_+)$, i.e., $! > !_+$ for the first-order transition. It implies that $g(h_x; h_z; y) > 0$ in Eq.(11) for the first-order transition, and $g(h_x; h_z; y) = 0$ determines the phase boundary between the first- and the second-order transition. In

FIG. 3: Phase diagram $h_x(y)$ for $h_z = 0.1$ and 0.2 .FIG. 5: Phase diagram $h_x(h_z)$ for $y = 0.2$ and 0.3 .FIG. 4: Phase diagram $h_x(h_z)$ for $y = 0$ and 0.1 .

this case the three parameters h_x, h_z, y should be treated simultaneously, which is not a simple problem. In the present work we will fix one parameter and then compute the boundary curve with the other two parameters. We first solve Eq.(11) numerically to obtain the phase boundary lines $h_x(y)$'s for several values of h_z , which are plotted in Fig.3. From Fig.3, an immediate observation is that the first-order region for a given h_z diminishes as y increases, which shows the same trend as the $h_z = 0$ case. Thus, it is evident that the region for the first-order transition is greatly reduced as the system becomes ferrimagnetic and there is no first-order transition in almost compensated antiferromagnetic particles. The result coincides with Ref.[17]. Fig.3 also shows that with increasing h_z the variety of first-order region is not a simple case. For the small y case with increasing h_z the first-order region is shrunk, while for the larger y case the longitudinal field h_z favors occurrence of the first-order transition. For instance, for the case of $y = 0.05$, the maximum values of the transverse field h_x for occurrence of the first-order transition are $h_x = 0.203; 0.195$ and 0.175 for $h_z = 0.1$ and 0.2 , respectively. On the other

hand, the first-order region vanishes beyond $y = 0.46; 0.55$ and 0.89 for $h_z = 0.1$ and 0.2 , respectively. This can be qualitatively understood from the consideration that the height of the effective potential barrier decreases as h_z increases, whereas the height increases as y increases, therefore there is a competition between the longitudinal field and the relative magnitude of the non-compensation. When $y = 0$, the fact that the region for the first-order transition decreases as the longitudinal field increases results from a flattening of the peak of the barrier.[11] For the small y case (i.e., the large noncompensation), the crossover behavior of the ferrimagnetic system still keeps qualitatively that of the ferromagnetic one. However, for the larger y case the exchange interaction plays the role of effective magnetic field and so, for a given small transverse field the region for the first-order transition increases as the longitudinal field increases. To illustrate further the effect of the longitudinal field h_z on quantum-classical transition, we next calculate the phase boundary lines $h_x(h_z)$'s for several values of y , which are shown in Figs.4 and 5. In Fig.4, for the case of $y = 0$ corresponding to uniaxial ferromagnetic system, the phase boundary line is plotted by a dotted line. Obviously the line coincides with Fig.13 in Ref.[11], in which quantum-classical transition in a uniaxial ferromagnetic system with a transverse magnetic field and a longitudinal one was investigated. For the case of $y = 0.1$, the phase boundary lines $h_x(h_z)$'s shift downwards and with h_z increasing the critical value of h_x decrease monotonically. Fig.5 gives another case, in which the phase boundary lines $h_x(h_z)$'s show a kind of non-monotonic behavior. For instance, for the case of $y = 0.3$, the first-order region vanishes beyond $h_x = 0.024$ for $h_z = 0$, while the maximum is $h_x = 0.112$ for $h_z = 0.332$.

It was shown that quantum tunneling shall show up at higher temperatures and higher frequencies in antiferromagnetic particles than in ferromagnetic particles of similar size.[25] Moreover, most ferromagnetic systems are ferrimagnetic, so nanometer-scale antiferromagnets are more interesting from experimental and theoretical

aspects. But a detailed comparison between the theory and experiment on quantum-classical transition remains a challenging task. It is very important to obtain the information on the magnitude of the quantity y for observing the first-order transition in real experiments. For the typical antiferromagnetic particle with 10^4 , $K_k = 10^6$ erg/cm³, and $m = 500$ emu/cm³, one can get the quantity $y \sim 10^{-4}$. [19] In this case, for the longitudinal field parameter $h_z = 0.4$, the range of the transverse field parameter for observing the first-order transition is $0 < h_x \leq 0.127$. It is noted that Wensdorfer et al have performed the switching field measurements on individual ferrimagnetic and insulating BaFeCoTiO nano-particle containing about 10^5 { 10^6 spins at very low temperature (0.1{6 K). [26] Below 0.4 K, experimental results are quantitatively in agreement with the predictions of the macroscopic quantum tunneling theory without dissipation. The BaFeCoTiO nano-particles have a strong uniaxial magnetocrystalline anisotropy. [26] Therefore, the material is expected as a candidate to investigate quantum-classical transition of the escape rate of uniaxial ferrimagnetic or antiferromagnetic particles in an arbitrarily directed field.

In conclusion, we have investigated quantum-classical escape rate transition for uniaxial antiferromagnetic particle with an arbitrarily directed magnetic field, i.e., the coexistence of the transverse and longitudinal magnetic fields. There are three parameters which can be controlled by experiment: relative magnitude of the non-compensation and two field parameters. The nonlinear perturbation method is used to obtain various phase diagrams for first- and second-order transition depending on the three parameters. It is shown that the effects of the applied longitudinal magnetic field on quantum-classical transition vary greatly for different relative magnitudes of the non-compensation.

The work was supported by China Postdoctoral Science Foundation under Grant No. 2002032138, National Natural Science Foundation of China under Grant Nos. 10174015 and 10234010, and Research Grant Council of Hong Kong.

Appendix

The coefficients in Eq.(10) are deduced by using software Mathematica 3.0.

$$p_0 = \frac{1}{2A_1} [n_1 \cos \theta_0 (n_1! + x_1!_+^2 \sin^2 \theta_0)$$

$$+ A_2 \frac{x_1^2}{1} - A_4 \frac{x_1^2}{1} - 2x_1!_+ \cos 2\theta_0 + b_x \sin 2\theta_0];$$

$$\begin{aligned} p_2 = & fB_1(A_4 \frac{x_1^2}{1} + A_2 \frac{x_1^2}{1}) + 2B_2 n_1!_+ \sin \theta_0 \\ & 8b_x^2 x_1^2!_+^2 \cos^3 \theta_0 \sin \theta_0 \\ & 2x_1! (2B_2 b_x n_1!_+ + B_1 b_x n_1!_+ + 2A_4 \frac{x_1^2}{1}! + 2A_2 \frac{x_1^2}{1}!) \sin^2 \theta_0 \\ & + 8b_x x_1^2!_+^2 (b_x n_1!_+ + n_1!) \sin^4 \theta_0 \\ & 4b_x n x_1^2!_+^2 \sin^3 \theta_0 + 2b_x x_1! \cos^2 \theta_0 B_1 n_1!_+ \\ & 6b_x x_1! - 2x_1!_+^2 + 4x_1!_+^2 \\ & + 2x (3b_x n_1! + n_1!_+^2 - 2n_1!_+^2) \cos 2\theta_0 \\ & + 4n_1! \sin \theta_0] \cos \theta_0 B_1 n_1!_+ \\ & + [2n_1! (n_1^2!_+ + 2B_2 b_x x_1) \\ & B_1 x_1 (4b_x n_1! - n_1!_+^2)] \sin \theta_0 \\ & 4n x_1! [3b_x n_1! + n_1!_+^2 - 2n_1!_+^2] \sin^2 \theta_0 \\ & + 4x^2! [4b_x^2 n_1! + 4b_x n_1!_+ (n_1!_+^2 + n_1!_+^2) \\ & - (2b_x^2 n_1^2!_+ + n_1!_+^2)] \sin^3 \theta_0] g = f 2B_1 (A_1 + 4x_1!^2) \\ & + 8!^2 [A_1 x + (n_1 - 2b_x \cos \theta_0)^2 \\ & + 4x^2!^2] \sin^2 \theta_0 + 32b_x x_1!^2 (2b_x \cos \theta_0 - n_1) \sin^3 \theta_0 \\ & + 32b_x^2 x_1^2!^2 \sin^4 \theta_0 g; \end{aligned}$$

$$\begin{aligned} q_2 = & f_1 (A_1 + 4x_1!^2) B_2 n_1! \cos \theta_0 \\ & + 2x_1! (b_x \sin 2\theta_0 + b_z \cos 2\theta_0) \\ & 2x_1!_+^2 \sin 2\theta_0] - \sin \theta_0 [n_1 \\ & 2x (b_z \cos \theta_0 + b_x \sin \theta_0)] [2n_1 n_1! \cos \theta_0 \\ & 4x_1 n_1! (b_x \sin 2\theta_0 + b_z \cos 2\theta_0) \\ & + x_1^2!_+^2 \sin 2\theta_0 - 2A_4 \frac{x_1^2}{1} - 2A_2 \frac{x_1^2}{1} g \\ & = f 8!^2 \sin^2 \theta_0 [n_1 - 2x (b_z \cos \theta_0 + b_x \sin \theta_0)]^2 \\ & + 2 (A_1 + 4x_1!^2) (4x_1!^2 \sin^2 \theta_0 - B_1) g; \end{aligned}$$

-
- [1] I. A. A.eck, Phys. Rev. Lett. 46, 388 (1981)
 [2] A. D. Linde, Phys. Lett. B 100, 37 (1981)
 [3] E. M. Chudnovsky and Phys. Rev. A 46, 8011 (1992)
 [4] D. A. Gorokhov and G. B. Latter, Phys. Rev. B 56, 3130

- (1997)
 [5] E. M. Chudnovsky and D. A. Garanin, Phys. Rev. Lett. 79, 4469 (1997)
 [6] D. A. Garanin and E. M. Chudnovsky, Phys. Rev. B 56,

- 11102 (1997)
- [7] H. J. W. Müller-Kirsten, D. K. Park, and J. M. S. Rana, Phys. Rev. B 60, 6662 (1999)
- [8] J.-Q. Liang, H. J. W. Müller-Kirsten, D. K. Park, and F. Zimmermann, Phys. Rev. Lett. 81, 216 (1998)
- [9] G. H. Kim, Phys. Rev. B 59, 11847 (1999)
- [10] G. H. Kim and E. M. Chudnovsky, Europhys. Lett. 52, 681 (2000)
- [11] D. A. Garanin, X. Martinez Hidalgo, and E. M. Chudnovsky, Phys. Rev. B 57, 13639 (1998)
- [12] D. A. Garanin and E. M. Chudnovsky, Phys. Rev. B 59, 3671 (1999)
- [13] X. Martinez Hidalgo and E. M. Chudnovsky, J. Phys. Condens. Matter 12, 4243 (2000)
- [14] B. Zhou, J.-Q. Liang, and F.-C. Pu, Physica B 301, 180 (2001)
- [15] S. Y. Lee, H. J. W. Müller-Kirsten, D. K. Park, and F. Zimmermann, Phys. Rev. B 58, 5554 (1998)
- [16] A. Caneschi, D. Gatteschi, C. Sangregorio, R. Sessoli, L. Sorace, A. Comia, M. A. Novak, C. Paulsen, and W. Wernsdorfer, J. Magn. Magn. Mater. 200, 182 (1999)
- [17] G. H. Kim, Europhys. Lett. 51, 216 (2000)
- [18] B. Zhou, J.-Q. Liang, and F.-C. Pu, Phys. Rev. B 64, 132407 (2001)
- [19] G. H. Kim, Phys. Rev. B 67, 144413 (2003)
- [20] E. M. Chudnovsky and D. A. Garanin, Phys. Rev. Lett. 87, 187203 (2001)
- [21] K. Park, M. A. Novotny, N. S. Dalal, S. Hill, and P. A. Rikvold, Phys. Rev. B 65, 014426 (2002)
- [22] B. Parks, J. Loomis, E. Rumberger, D. N. Hendrickson, and G. Christou, J. Appl. Phys. 91, 7170 (2002)
- [23] K. M. Mertes, Y. Suzuki, M. P. Sarachik, Y. Paltiel, H. Shtrikman, E. Zeldov, E. Rumberger, D. N. Hendrickson, and G. Christou, Phys. Rev. Lett. 87, 227205 (2001)
- [24] J. M. Hernandez, F. Torres, J. Tejada, and E. Molins, Phys. Rev. B 66, 161407 (2002)
- [25] E. M. Chudnovsky, J. Magn. Magn. Mater. 140-144, 1821 (1995); B. Barbara and E. M. Chudnovsky, Phys. Lett. A 145, 205 (1990)
- [26] W. Wernsdorfer, E. Bonet Orozco, K. Hasselbach, A. Benoit, D. Mailly, O. Kubo, H. Nakano, and B. Barbara, Phys. Rev. Lett. 79, 4014 (1997)

$$-V(\theta, \phi = 0)$$









