

Novel Families of Fractional Quantum Hall States : A Berry Phase Approach

B. Basu* and P. Bandyopadhyay†
Physics and Applied Mathematics Unit
Indian Statistical Institute
Kolkata-700108

We have studied here the newly observed families of fractional quantum Hall states in the framework of Berry phase. It has been shown that this approach embraces in a unified way the whole spectrum of quantum Hall states with their various characteristic features. The newly observed states can be well accommodated within the primary sequence of FQH states and need not be considered as *second generation* FQH states as suggested by some authors.

PACS numbers: 73.43.-f; 73.43.Cd; 03.65.Vf

I. INTRODUCTION

The recent experiments [1] on high mobility samples at very low temperatures predicted the existence of fractional quantum Hall (FQH) states at some unusual filling factors such as $\nu = 4/11, 5/13, 5/17, 6/17, 4/13$ and $7/11$. For the states $\nu = 4/11$ and $5/13$ a deep minimum in ρ_{xx} and also a respectable plateau in ρ_{xy} was observed. But for $\nu = 6/17, 4/13, 5/17$ and $7/11$ states there was no plateau in ρ_{xy} and also minimum in ρ_{xx} was not pronounced. It has been observed that the new state at $\nu = 4/11$ is a fully polarized FQH state. A weak state was also observed in $\nu = 3/8$ and at $\nu = 3/10$. These sequences of fractions do not fit into the standard series [2] of integral quantum Hall effects (IQHE) of composite fermions (CF) at $\nu = \frac{p}{2mp \pm 1}$, but the states $\nu = 4/11$ or $4/13$ appear in the hierarchy of quasiparticle condensates [3, 4]. However, others such as $\nu = 3/8$ or $3/10$ do not belong to this hierarchy and the origin of their incompressibility seems to be puzzling. It has been proposed [1] that these states may be regarded as FQHE of composite fermions attesting to residual interactions between these composite particles. Finite size diagonalization of small clusters of electrons by Mandal and Jain [5] and Chang et.al.[6] suggest that the state at $\nu = 4/11$ is partially polarized which differs from the experimental findings of Pan et.al.[1].

A significant analysis has been done by Quinn et. al.[7, 8] in which they interpret these states as a novel family of FQH states involving pairing correlations among the quasiparticles (QP). The correlations depend upon the behaviour of the QP-QP pseudopotential $V_{QP}(L')$ where the interaction energy of a pair depends on the angular momentum. These pairs are proposed to have Laughlin correlations with one another and to form condensed states at a sequence which include all new fractions found in experiments.

Besides, it is also proposed [9] that the new composite particles consist each of a composite fermion of the first generation and a vortex like excitation which is based on the framework of Hamiltonian theory of Murthy and Shankar[10]. Very recently, Lopez and Fradkin [11] have proposed that the new states may be viewed as the hierarchical Jain states such that these are the quasiparticles and quasiholes of the primary states of the Jain sequence. Indeed, they have constructed the FQH states observed by Pan. et. al. as the fully polarized hierarchical descendants of the Jain series. All these approaches suggest that these new states effectively represent the **second generation** of FQH states as these are expressed as FQH states of FQH states.

In this note we shall show that all these new states may be treated as the FQH states in the primary sequence when we analyze the series from the viewpoint of Berry phase. Our analysis suggests the existence of FQH states which include apart from the states observed by Pan et. al. as well as all the predicted states by Quinn et. al. and also by Lopez and Fradkin. Besides, in this formalism these states are found to be fully polarized and their particle-hole conjugate states are found to be unpolarized. This

*Electronic address: banasri@isical.ac.in

†Electronic address: pratul@isical.ac.in

is consistent with the observation in experiments for the state with $\nu = \frac{4}{11}$ and its particle-hole conjugate state $\frac{7}{11}$.

In some earlier papers [12, 13] we have analyzed the sequence of quantum Hall states from the viewpoint of chiral anomaly and Berry phase. To this end, we have taken into account quantum Hall states on the two dimensional surface of a 3D sphere with a magnetic monopole of strength μ at the centre. In this spherical geometry we can analyze quantum Hall states in terms of spinor wave functions and take advantage of the analysis in terms of chiral anomaly which is associated with the Berry phase. In this geometry the angular momentum relation is given by

$$\mathbf{J} = \mathbf{r} \times \mathbf{p} - \mu \hat{\mathbf{r}}, \quad \mu = 0, \pm 1/2, \pm 1, \pm 3/2, \dots \quad (1)$$

From the description of spherical harmonics $Y_\ell^{m,\mu}$ with $\ell = 1/2$, $|m| = |\mu| = 1/2$, we can construct a two-component spinor $\theta = \begin{pmatrix} u \\ v \end{pmatrix}$ where

$$\begin{aligned} u &= Y_{1/2}^{1/2,1/2} = \sin \frac{\theta}{2} \exp[i(\phi - \chi)/2] \\ v &= Y_{1/2}^{-1/2,1/2} = \cos \frac{\theta}{2} \exp[-i(\phi + \chi)/2] \end{aligned} \quad (2)$$

Here μ corresponds to the eigenvalue of the operator $i \frac{\partial}{\partial \chi}$.

Then the N -particle wave function for the quantum Hall fluid state at $\nu = \frac{1}{m}$ can be written as [3, 12]

$$\psi^{(m)}_N = \prod_{i < j} (u_i v_j - u_j v_i)^m \quad (3)$$

m being an odd integer. Here $u_i(v_j)$ corresponds to the i -th (j -th) position of the particle in the system.

It is noted that $\psi^{(m)}_N$ is totally antisymmetric for odd m and symmetric for even m . Following Haldane [3] we can identify m as $m = J_i + J_j$ for the N -particle system where J_i is the angular momentum of the i -th particle. It is evident from eqn. (1) that with $\mathbf{r} \times \mathbf{p} = 0$ and $\mu = \frac{1}{2}$ we have $m = 1$ which corresponds to the complete filling of the lowest Landau level. From the Dirac quantization condition $e\mu = \frac{1}{2}$, we note that this state corresponds to $e = 1$ describing the IQH state with $\nu = 1$.

The next higher angular momentum state can be achieved either by taking $\mathbf{r} \times \mathbf{p} = 1$ and $|\mu| = \frac{1}{2}$ (which implies the higher Landau level) or by taking $\mathbf{r} \times \mathbf{p} = 0$ and $|\mu_{eff}| = \frac{3}{2}$ implying the ground state for the Landau level. However, with $|\mu_{eff}| = \frac{3}{2}$, we find the filling fraction $\nu = \frac{1}{3}$ which follows from the condition $e\mu = \frac{1}{2}$ for $\mu = \frac{3}{2}$. Generalizing this we can have $\nu = \frac{1}{5}$ with $|\mu_{eff}| = \frac{5}{2}$.

It may be remarked that as μ here corresponds to the monopole strength, we can relate this with the Berry phase. Indeed $\mu = \frac{1}{2}$ corresponds to one flux quantum and the flux through the sphere when there is a monopole of strength μ at the centre, is 2μ . The Berry phase of a fermion of charge q is given by $e^{i\phi_B}$ with $\phi_B = 2\pi q$ (number of flux quanta enclosed by the loop traversed by the particle). It may be mentioned that for a quantum Hall particle the charge is given by $-\nu e$ when ν is the filling factor.

If μ is an integer, we can have a relation of the form

$$\mathbf{J} = \mathbf{r} \times \mathbf{p} - \mu \hat{\mathbf{r}} = \mathbf{r}' \times \mathbf{p}' \quad (4)$$

which indicates that the Berry phase associated with μ may be unitarily removed to the dynamical phase. Evidently, the average magnetic field may be considered to be vanishing in these states. The attachment of $2m$ vortices (m an integer) to an electron effectively leads to the removal of Berry phase to the dynamical phase. So, FQH states with $2\mu_{eff} = 2m + 1$ can be viewed as if one vortex line is attached to the electron. Now we note that for a higher Landau level we can consider the Dirac quantization condition $e\mu_{eff} = \frac{1}{2}n$, with n being a vortex of strength $2\ell + 1$. This can generate FQH states having the filling factor of the form $\frac{n}{2\mu_{eff}}$ where both n and $2\mu_{eff}$ are odd integers. Indeed, we can write the filling factor as [12, 13]

$$\nu = \frac{n}{2\mu_{eff}} = \frac{1}{\frac{2\mu_{eff}}{n} \mp 1 \pm \frac{1}{n}} = \frac{n}{2mn \pm 1} \quad (5)$$

where $2\mu_{eff} \mp 1$ is an even integer given by $2mn$.

In this scheme, the FQH states with ν having the form

$$\nu = \frac{n'}{2mn' \pm 1} \quad (6)$$

(n' an even integer) can be generated through particle-hole conjugate states

$$\nu = 1 - \frac{n}{2mn \pm 1} = \frac{n(2m-1) \pm 1}{2mn \pm 1} = \frac{n'}{2mn' \pm 1} \quad (7)$$

Now, from eqn.(5),

$$\frac{n}{2\mu_{eff}} = \frac{n}{2m' \pm 1} = \frac{n}{2mn \pm 1} \quad (8)$$

we note that the integer m' has been taken to be the product mn . However, not all m' can be written in the product form mn . So we can write in a generalized way,

$$m' = mn \pm \tilde{m}, \quad \tilde{m} \text{ being an integer}$$

Indeed replacing n by q where q is any integer we can express the generalized relation as

$$\frac{q}{2\mu_{eff}} = \frac{q}{2m' \pm 1} = \frac{n}{2mq \pm (2\tilde{m} \pm 1)} \quad (9)$$

where we have taken $m' = mq \pm \tilde{m}$, with \tilde{m} being an integer. The relation (5) can be expressed as

$$\nu = \frac{1}{2m \pm \left(\frac{2\tilde{m} \pm 1}{q}\right)} = \frac{1}{2m \pm p/q} \quad (10)$$

where $p \geq 1$ an odd integer. It is noted that for $p = 1$ implying $\tilde{m} = 0$ corresponds to the Jain sequence.

The even denominator filling factors are obtained when μ_{eff} is an integer. As we have pointed out that for integer μ , the Berry phase can be removed to the dynamical phase, these states can only be observed when they appear as pair states [14, 15]. This corresponds to the non-Abelian Berry phase and represents non-Abelian quantum Hall fluid. In this case, the relation (10) will take the form

$$\nu = \frac{1}{2m \pm \frac{2\tilde{m}}{q}} = \frac{1}{2m \pm p/q} \quad (11)$$

where p (q) is an even(odd) integer.

Now to have a physical interpretation of the states given by (10) and (11) we note that these correspond to the attachment of p vortices (flux quanta) in a cluster of q electrons in the lowest Landau level.

For $p > 1$ and q odd we can take

(i) $q - 2$ electrons of which each one is attached with a vortex is coupled with a residual boson composed of two electrons

(ii) $q - 1$ electrons, each of them being attached with a vortex is coupled with the residual fermion.

In a similar way for q even, we may view the relation (10) such that $q - 1$ electrons, each having one vortex attached with it is coupled with the residual fermion. It is noted that when an electron is attached with a magnetic flux, its statistics changes and it is transformed into a boson. These bosons condense to form a cluster which is coupled with the residual fermion or boson composed of two fermions. Indeed the residual boson or fermion will undergo a "statistical" interaction tied to a geometric Berry phase effect that winds the phase of the particle as it encircles the vortices. This suggests that for odd q , p takes the values $q - 2$ and $q - 1$ and for even q , p is given by $q - 1$. Also we observe that the attachment of vortices to electrons in a cluster will make the fluid an incompressible one. Indeed as two vortices cannot be brought very close to each other, there will be a hard core repulsion in the system which accounts for the incompressibility of the quantum Hall fluid.

Now we consider some specific cases:

A : q odd, $p > 1$ and $p = q - 1$ or $q - 2$

q	p	general form of ν	values of m	ν
3	2	$\frac{1}{2m \pm 2/3}$	1	3/8 , 3/4
			2	3/14, 3/10
5	3	$\frac{1}{2m \pm 3/5}$	1	5/13 , 5/7
			2	5/23, 5/17
7	4	$\frac{1}{2m \pm 4/5}$	1	5/14, 5/6
			2	5/24, 5/16
7	5	$\frac{1}{2m \pm 5/7}$	1	7/19, 7/9
			2	7/33, 7/23
7	6	$\frac{1}{2m \pm 6/7}$	1	7/20, 7/8
			2	7/34, 7/22

B : q even, $p > 1$ and odd

q	p	general form of ν	values of m	ν
4	3	$\frac{1}{2m \pm 3/4}$	1	4/11, 4/5
			2	4/19, 4/13,
6	5	$\frac{1}{2m \pm 5/6}$	1	6/17 , 6/7
			2	6/29, 6/19

Values with bold faces in the above Tables have been reported in ref. [1]. The values $\nu = 4/5$ and $\nu = 5/7$ have been observed by Du. et. al.[16] which appear as weak depressions in the longitudinal resistivity. In a similar way, we can carry on with other q and p -values.

In this context we may add that the states we have obtained include all the states predicted by Quinn et al.[7, 8]. Indeed, their classification scheme suggests the relations

$$\nu^{-1} = 2\tilde{p} + 1 \pm (2 + \tilde{q}/2)^{-1} \quad (12)$$

where \tilde{p} and \tilde{q} are integers.

As mentioned earlier, the even denominator states are expected to appear as paired states. Indeed, in this case μ_{eff} is an integer and hence can be removed to the dynamical phase. These states can only be observed in paired states. This suggests that the newly observed states $\nu = 3/8$ and $\nu = 3/10$ should appear in paired states which has also been suggested by some other authors [7, 8, 17]. From our analysis, it appears that these states correspond to non-Abelian Berry phase and represent non-Abelian quantum Hall fluid.

In a recent paper [18] we have analyzed the polarization of quantum Hall states in the framework of the hierarchical analysis in terms of the Berry phase. There it is observed that the states in the lowest Landau level corresponding to the filling factors $\nu = 1$ and $\nu = \frac{1}{2m+1}$ with m an integer correspond to the fully polarized states. Indeed, in such a system even number of vortices are gauged away and the attachment of one vortex (magnetic flux) to an electron leads to the fully polarized state. However, in the higher Landau level we have the filling factor given by

$$\nu = \frac{n}{2\mu_{eff}} = \frac{1}{\frac{2\mu_{eff} \mp 1}{n} \pm \frac{1}{n}} = \frac{1}{2m \pm \frac{1}{n}} = \frac{n}{2mn \pm 1}$$

where n is an odd integer corresponding to a vortex of of strength $2l + 1$. As this effectively corresponds to the attachment of $\frac{1}{n}$ flux unit attached to an electron, the state will be partially polarized. Again the particle-hole conjugate states given by

$$\nu = 1 - \frac{n}{2mn \pm 1},$$

n being an odd integer will correspond to unpolarized states. This analysis suggests that the newly observed states as reported in [1] which appear in the lowest Landau level will correspond to fully polarized states. Indeed, the states corresponding to $\nu = 4/11$ has been observed to be fully polarized and the particle-hole conjugate state $\nu = 7/11$ is found to be unpolarized [1].

Finally, we may mention here that the hierarchical interpretation of the Haldane-Halperin scheme was questioned [19] because of the specific form of the QP-QP interaction. Besides, as mentioned earlier, the newly observed states $\nu = 3/8$ and $\nu = 3/10$ do not belong to this hierarchy. On the other hand, the Jain classification scheme effectively reveals a fundamental connection between IQHE and FQHE as the FQH states are considered to be IQH states of composite fermions. However, to interpret the newly observed states this fundamental concept has to be abandoned and we have to take into account the residual CF-CF interaction. A specific form of this interaction [7, 8] based on the pseudopotentials such that the pair interaction energy depends on the relative angular momentum nicely explains the new states. But some other states observed by Du. et.al [16] which appear as weak depression in the longitudinal resistivity cannot be accommodated in this scheme. Our present analysis suggests that these *second generation* FQH states are not truly second generation states, rather these appear in the primary sequence of the FQH states. This classification scheme can explain all the states observed to date [1, 16]. This also predicts the polarization of the states $\nu = 4/11$ and $\nu = 7/11$ consistent with experiment.

-
- [1] W. Pan, H. L. Störmer, D. C. Tsui, K. W. Baldwin and K. W. West, Phys. Rev. Lett. **90**, 016801 (2003).
 - [2] J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
 - [3] F. D. M. Haldane, Phys. Rev. Lett. **51**, 605 (1983).
 - [4] B. I. Halperin, Phys. Rev. Lett. **52**, 1583 (1994)
 - [5] S. S. Mandal and J. K. Jain, Phys. Rev. **B66**, 155302 (2002).
 - [6] C. C. Chang, S. S. Mandal and J. K. Jain, Phys. Rev. **B67**, 121305 (2003).
 - [7] J. J. Quinn, A. Wojs, K. S. Yi, cond-mat/0309435
 - [8] A. Wojs, K. S. Yi and J. J. Quinn, cond-mat/0304130
 - [9] M. O. Goerbig, P. Lederer, C. M. Smith, cond-mat/0308021
 - [10] G. Murthy and R. Shankar, Phys. Rev. **B 65**, 245309 (2002)
 - [11] A. Lopez, E. Fradkin, cond-mat/0310128
 - [12] B. Basu and P. Bandyopadhyay, Int. J. Mod. Phys. **B 12**, 2649 (1998).
 - [13] B. Basu and P. Bandyopadhyay, Int. J. Mod. Phys. **B 12**, 49 (1998).
 - [14] B. Basu and P. Bandyopadhyay, Int. J. Mod. Phys. **B 11**, 2707 (1997).
 - [15] B. Basu, D. Banerjee and P. Bandyopadhyay, Phys. Lett. **A 236**, 125 (1997).
 - [16] R. R. Du, H. L. Störmer, D. C. Tsui, A. S. Yeh, L. N. Pfeiffer and K. W. West, Phys. Rev. Lett. **73**, 3274 (1994)
 - [17] V. W. Scarola, J. K. Jain and E. H. Rezayi, Phys. Rev. Lett. **88**, 216804 (2002)
 - [18] B. Basu, S. Dhar and P. Bandyopadhyay : Polarization of Quantum Hall States, Skyrmions and Berry Phase (submitted for publication) : cond-mat/0301368.
 - [19] A. Wojs and J. J. Quinn, Phys. Rev. **B 61**, 2846 (1999)