

# Spin nematics in the bilinear-biquadratic S=1 spin chain

A. Läuchli,<sup>1,\*</sup> G. Schmid,<sup>2</sup> and S. Trebst<sup>2,3</sup>

<sup>1</sup>Laboratoire de Physique Théorique, CNRS UMR-5152, Univ. Paul Sabatier, F-31062 Toulouse, France

<sup>2</sup>Theoretische Physik, ETH Hönggerberg, CH-8093 Zürich, Switzerland

<sup>3</sup>Computational Laboratory, ETH Zentrum, CH-8092 Zürich, Switzerland

(Dated: March 22, 2022)

We report the existence of an extended critical, nondimerized region in the phase diagram of the bilinear-biquadratic spin-one chain. The dominant power law correlations are ferroquadrupolar *i.e.* spin nematic in character. Another known critical region is also characterized by dominant quadrupolar correlations, although with a different wave vector. Our results show that spin nematic correlations play an important role in quantum magnets with spin  $S \geq 1$  in regions between antiferromagnetic and ferromagnetic phases.

PACS numbers: 75.10.Jm, 75.10.Pq, 75.40.Mg, 75.40.Cx

The recent experimental demonstration [1] of the transition to a Mott insulating state of atoms in an optical lattice from a superfluid state has opened the way to novel realizations of effective quantum lattice models with widely tunable control parameters. Quantum magnetic systems can be realized by spinor atoms in an optical lattice, e.g.  $^{23}\text{Na}$  with a total  $S = 1$  moment. Confining  $S = 1$  atoms to an optical lattice there are two scattering channels for identical atoms with total spin  $S = 0, 2$  which can be mapped to an effective bilinear and biquadratic spin interaction [2, 3]:

$$H = \sum_{\langle i,j \rangle} \left[ J (\mathbf{S}_i \cdot \mathbf{S}_j) + J_q (\mathbf{S}_i \cdot \mathbf{S}_j)^2 \right], \quad (1)$$

where we adopt the standard parametrization  $J = \cos \theta$  and  $J_q = \sin \theta$ . In one dimension, the bilinear-biquadratic spin-one model has a rich phase diagram (see Fig. 1), which includes the well-known Haldane gap phase [4], a dimerized phase [5], a ferromagnetic phase and - as we will show in this Letter - two critical spin nematic phases. In a pioneering paper [6] Chubukov suggested the existence of a gapped, nondimerized phase with dominant spin nematic correlations close to the ferromagnetic region of the phase diagram. Subsequent numerical work [7] could however not substantiate this claim and it was therefore believed that the dimerized phase would extend up to the ferromagnetic phase boundary. Recent quantum Monte Carlo calculations [8] and field theoretical work [9] suggest that this picture might need to be reconsidered, especially in the light of possible experimental verifications in Bose-Einstein condensate systems [2].

In dimensions higher than one, the bilinear-biquadratic spin-one model is well understood in the relevant parameter regime. It exhibits a spin nematic phase with a finite spin quadrupole moment in the groundstate [10, 11].

In this Letter we propose a novel scenario for the phase diagram of the one-dimensional model in between the dimerized and the ferromagnetic phases. In particular, we substantiate the existence of an extended critical,

nondimerized region with dominant spin nematic correlations using extensive numerical simulations. We further characterize a second extended critical phase between the Haldane and the ferromagnetic phase, as also having prevalent quadrupolar correlations, albeit with a different wavevector.

The outline of this paper is as follows. In the first part we present various numerical results providing evidence for an extended critical phase without spontaneous dimerization for  $\theta \in (-3\pi/4, -0.67\pi]$ . We show that the quintuplet gap vanishes over this whole interval and that the transition between the dimerized and the critical phase is of a generalized Beresinski-Kosterlitz-Thouless (BKT) type. The absence of finite dimerization is discussed. We further determine conformal field theory parameters, such as scaling dimensions and the central charge for the critical region. Finally we address the second extended critical phase  $\theta \in [+ \pi/4, +\pi/2]$ , which we exhibit to have prevailing quadrupolar correlations at wavevectors  $\pm 2\pi/3$ .

*Numerical techniques* — Our results constitute the




FIG. 1: Phase diagram of the bilinear-biquadratic spin-one chain. The new elements are (a) an extended, critical phase with dominant ferroquadrupolar correlations for  $-3\pi/4 < \theta \lesssim -0.67\pi$  and (b) the characterization of the other critical region  $\pi/4 \leq \theta < \pi/2$  as having dominant  $k = \pm 2\pi/3$  quadrupolar correlations.




FIG. 2: Extrapolated DMRG  $S = 2$  gaps as a function of  $\theta$ . The gray region denotes the estimated onset of the critical behavior. The filled diamond is the exactly known gap result at  $\theta = -\pi/2$ . Inset: finite size extrapolation of the  $S = 2$  gaps (32 to 512 sites).  $\theta/\pi$  varies from  $-0.5$  to  $-0.7$  with decrements of  $0.05$  from top to bottom.

synthesis of different computational methods: i) Exact diagonalization (ED) of periodic chains up to 18 sites, using both static and dynamic observables, ii) Large scale Density Matrix Renormalization Group (DMRG) [12] calculations using the finite size algorithm on systems up to 512 sites while keeping 1000 states, iii) Strong coupling series expansions [13] around the dimerized limit up to 10th order.

*Closing of the Gap* — We first track the evolution of the energy gap to the first excitation using DMRG calculations on long open chains. A phase transition is signaled by the closing of the gap. The lowest excited state on open chains carries spin two for  $\theta \in (-3\pi/4, -\pi/2]$  [7]. Our results shown in Fig. 2 are clearly consistent with a finite gap of the  $S = 2$  excitation for  $\theta \geq -0.65\pi$ . However, for  $\theta < -0.65\pi$  the extrapolated gap is very small (of the order of  $10^{-3}$  for  $\theta = -0.7\pi$ ) which suggests the existence of a phase transition around  $-0.67\pi$  below which the gap is zero. The data shows no evidence for a reopening of the  $S = 2$  gap in the interval  $\theta \in (-3\pi/4, -0.67\pi]$ , at variance with the initial proposal [6]. In order to investigate the closing of the gap by different means we have calculated strong coupling series expansions of the  $S = 2$  single particle gap starting from the dimerized limit up to 10th order in the interdimer coupling  $\lambda$  for fixed  $\theta$ . A *direct* evaluation of the quintuplet gap shows a closing of the gap around  $\theta = -0.67\pi$  which is illustrated in the left panel of Fig. 3. Using DLog Padé approximants we have determined the critical interdimer coupling  $\lambda_c(\theta)$  where the gap closes, see upper right panel of Fig. 3. Consistent with previous estimates by Chubukov [6] we find that the gap closes for the uniformly coupled ( $\lambda = 1$ ) chain around  $\theta \approx -0.67\pi$ .

Having substantiated the closure of the gap in both DMRG and series expansions with indications of an extended gapless region  $-3\pi/4 < \theta < -0.67\pi$ , we now consider the existence of a generalized BKT phase transition.




FIG. 3: Strong coupling dimer expansion of the  $S = 2$  gap. Left panel:  $S = 2$  gap as a function of  $\theta$ . The gap closes at  $\theta_c \approx -0.67\pi$ . Upper right panel: Critical value  $\lambda_c$  where the gap vanishes. The uniformly coupled chain of interest here corresponds to  $\lambda = 1$  (dashed line). Lower right panel: Critical exponent of the  $S = 2$  gap opening as a function of the dimerization ( $\Delta \sim |\lambda - \lambda_c|^\gamma$ ) calculated by various DLog Padé approximants.




FIG. 4: Finite size scaling of the crossing points between the singlet gap at momentum  $\pi$  and the quintuplet ( $S = 2$ ) gap at momentum  $0$ . The extrapolation yields a critical value  $\theta_c \approx -0.67\pi$  in the thermodynamic limit.

To determine the critical point we use phenomenological level spectroscopy. We calculate the level crossing  $\theta_c^L$  of the lowest singlet excitation at  $k = \pi$  with the lowest spin-2 level at  $k = 0$  for different system sizes  $L$  with ED and extrapolate  $L \rightarrow \infty$ . The results are shown in Fig. 4. The extrapolation is performed with  $1/L$  and  $1/L^2$  corrections, fitting only the largest four system sizes. The extrapolated critical point is  $\theta_c = (-0.67 \pm 0.02)\pi$ , in good agreement with estimates obtained by other techniques [2, 6]. The coefficient of the  $1/L$  correction is small, and fitting without this term yields slightly larger values  $\theta_c > -0.67\pi$ .

*Dimerization and quadrupolar correlations* — Dimerization in the phase diagram of the spin-one chain has been firmly established by exact results obtained for the  $\theta = -\pi/2$  point [5]. It should be noted that although the system is dimerized, it is rather poorly de-




FIG. 5: Static and generalized susceptibilities for the bilinear dimerization (left panel) and the ferroquadrupolar correlations (right panel), obtained by ED on systems of 8 to 16 sites. Solid symbols: Static structure factors. Open symbols: Corresponding generalized susceptibilities.

scribed by a simple product wavefunction of alternating singlets, as can be seen from the small gap and the large correlation length over the whole dimerized phase. The dimerization order parameter [14] considered here is:  $\langle (\mathbf{S}_{i-1} \cdot \mathbf{S}_i) - \langle \mathbf{S}_i \cdot \mathbf{S}_{i+1} \rangle \rangle$ . The second kind of correlations expected to be important in this region of the phase diagram [10, 11] are the *quadrupolar* spin fluctuations  $\mathcal{C}_{\text{quad}}(r) \equiv \langle [(S_0^z)^2 - 2/3][(S_r^z)^2 - 2/3] \rangle$ .

We have calculated the standard static structure factor (SF) and the generalized nonlinear susceptibility (GNS) [15] for both kinds of correlations. We present ED results regarding the dimerization in the left panel of Fig. 5. The SF and the GNS both diverge as  $L \rightarrow \infty$  for  $\theta = -\pi/2$ , where long range dimer order is established [5]. Similar behavior is found for  $\theta \gtrsim -0.67\pi$ . For  $\theta \in (-3\pi/4, -0.67\pi)$  however, we find that the SF and the GNS are both *decreasing* with system sizes, indicating the absence of static dimerization in this gapless region. In the right panel we display the corresponding quantities for the  $k=0$  (*ferroquadrupolar*) mode of the quadrupolar correlations. Here the behavior is different: while there are only short range correlations deep in the dimerized phase, the ferroquadrupolar correlations increase drastically with system sizes for  $\theta \in (-3\pi/4, -0.67\pi)$ . Since quadrupolar long ranged order is forbidden at  $T=0$  in this 1D model – apart from  $\theta = -3\pi/4, +\pi/4$ , where the order parameter commutes with the Hamiltonian – we therefore expect a power-law behavior. Note that exactly at  $\theta = -3\pi/4$  the system is a  $SU(3)$  ferromagnet and shows ferromagnetism and ferroquadrupolar long range order at the same time [16].

Finally we want to discuss the relation of the proposed scenario to the numerical results presented in Ref. [7] where an intermediate phase as proposed by Chubukov [6] could not be found. We agree with the authors on the absence of a gapped, nematic-like phase. However, we




FIG. 6: Conformal field theory parameters calculated in ED on chains of 8 up to 18 sites. The scaling dimensions of the singlet ( $k = \pi$ ), and the quintuplet ( $k = 0$ ) field are shown. Inset: effective central charge. The value at the boundary of the critical region ( $\theta \approx -0.67\pi$ ) is  $c \approx 3/2$ .

think that the numerical results in [7] are compatible with the existence of an intermediate *critical*, nondimerized phase, though the authors have not explicitly considered such a phase.

*Field theoretical description* — In order to discriminate between potential conformal theories describing the phase transition and the extended gapless region, we calculate relevant field theory parameters, i.e. the scaling dimensions (SD)  $x$  of several fields and the central charge  $c$  (Fig. 6). The calculation of these parameters has been performed with ED and relies on finite size scaling properties of groundstate and excited states energies [17]. Leading logarithmic corrections have been taken into account. The SD of the  $k = \pi$  singlet field ( $x_s$ ) and the  $k = 0$  quintuplet field ( $x_q$ ) are both close to  $3/8$  at the onset of criticality near  $\theta \approx -0.67\pi$ .  $x_s$  significantly increases, while  $x_q$  decreases and seems to approach 0 as  $\theta \rightarrow -3\pi/4$ , compatible with quadrupolar long range order precisely at  $\theta = -3\pi/4$  [16]. Our results for the effective central charge are unexpected. The central charge  $c$  does not seem to be constant throughout the critical region. The smallest value is found at the onset ( $c \approx 3/2$ ), and then seems to increase monotonously. While this behavior can not directly be ruled out on field theoretical grounds, it is rather uncommon. It remains to be understood whether  $c$  is really continuously increasing or whether we are facing a crossover phenomenon. The critical theory at  $\theta \approx -0.67\pi$  we however believe to be well characterized by a level two  $SU(2)$  Wess-Zumino-Witten model given both the scaling dimensions ( $3/8$ ) and the central charge ( $3/2$ ). Further support for this claim comes from the indirect calculation of  $x_s$  within our series expansions. There the critical exponent  $\gamma$  of the  $S = 2$  gap is related to the scaling dimension  $x_s$  by  $\gamma = 1/(2 - x_s)$ . In the lower right panel of Fig. 3 we show the critical exponent calculated by DLog Padé approximants to the 10th order series. In the vicinity of  $\theta \approx -0.67\pi$  the various approximants reveal only small




FIG. 7: Static structure factors for spin and quadrupolar correlations at momentum  $k = 2\pi/3$  as a function of  $\theta$  for different system sizes. At the  $SU(3)$  point  $\theta = +\pi/4$  the two correlation functions are symmetry related. By going deeper into the critical region the spin correlations get weaker, while the quadrupolar correlations are enhanced. Inset: the central charge, the excitation velocity and the scaling dimension of the quintuplet field in the critical region.

spreading. Within the precision of a 10th order calculation the exponents comply with a scaling dimension of  $x_s = 3/8$  (dashed line).

*The period 3 phase* — The existence of an extended critical phase with soft modes at  $k = \pm 2\pi/3$  has first been reported in [18] and further analyzed and interpreted in [19, 20, 21]. Our results presented in the first part raise the question whether this critical region is of spin nematic character as well. Further hints come from the field theoretical work of Itoi and Kato [21], which shows that throughout the critical region the spin 2 mode at  $k = \pm 2\pi/3$  has correlation-enhancing logarithmic corrections. This is consistent with the earlier observation, that the lowest level at  $k = \pm 2\pi/3$  carries spin 2 [18]. We have calculated the static spin structure factor ( $S = 1$ ) and the spin quadrupolar structure factor ( $S = 2$ ) both at momentum  $2\pi/3$ . The results shown in Fig. 7 display nicely that the quadrupolar correlations become more important than the corresponding spin-spin correlations as one leaves the  $SU(3)$  symmetric point. Although all correlation exponents (dimer, spin, quadrupolar) are equal to  $\eta = 4/3$ , logarithmic corrections render the quadrupolar correlations dominant in this phase. Note that this phenomenon is similar to the  $S = 1/2$  antiferromagnetic Heisenberg chain, where both spin-spin and dimer-dimer correlations have equal correlation exponents  $\eta = 1$ , but the logarithmic corrections choose the spin-spin correlations to decay more slowly. For completeness we show selected field theory parameters for this critical region in the inset of Fig. 7. The results are in good agreement with field theoretical predictions [21] and support the applicability of the used ED techniques to determine field theory parameters.

*Conclusions and Outlook* — To summarize, we have shown that spin nematic correlations play an impor-

tant role in the phase diagram of the  $S = 1$  bilinear-biquadratic chain. Applying a combination of numerical techniques we reconsidered Chubukov's proposal [6] of a gapped, nondimerized and nematic-like phase between the dimerized and the ferromagnetic phases. Our numerical results provide evidence for a modified picture with a phase transition at  $\theta \approx -0.67\pi$  where the dimerized phase terminates. However, the phase in the window  $\theta \in (-3\pi/4, -0.67\pi]$  remains critical, i.e. gapless and nondimerized. The dominant correlations are spin-quadrupolar correlations at wavevectors  $k = \pm 2\pi/3$ . It will be interesting to compare our field theoretical parameters to future analytical calculations.

We would like to thank F. Alet, N. Laflorencie, P. Lecheminant, F. Mila, D. Roggenkamp, M. Troyer and S. Wessel for valuable discussions. The authors acknowledge support from the Swiss National Science Foundation.

---

\* Electronic address: laeuchli@comp-phys.org

- [1] C. Orzel *et al.*, Science **291**, 2386 (2001); M. Greiner *et al.*, Nature **415**, 39 (2002).
- [2] S.K. Yip, Phys. Rev. Lett. **90**, 250402 (2003).
- [3] A. Imambekov *et al.*, cond-mat/0306204.
- [4] F.D.M. Haldane, Phys. Lett. **93A**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
- [5] M.N. Barber and M.T. Batchelor, Phys. Rev. B **40**, 4621 (1989); A. Klümper, Europhys. Lett. **9**, 815 (1989); Y. Xian, Phys. Lett. A **183**, 437 (1993).
- [6] A.V. Chubukov, Phys. Rev. B **43**, 3337 (1991).
- [7] G. Fáth and J. Sólyom, Phys. Rev. B **51**, 3620 (1995).
- [8] N. Kawashima, Prog. Theor. Phys. Suppl. **145**, 138 (2002).
- [9] B.A. Ivanov and A.K. Kolezhuk, Phys. Rev. B **68**, 052401 (2003).
- [10] N. Papanicolaou, Nucl. Phys. **B305**, 367 (1988).
- [11] K. Harada and N. Kawashima, Phys. Rev. B **65**, 052403 (2002).
- [12] S.R. White, Phys. Rev. Lett. **69**, 2863 (1992).
- [13] M.P. Gelfand and R.R.P. Singh, Adv. Phys. **49**, 93 (2000).
- [14] We only discuss the bilinear dimerization, but we have checked that the biquadratic dimerization shows similar behavior regarding the phase boundaries.
- [15] The generalized nonlinear susceptibility is defined as the second derivative of the groundstate energy with respect to the addition of a symmetry breaking perturbation. This approach has been discussed in L. Capriotti, Int. J. Mod. Phys. B **15**, 1799 (2001).
- [16] C.D. Batista *et al.*, Phys. Rev. B **65**, 180402(R) (2002).
- [17] H.W.J. Blöte *et al.*, Phys. Rev. Lett. **56**, 742 (1986); I. Affleck, Phys. Rev. Lett. **56**, 746 (1986); J.L. Cardy, J. Phys. A: Math. Gen. **19**, L1093 (1986).
- [18] G. Fáth and J. Sólyom, Phys. Rev. B **44**, 11836 (1991).
- [19] Y. Xian, J. Phys. Condens. Matter **5** 7489, (1993).
- [20] A. Schmitt *et al.*, Phys. Rev. B **58**, 5498 (1998).
- [21] C. Itoi and M. Kato, Phys. Rev. B **55**, 8295 (1997).