Superconductors with Superconducting Transition Tem peratures Tc = 91K (1999), 120K (1994), 340K (2000), and 371K (1995): Experim ental Errors or a Technological Puzzle? Two-C om ponent N onstoichiom etric C om pounds and the Insulator{Superconductor{M etal Transition.

V.N.Bogom olov A.F.Io e Physical & Technical Institute, Russian Academy of Science, 194021 St. Petersburg, Russia (Dated: March 22, 2024)

O ne of the reasons for the lack of understanding of both the mechanism s underlying the H T SC phenom enon and of the instability of materials with $T_c > 300$ K may be the widely accepted but wrong ideas about the types of chem ical bonding in a substance and the radii of the atom s and ions. A revision of these concepts started in the beginning of the XX century in connection with the investigation of nonstoichiom etric compounds (the berthollides) but did not reach a critical level until recently. Most of the HTSC materials, how ever, are actually two-component nonstoichiom etric nanocom posites (the berthollides), whose com ponents "dilute" or "stretch" one another. Each component resides in an "interm ediate" state, which still remains poorly studied. For instance, in a system of particles having two paired electrons each (divalent particles), the unbroken electron pairs may start tunneling at a certain "medium" concentration (or in the tem perature interval $T_1 < T_c < T_2$;) with the system becoming a Bose superconductor (the state between the insulator and the metal with BCS superconductivity). For univalent atom s (Na, Aq,), however, such possibility realizes neither in the interm ediate (the M ott transition) nor in the nal state. Univalent m etals are not superconductors. In the berthollides, however, a possible Jahn {Teller{Peierls-type instability may give rise to form ation of diatom ic molecules (Na2, Ag2,) with electron pairs, and superconductivity can set in. It is possibly such system s that were obtained by chance in experiments with univalent components and reported to have T $_{\rm c}$ of up to 371K : Structures of a number of HTSC m aterials are considered.

PACS num bers: 71.30.+ h, 74.20.-z, 74.25.Jb

In the recent decade, several reports from independent sources announced synthesis of substances with the superconducting transition occurring at 91K [1], 120K [2], 340K [3], and 371K [4]. These experiments, how ever, neither have received con mation nor were continued. In all the cases, the reason for this was the poor reproducibility of the synthesis and the instability of the sam ples. It is this feature, how ever, that is the main structural characteristic of H T SC substances with the model of superconductivity used to interpret the metallization of xenon under pressure [5] and, later, applied to two-component nonstoichiom etric compounds with incommensurate parameters of the constituents (matrix system s) [6]. In addition to the instability common to all the four substances, they have also certain common physical features characteristic of the given superconductor model, which permits one to consider their appearance not as due to experimental errors but rather as chance occurrences of technological nature associated with structural instabilities.

Synthesis of superconductors with $T_c > 300$ K and elucidation of the mechanism s underlying high-tem perature superconductivity are issues that are still awaiting solution. One of the reasons for this may be the widely accepted but inadequate concepts concerning the part played by the atom ic and ionic states (and by their radii) in the structure of a substance. A revision of these concepts started in the beginning of the XX century in connection with the investigation of nonstoichiom etric compounds (berthollides). The lack of understanding of their nature has not, how ever, reached a critical level until recently. Indeed, we may recall that most of the HTSC compounds are actually berthollides, and their physical properties cannot be understood properly without a correct description of their chemical structure. Rather than being uniform chemical substances, the berthollides can be, for instance, twocomponent nonstoichiom etric nanocomposites, with each of the components "diluting" or "stretching" the other. They stabilize one another in an unusual, "interm ediate" state, which remains poorly studied. If, for instance, particles having two paired electrons each (divalent particles) approach one another, a real state may form within a narrow interval of concentrations (or tem peratures) between the insulator and the metal, a state where pair tunneling has already started, and where, while their binding energy has already begun to decrease, the electron pairs have not yet split into ferm ions (as is the case with the nal, metallic, equilibrium state for $T > T_c$): The system

E lectronic address: V Bogom olov@ m ail.io e.rssi.ru

transfers to the state of a Bose condensate. A Bose superconductor forms between the insulator and the metal with BCS superconductivity within a narrow and structurally unstable region. For univalent atoms (Na, Ag,), however, such possibility realizes neither in the intermediate (the M ott transition) nor in the nal state. Univalent metals are not superconductors. In the berthollides, however, a possible Jahn {Teller{Peierls-type instability may give rise to form ation of diatom ic molecules (Na₂, Ag₂,) with electron pairs and superconductivity can set in. It is possibly such systems that were obtained by chance in experiments with univalent components and reported to have T_c of up to 371K. In the berthollides, stability is a consequence primarily of their architecture rather than of chem ical interactions, a point that is frequently disregarded, thus bringing about a distorted picture of physical phenom ena. This may come from a wrong determ ination of the type of chem ical bonding in the given structure because of the use of wrong values of the radii and charges of the particles, a possibility substantiated by a com parison of experimental data with quantum -m echanical calculations of the radii of atom s and, particularly, of ions. For instance, the radius of the 0² ion is not 1:4A but 0:6A, and that of Cu²⁺; 0:33A rather than 0:73A [7, 8].

Two-component nonstoichiom etric compounds are relatively inert matrices with voids of diameter D containing not ions but atom s, whose diameters d are slightly larger than those of the free atom s but smaller than of atom s in a metal. W hen atom s are removed from the metals (or when the number of the nearest neighbors decreases), their diameter decreases by only 10% 15% (the Goldschmidt contraction). Each component is a stoichiom etric substance, whose structure may be not necessarily stable in itself. W hile atom s of metals, for instance, can have close-to-metallic radii, the component itself may turn out to be an unusual, "stretched" or "dilute" metal, and not a Ferm i system at all. Stoichiom etric components are separated by a "gap" taking up a sizable part of the nanocomposite volume [9].

(1) The case of D (3 10)d: Belonging here are, for instance, the zeolites with substances embedded in the structural voids [10]. The clusters in the voids are domains of the second phase. The system as a whole may be treated also as a "supercrystal" [11].

(2) The case of D . d: A tom s of the second component (but not ions) stretch the matrix. This case, well known in chem istry, relates directly to the nonstoichiom etric compounds (the berthollides or interstitial compounds). It can be illustrated by the M eB₆ system . A sublattice of the m etal atom s rather than of the matrix-doping ions is inserted into the voids of the fram ew ork between the B₆ octahedra. This is indicated by the fram ew ork stretching, with d, the diam eter of the atom, exceeding D, which is the M eB₆ lattice parameter. For Me = Ce, La, Ca, Sr, Ba the ratio D =d is Ce (4:14=3:26); La (4:15=3:74); Ca (4:15=3:95), Sr (4:19=4:30); and Ba (4:27=4:34): The sublattices of Ce, La, and C a are stretched, and those of Sr and Ba, com pressed, but at the sam e time the B₆ fram ew ork is stretched. M etal ions could not stretch the fram ew ork. All these com pounds feature about the sam e melting points T 2200K; which are determ ined by the rigid covalent B₆ fram ew ork [12]. The B {B bond lengths of the venearest neighbors are 1:72 A.

(3) If D & d (perovskite-like structures), the atom s of the second M e component inserted in the voids of the matrix M t can undergo a 3D Jahn {Teller{Peierls transition with form ation of diatom ic molecules made up of atom s which occupy neighboring voids [6]. A firer all the voids have lled (M q M t_1 ;) one can conceive of the following possibilities: (A) form ation of regular structures of diatom ic molecules;

(B) form ation of disordered structures.

In both cases, a matrix-stabilized second phase will appear. This phase is at the same time "diluted" by the matrix. If not all of the voids are led (a technological problem), the composition of the substance can be written as $M e_x M t_1$ for x < 1; which likewise suggests two possibilities:

(a) - dom ains with all voids lied will form (a version of cases A and B);

(b) - a component with a low concentration of diatom ic molecules form s.

Realization of version (b) is a still more complex technological problem, but it is such systems with very high T_c that were apparently obtained accidentally [1, 2, 3, 4].

I. If the inserted atom s are univalent, the diatom ic m olecules will have two paired electrons and a binding energy $E_m = E_0 (D - d) = d$, where E_0 is the binding energy of the M e m et al [6]. As the concentration N of such m olecules increases, or the tem perature decreases, the therm alde B roglie wavelengths dB T $^{1=2}$ of the electron pairs and the mean interm olecular distances N¹⁼³ may become comparable. It gives rise to Bose-Einstein condensation $\frac{3}{dB}$ N > 2:612 [13]. The condensation phenom enon of the Bose E instein statistics is distorted, of course, by the at presence of m olecular forses and by fact that it m anifests itself not in the gaseous state. But the sam e problem s exist in the case of liquid helium as well. The system will transfer from the insulator to the Bose superconductor state with the transition temperature $kT_{cn} = kT_B = 3.31 (h=2)^2 N^{2=3} = M = 1.04 (h=2)^2 n^{2=3} = m$; where n = 2Nand m = M = 2 are the electron concentration and mass, respectively. For $T_B = (15; 70; 325) K$, we obtain $n = (2 \ 1b^{9}; 2 \ 1b^{9}; 2 \ 1b^{9})$ cm³: The transition temperature can, how ever, be governed by the E_{m} energy as well, as is the case with m etals (T_{cm}). As the tem perature increases to $kT > kT_{cm}$ E_m ; or the concentration rises to $n > n_{crit}$ (ultimately to x = 1), the bosons will separate into the fermions, and the system will become Fermi type with the tem perature T_F > T_B: Therefore, such system s can become superconducting only within a narrow region of concentrations of atom s (m olecules) or tem peratures ($T_1 < T_c < T_2$). These are either doped sem iconductors,

or "stretched" or "diluted" m etals; being actually real system s, they are not Ferm i type but rather unstable, and it is such system s that are dealt with here. Such system s can become superconducting not only through form ation of electron pairs (T_{cm}) as is the case with m etals, but as a result of B ose condensation of the bosons present in the system ($T_{cn} = T_B$). An increase in the concentration of M e atoms drives the insulator{superconductor{m etal transition.

II. If two paired electrons, rather than belonging to molecules, are outer electrons of a divalent atom, for instance, of M g, the components need not necessarily be incommensurate. One has only to produce in the voids of an inert matrix a local concentration of the metal atoms M e_x within a certain interval x < 1 (provided there are no domains with x = 1, as is the case with an equilibrium Ferm imetal). The mechanism of preservation of electron pairs in metals, if they form for $T < T_{cm}$ from the outset as superconductors, is described by BCS theory (the C coper pairs). Univalent metals do not have such "relic" states, and this is what accounts for their not exhibiting superconductivity. C om pression of a metal usually reduces $T_c: H ence$, C coper pairs in a "stretched" metal are bound stronger. They are much closer to the state of atom ic electron pairs.

M gB₂. It is a version of such a "diluted m etal" that possibly is realized in the M gB₂ system. The divalent m agnesium is "diluted" by boron layers in the form of a covalently bonded hexagonal network. The M gB₂ lattice parameters are c = 3:40 A, a = 3.05 A [14]. Each B atom has three nearest neighbors. The B (B distance, 1:77 A, is larger than the dimensions of the atom, 1:55 A [8], than that in the B₂ m olecule (1:59 A), and that in M eB₆ (1:72 A), which has venearest neighbors. Having a nearly covalent diameter (3:08 A), the m agnesium atoms stretch the boron atom network. This produces conditions favorable for the form ation of B₂ m olecules. The M g atom s con ned between the B layers are also slightly stretched along the c axis as compared to their diameters in the m etal (3:20 A). This system revealed two energy gaps proportional to $T_{cm 1} = 15K$ and $T_{cm 2} = 45K$.

 $M e_3 C_{60}$: A nother illustration of superconductors with a stretched component (version D . d) is provided by the $M e_3 C_{60}$ fullerides. Their $T_c = T_{cm}$ $E_0 (D D_0) = D_0$; where E_0 is the binding energy of the C_{60} spheres, D_0 is the fullerene lattice parameter, and D is this parameter increased by incorporation of $M e_3$ clusters into the octahedral voids [6]. $T_{cm} a_X$ 40K.

 $Na_x CoO_2$ 1:3H₂O. The superconductivity (T_c 4K) observed in the $Na_x CoO_2$ 1:3H₂O system is due to the univalent sodium, possibly as a result of formation of weakly bound Na_2 molecules. The optimum concentration of Na is x = 0.3 [15], i.e., it is "interm ediate", as in many other cases. It is stabilized by the H₂O molecules (an "additional diluent" or third component).

 Na_x-NH_3 . The same molecules (Na_2) formed undoubtedly in the Na_x-NH_3 solutions as well (T_c 200 K) [16]. The instability of this system originates from fast phase separation in the liquid, and the decrease in the solution density can be traced to the increase of the atom ic volume of sodium in diatom ic molecules and the interphase "gap" [6, 9] rather than to the formation of "bubbles" [16].

 $Y_{2}Ba_{4}Cu_{6}O_{14}$. The well studied system $Y_2Ba_4Cu_6O_{14}$ (85K) [17] can also be treated in terms of the above scheme. Its lattice parameters are c = 11:681 A = 3 (3:894)A, a = 3:886 A, b = 3:827 A [17]. This structure m ay be considered also as a slightly distorted sim ple cubic lattice of spheres with an average diam eter of 3:86 A. It corresponds to the average diam eter of the Y and Ba atom s in these spheres of 3:87 A. The diam eters of the free atom same 3:38 A and 4:12 A, and those of the ions Y³⁺ and Ba²⁺; 1:28 A and 1:73 A. The diam eters of the atom s in the metalare 3:56 A and 4:34 A. The density of this lattice is 0:52 ("dilute metal"). The remainder of the volume, 0:48, is taken up by the Cu {O octahedra and pyramids and the interphase "gap" [9]. Each oxygen atom is also bonded covalently to six or eight nearest oxygen atom s. Sulfur, an analog of oxygen, also exhibits a tendency to form ation of structures of a molecular type (S_2, S_6, S_8) . In the case of oxygen, this process requires a "seed" (Cu) for its realization. The Cu₃O₇ component is a stoichiom etric compound with O {O covalent bonding [18]. It is stable naturally only in the framework of the Y-Ba₂ lattice. Both components stabilize one another. The Y and Ba sublattices are "diluted", and the Y sublattice, "stretched" as well (Y $_2$). Hence, the volume of the C u_3 O $_7$ com ponent is not governed by the ionic diam eters, either traditionally accepted (1:46 A and 2:8 A) or calculated (0:65 A and 1:20 A) [7,8].

N $a_{0:05}WO_3$: The properties of two-component nonstoichiom etric compounds become most clearly pronounced in the M e_xWO_3 tungsten bronzes [6]. The bronze with an average composition N $a_{0:05}WO_3$ was found to be superconducting with $T_c = 91K$ and to have an energy gap E 160K [1]. The substance is unstable. The parameter D = 3:78 A. In the N am etal, d = 3:72 A. In this case, for $E_0 = 1$ eV in sodium we obtain for the gap energy $E_m = E_0(D - d) = d = 0.016$ eV, which in plies that the transition is energy driven. Free N a_2 molecules have a binding energy of 0:35 eV. In the W O₃ matrix, their energy is 20 times smaller (0:35=0.016). Their detection is made di cult by the fact that the modulation of the period is only 0.016 . W O₃ is a standard stoichiom etric com pound (M $^{6+} - O^2$): How ever, in this compound each oxygen atom has likew ise eight nearest oxygen atoms. Therefore, its physico-chem ical bonding is also a quantum -m echanical superposition of several types of states [18]. Rem arkably, replacement of N a by Rb or C s, for which D < d , low ers sharply T_c (from 91K down to 2 3K). Substitution of N a by Lior Ag (d 3:10 A) was not tried, because N a, Rb, and C s were considered as dopants only (i.e., in the ionic form), rather than a second

component in a nonstoichiom etric compound [1].

Ag{Y {Ba{Cu{O.Substitution of this type was made earlier in the Y {Ba{Cu{O system . One doped it with silver, which raised T_c to 120K [2].

A $g_x P b_6 C O_9$. The use of silver in the A $g_x P b_6 C O_9$ system also brought about an increase in T_c to 340K [3]. The "colossal electrical conductivity > 10⁹ O hm⁻¹ cm⁻¹" observed to occur in the A $g_5 P b_2 O_6$ system between 210K and 525K is assigned [19] to a heating-induced segregation of the A g component in channels of the A $g_3 P b_2 O_6$ lattice. The composition proposed for this compound in [19] is A $g_2 A g_3 P b_2 O_6$]: If this state is indeed superconductivity, it becomes realized within the interval ($T_1 < T_c < T_2$) (i.e., under heating).

 $Y Ba_2Cu_3Se_7$. T_c can be increased not only by reducing d but by increasing D as well. The oxygen in the standard $Y \{Ba_4Cu_4O \text{ system was replaced by selenium, which brought about an increase in the lattice parameter, after which <math>Y Ba_2Cu_2Se_7$ exhibited T_c 371K [4].

O ne can conceive also of a speci c class of two-com ponent nanocom posites of a "dynam ic" type. These are condensates of m olecular, or noble gases. In these condensates, a certain number of virtual exciner m olecules, for instance, $X e_2 * [20]$ with two paired electrons, are present in the lattice of X e atom s residing in the ground state ("m atrix"). It is supposed, that increasing the $X e_2 *$ concentration either by applying pressure or through the action of catalysts (atom s of m etals) could give rise to B ose superconductivity with high T_c [5]. The investigation of the M ott transition in m etal{ inert-gas nanocom posites and of adsorption forces (sorption com pounds) is dealt with in a large number of publications (see, e.g., ([21, 22, 23, 24]). It is quite possible that a W -A r sorption com pound forms in the gas sheath surrounding the tungsten lament in the course of m etal vaporization in conventional gas- led tubes. M etallization of this layer brings about a drop in the voltage across the lament with increasing current [25]. O ne cannot exclude here, how ever, the possibility that the increase in the conductivity of the gas is actually the result of its ionization.

Thus, the major technological di culty in the way of developing HTSC materials consists in preparing stable systems of particles with paired electrons (for instance, divalent atom s or molecules) in concentrations at which the electron pairs already are capable of tunneling (B ose condensation) while not yet separating into single electrons, either as a result of conventional chem ical interactions or through a decrease in their binding energy in an elective dielectric medium. Such an intermediate state corresponds to a "stretched" or "diluted" substance, which can be stabilized by using "solid solvents" as matrices. The highest T_c were obtained with univalent metals (not superconductors), because their diatom ic molecules can exist in the matrices in low enough concentrations. In some cases such nonequilibrium concentrations are possibly "frozen" by chance uctuations in the technology of synthesis ([1, 2, 3, 4]). Stability has thus far been reached apparently at the expense of a low T_c : In searching for electric moment, for instance) a better understanding of the quantum -mechanical interactions in the structure of each component, for instance) a better understanding of the quantum -mechanical interactions in the structure of each component is of crucial importance. Such methods of stabilization can be found only by using realistic atom ic and ionic radii rather than conventional quantities, which are nothing else but a consequence of postulated bonding types. The matrices in such nanocom posites also exist in an unusual state because of the unusual structure and contact interaction with the second component. This complicates greatly the investigation and description of such systems.

[10] Bogom olov, V N. Liquids in ultrathin channels (thread and cluster crystals). Usp. Fiz. Nauk 124, 171 - 182 (1978). [Sov.Phys.Usp. 21, 77 - 86 (1978)].

^[1] Shengelaya, A., Reich, S., Tsabba, Y., and Muller, K.A. Electron spin resonance and magnetic susceptibility suggest superconductivity in Na doped W O₃ samples. Eur. Phys. J. B 12, 13-15 (1999).

^[2] Yanmaz, E., Multu, I.H., Kucukomeroglu, T., Altunbas, M. Ag-doped 120K YBa₂Cu₃O_{7 d} superconductors prepared by the ame-quench-melt-growth (FQMG) method Supercond. Sci. Technol. 7, 903 - 907 (1994).

^[3] D jurek, D., M edunic, Z., Tonejc, A., Paljevic, M. superconductivity from Pb₃CO₅ - Ag₂O (PACO) system Physica C 341-348, 723-725 (2000).

^[4] Shabetnik, V D., Butuzov, S.Yu., Plaksii, V.J. High-tem perature superconducting compound YBa₂Cu₃Se₇ with T_c = 371K. Techn. Phys. Lett. 21, 382 - 384 (1995).

^[5] Bogom olov, VN. M etallization of m olecular condensates and superconductivity. Techn. Phys. Lett. 28, 211-215 (2002).

^[6] Bogom olov, VN. Superconductivity of the two-component non-stoichiom etric compounds with incom ensurate sublattices. e-print, cond-m at/0304561.

^[7] Slater, J.C. A tom ic radii in crystals. J. Chem. Phys. 41, 3199-3204 (1964).

^[8] W aber, J.T., Cromer, Don T. Orbital radii of atom s and ions. J. Chem. Phys. 42, 4116-4123 (1965).

^[9] Bogom olov, V N. Capillarity phenom ena in extrem ely thin zeolite channels and m etal-dielectric interaction. Phys. Rev. 51, 17040 - 17045 (1995).

^[11] A rita, R., M iyake, T., K otan, T., Schilfgaarde, M. van., O ka, T., K uroki, K., N ozue, Y., A oki, H. E lectronic properties of alcalim etal loaded zeolites-"supercrystal" M ott insulator. e-print, cond-m at/0304322.

^[12] M andelcom, L. (ed) in Non-Stoichiom etric Compounds (A cadem ic Press, N.Y. and London, 1964)...

- [13] London, F.On the Bose Einstein condensation. Phys Rev., 54, 947-954 (1938).
- [14] Xu, M., Kitazawa, H., Takano, Y., Ye, J., Nishida, K., Abe, H., Matsishita, A., Kido, G. Single crystal M gB₂ with anisotropic superconducting properties. e-print, cond-m at/0105271.
- [15] Schaak, R E., K lim czuk, T., Fool, M L., Cava, R J. Superconductivity phase diagram of Na_xCoO₂ -1.3 H₂O .Nature 424, 527-529 (2003).
- [16] Ogg, R A Jr. Bose-Einstein condensation of trapped electron pairs. Phase separation and superconductivity of metalammonia solution. Phys. Rev. 69, 243-244 (1946).
- [17] Izum i, F., Asano, H., Ishigaki, T., Ono, A., Okamura, P.Crystal structure of Ba-Y-Cu-O superconductor as revealed by Rietveld analysis of x-ray powder di raction data. Jap. J. Appl. Phys. 26, L611-L612 (1987).
- [18] K rebs, H. G rundzuge D er Anorganischen K ristallchem ie (ed Enke, F., Stuttgart, 1968).
- [19] D jurek, D., M edunic, Z., Paljevic, M., Tonejc, A. Colossal electric conductivity in Ag-defect Ag₅Pb₂O₆. e-print, condm at/0310011.
- [20] Tilton, RA., Flynn, CP. Sharp optical spectra of in purities in metal. Phys. Rev. Lett. 34, 20-23 (1975).
- [21] Phelps, D. J., Avci, R., Flinn, C. P. Metal-insulator transition in metal-rare gas-alloys. Phys. Rev. Lett., 34, 23-26 (1975).
- [22] Endo, H., Eatah, A.J., Wright, J.G., Cusack, N.E.A. metal-nonmetal transition in argon-copper system. J. Phys. Soc. Of Japan, 34, 666-671 (1973).
- [23] Leung, A W K., Kaup, J.G., Bellert, D., M oG a rey, J.G., Breckenridge, W H. Spectroscopic characterization of excited Ca(4s4d 3D_J)RG (³_{1;2}) states (RG = Ar, Kr, Xe): no "heavy-atom " m ixing of RG (nd) character into the wave functions. J. Chem. Phys., 111, 981-987 (1999).
- [24] Burda, J.V., Runeberg, N., Pyykko, P. Chem ical bond between noble metals and noble gases. Ab initio study of neutral diatom ic NiXe, PdX e and PtX e. Chem. Phys. Lett., 288, 635–641 (1998).
- [25] Markov, G A., Melnikova, N.V., Khon, Yu. A nom alously high conductivity of deformed metals at elevated temperatures. Techn. Phys. Lett., 22, 680-681 (1996).