arXiv:cond-mat/0311302v1 [cond-mat.dis-nn] 13 Nov 2003

Density of states of disordered Dirac particles: Infinitely many operators with negative scaling dimensions and freezing transitions

Shinsei Ryu
Department of Applied Physics, University of Tokyo, 7-3-1 Hongo Bunkyo-ku, Tokyo 113-8656, Japan

Christopher Mudry
Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland and Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

Akira Furusaki
Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: March 22, 2022)

The global density of states (GDOS) close to the band center $\varepsilon = 0$ for a particle hopping on a square lattice and subjected to disorder that preserves the bipartite symmetry of the lattice is computed using field theoretical methods. The GDOS diverges like $|\varepsilon|^{-1}\exp\left(-c|\ln\varepsilon|^{\kappa}\right)$ with $\kappa = 2/3$ in agreement with a prediction by Motrunich *et al.*[8] and in disagreement with an older prediction by Gade[9] ($\kappa = 1/2$).

PACS numbers:

The problem of two-dimensional ($2D$) disordered Dirac particles, particles with a relativistic dispersion subjected to several types of quenched randomness, has attracted a lot of interest since 1994.[1] Indeed, it is related to a broad class of models such as the $2D$ random phase XY model, dirty d-wave superconductors, or the Chalker-Coddington model for the plateau transition in the integer quantum Hall effect, in which disorder plays an essential role. In spite of its relative simplicity, it can exhibit a rich variety of phenomena which appears to be specific to critical behavior induced by disorder. Multifractality is one example thereof.

Multifractal scaling implies that the scaling regime in the vicinity of criticality is encoded by an *infinite* set of independent relevant scaling exponents, the multifractal spectrum in short. On general grounds, the multifractal spectrum must fulfill the condition of monotonicity. Implementing this condition is a challenge to the common lore of *local* field theoretical techniques.

We consider a problem of Anderson localization defined on a $2D$ lattice where a particle is hopping on a square lattice in the background of π-flux phase and subjected to weak uncorrelated bond-disorder that preserves the sublattice symmetry and time-reversal invariance.[2] The trademark of this model, the HWK model, is the sublattice symmetry present for any given realization of the disorder. Numerical studies offer strong evidences that typical wavefunctions at the band center are neither localized nor delocalized but are multifractal[2] and that the density of states (DOS) is singular[3]. This disorder induced critical behavior at the band center is caused by the sublattice symmetry. After taking the continuum limit, the HWK model reduces to two flavors of Dirac particles subjected to an imaginary Abelian random vector potential and a complex valued random mass.

It is known that monotonicity of the multifractal spectrum for the zero-modes of one flavor of $2D$ Dirac particles subjected to a white-noise random vector potential is implemented in a rather dramatic fashion, namely by the termination of the multifractal spectrum.[4] In analogy to thermodynamics, one can recast the termination of the multifractal spectrum as a freezing transition.[5] From a field theoretical perspective, the termination of the multifractal spectrum in this model is closely related to the existence of an infinite set of operators with negative scaling dimensions whose renormalization group (RG) flows are governed by a non-linear diffusion equation, the Kolmogorov-Petrovsky-Piscounov (KPP) equation.[6,7] In this article, we sketch how, within field theory, the global (i.e., self-averaging) density of states (GDOS) in the HWK model is governed by the *same* KPP equation and diverges at the band center $\varepsilon = 0$ as $|\varepsilon|^{-1}\exp\left(-c|\ln\varepsilon|^{\kappa}\right)$ with $\kappa = 2/3$ in agreement with a prediction by Motrunich *et al.*[8] and in disagreement with an older prediction by Gade[9] ($\kappa = 1/2$).

We start from the continuum limit of the HWK model. The Hamiltonian is given by (σ's are the Pauli matrices and the unit 2×2 matrix)

$$H_{\text{HWK}} = \begin{pmatrix} 0 & D_{\text{HWK}} \\ D_{\text{HWK}}^{\dagger} & 0 \end{pmatrix},$$
$$D_{\text{HWK}} = i\sigma_\mu\partial_\mu + i\sigma_\mu A_\mu + i\sigma_0 A_0 + \sigma_3 A_3, \qquad (1)$$

where $\boldsymbol{A} = (A_1, A_2) \in \mathbb{R}^2$ is the *purely imaginary* random vector potential, $A_0 \in \mathbb{R}$ the *purely imaginary* random scalar potential (chemical potential), and $A_3 \in \mathbb{R}$ the random mass potential. We assume that all potentials are white-noise Gaussian distributed with vanishing means and with the variances specified by $\overline{A_\mu(x)A_\nu(y)} = g_A\delta_{\mu\nu}\delta(x-y)$, for $\mu,\nu = 1,2$, and $\overline{A_\mu(x)A_\nu(y)} = g_M\delta_{\mu\nu}\delta(x-y)$, for $\mu,\nu = 0,3$. Disorder averaging is denoted by an overline. We employ the supersymmetric

(SUSY) method and introduce four-component Grassmann fields $\bar{\psi}$, ψ, and their bosonic partners $\bar{\beta}$, β to represent the resolvent $(H_{\rm HWK} - \varepsilon + i\eta)^{-1}$ of the HWK Hamiltonian in a given realization of the disorder. The advantage of the SUSY representation is that disorder averaging over the resolvent reduces to a Gaussian integration and induces an effective interacting theory with action $S_{\varepsilon - i\eta}$ for the fields $\bar{\psi}$, ψ, $\bar{\beta}$, and β.

Before characterizing the statistical distribution of the DOS, we need to understand the action $S_{\varepsilon - i\eta = 0}$. When the random mass perturbation is switched off, $g_M = 0$, and at the band center, $\varepsilon - i\eta = 0$, the resulting effective interacting field theory S_* defines a conformal field theory (CFT) with $gl(2|2)$ current algebra symmetry for any value of g_A.[10] It is also known that $g_M > 0$ is exactly marginal whereas g_A is marginally relevant with the non-perturbative beta function $\beta_{g_A} = (g_M \zeta)^2/(2\pi^2)$, $\zeta = (1 + g_M/\pi)^{-1}$.[10] More precisely, switching on a finite $g_M > 0$ demotes the CFT to the status of a nearly conformal invariant theory (NCFT).[10] Although conformal invariance is now broken, correlators for the currents generating the $gl(2|2)$ algebra can nevertheless be computed non-perturbatively in g_A and g_M. Furthermore, a weak form of Wick theorem still holds in the NCFT. This is all we will need to compute the GDOS.

To calculate the energy dependence of the GDOS, which we assume to be identical to that of the disorder-average local density of states (LDOS) $\nu_{\rm av}(\varepsilon)$, one must reinstate a finite complex energy, i.e., a $GL(2|2)$ symmetry breaking perturbation $\mathcal{A}_{N=1} = \bar{\psi}\psi + \bar{\beta}\beta$ is added to the action $S_{\varepsilon - i\eta = 0}$ with the complex energy $\varepsilon - i\eta$ as a coupling constant. The RG flow of the expectation value for $\psi(x)\bar{\psi}(x) + \beta(x)\bar{\beta}(x)$ in the critical theory perturbed by a finite but small energy under an infinitesimal rescaling $\mathfrak{a} \to \mathfrak{a} e^{dl}$ of the ultra-violet cutoff $\mathfrak{a}$ gives the Callan-Symanzik equation obeyed by $\nu_{\rm av}(\varepsilon)$.

The energy perturbation $\mathcal{A}_1$ is one member of an infinite set of operators with negative anomalous scaling dimensions.[11] Other members, $\mathcal{A}_N = \frac{1}{N!}(A_1)^N + \cdots$, from this set are generated through the operator product expansion (OPE)[11] $\mathcal{A}_N \times \mathcal{A}_{N'} = \begin{pmatrix} N+N' \\ N \end{pmatrix} \mathcal{A}_{N+N'} + \cdots$. The anomalous scaling dimension x_N of $\mathcal{A}_N$ is[11]

$$x_N = \zeta[1 + \mathcal{O}(g_M)]N - \left[\frac{g_A}{\pi}\zeta^2 + \mathcal{O}(g_M)\right] N^2. \qquad (2)$$

As the relevance of $\mathcal{A}_N$ grows with N or g_A, all $\mathcal{A}_N$ must be taken into account on an equal footing when performing a RG analysis of $\nu_{\rm av}(\varepsilon)$.

The perturbed action needed to perform a *consistent* RG analysis of $\nu_{\rm av}(\varepsilon)$ is obtained by adding to $S_{\varepsilon - i\eta = 0}$ the perturbations $-\sum_{N=1}^{\infty} Y_N \mathfrak{a}^{x_N - 2} \int d^2 r\, \mathcal{A}_N(r)$. An expansion of the one-point correlation function $\nu_{\rm av}(\varepsilon)$ up to second order in the "fugacities" Y_N shows that it obeys a Callan-Symanzik equation characterized by infinitely many one-loop β-functions $\beta_{Y_N} \equiv dY_N/dl$,

$$\beta_{Y_N} = (2 - x_N) Y_N + \pi \sum_{N'=1}^{N-1} \begin{pmatrix} N \\ N' \end{pmatrix} Y_{N'} Y_{N - N'} \qquad (3)$$

with the initial condition $Y_N(l = 0) = (\varepsilon - i\eta)/\mathfrak{a}^{x_1 - 2}$, together with an anomalous scaling *operator* that couples the expectation value of $\psi(x)\bar{\psi}(x) + \beta(x)\bar{\beta}(x)$ to *all* higher powers thereof. This Callan-Symanzik equation is thus an infinite set of coupled differential equations.

To treat all $\mathcal{A}_N$ on equal footing, we define the generating function

$$\widetilde{G}(y, l) = 1 + \frac{\pi}{2} \sum_{n=1}^{\infty} \frac{(-1)^n e^{-ny}}{n!} Y_n(l). \qquad (4)$$

The RG equation for this generating function reduces to a non-linear diffusion equation of the KPP-type,[7]

$$\partial_l \widetilde{G} = \left(\zeta \partial_y + \frac{g_A \zeta^2}{\pi} \partial_y^2\right) \widetilde{G} + 2\widetilde{G}(\widetilde{G} - 1) \qquad (5)$$

with the initial condition $\widetilde{G}(y, 0) = \exp\left(-\frac{\pi}{2} \frac{\varepsilon - i\eta}{\mathfrak{a}^{x_1 - 2}} e^{-y}\right)$. A detailed study of the asymptotic ($l \to \infty$) solution of Eq. (5) suggests that the GDOS $\nu_{\rm av}(\varepsilon)$ is governed by the "*equivalent*" Callan-Symanzik equation[11]

$$0 = \left[z_A \varepsilon \frac{\partial}{\partial \varepsilon} + \beta_{g_A} \frac{\partial}{\partial g_A} - (2 - z_A)\right] \nu_{\rm av}(\varepsilon). \qquad (6)$$

The infinite set of coupling constants Y_N together with the infinite dimensional anomalous scaling operator have been replaced by a *single* coupling constant z_A, the dynamical scaling exponent, that relates a change of the *typical energy scale* to a rescaling of length. From the perspective of Eq. (5), z_A is *uniquely* defined as the velocity of the asymptotic wave front solution. It scales as g_A for $g_A \zeta^2 < 2\pi$ whereas it scales as $\sqrt{g_A}$ for $g_A \zeta^2 \geq 2\pi$. A freezing transition must necessarily take place at $g_A \zeta^2 = 2\pi$ as g_A is marginally relevant. Equation (6) can be solved. The most interesting case is that of weak disorder $g_A(l = 0)\zeta^2 \ll 1$ at the bare level. Sufficiently far away from the band center, i.e., when the sublattice symmetry is completely broken by the perturbing bare energy, the GDOS is flat. Upon approaching the band center, the sublattice symmetry manifests itself by a power-law behavior of the GDOS when the marginal relevance of the Abelian random vector potential can be neglected. At lower energies the relevance of $g_A(l)\zeta^2 \sim 1 < 2\pi$ cannot be neglected anymore and the GDOS starts to increase in a Gade-like manner. Arbitrary close to the band center the relevance of $g_A(l)\zeta^2 \geq 2\pi$ dominates everything else in that the random vector potential induces a strongly fluctuating LDOS (in a log-normal manner). In the strong disorder regime $g_A(l)\zeta^2 \geq 2\pi$, z_A has undergone the freezing transition that is responsible for zero-modes displaying localized behavior instead of multifractality[4] and the GDOS diverging with the exponent $\kappa = 2/3$ instead of $\kappa = 1/2$[8].

In summary, the HWK model is described in the continuum limit by a NCFT in which we identified infinitely many operators with negative anomalous scaling dimensions. These operators reflect the broad distribution of the LDOS and enter in the field theoretical computation of the GDOS through a Callan-Symanzik equation made up of infinitely many differential equations. From a detailed study of the RG equation obeyed by a generating function for an infinite subset of relevant fugacities we identified a scale dependent dynamical exponent that quantifies the change in the typical energy scale due to a rescaling of length. We then conjectured that the GDOS obeys an equivalent Callan-Symanzik equation made of a finite number of differential equation and whose solution agrees with the prediction from Motrunich *et al.* for the GDOS in the HWK model.

[1] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, Phys. Rev. B **50**, 7526 (1994).

[2] Y. Hatsugai, X.-G. Wen, and M. Kohmoto, Phys. Rev. B **56**, 1061 (1997).

[3] S. Ryu and Y. Hatsugai, Phys. Rev. B **65**, 033301 (2002).

[4] C. C. Chamon, C. Mudry, and X.-G. Wen, Phys. Rev. Lett. **77**, 4194 (1996).

[5] H. E. Castillo, C. de C. Chamon, E. Fradkin, P. M. Goldbart, and C. Mudry, Phys. Rev. B **56**, 10668 (1997).

[6] C. Mudry, C. Chamon, and X.-G. Wen, Nucl. Phys. **B466**, 383 (1996).

[7] D. Carpentier and P. Le Doussal, Nucl. Phys. **B588**, 565 (2000). See also B. Horovitz and P. Le Doussal, Phys. Rev. B **65**, 125323 (2002).

[8] O. Motrunich, K. Damle, and D. A. Huse, Phys. Rev. B **65**, 064206 (2002).

[9] R. Gade, Nucl. Phys. B **398**, 499 (1993); R. Gade and F. Wegner, *ibid.* **360**, 213 (1991).

[10] S. Guruswamy, A. LeClair, and A. W. W. Ludwig, Nucl. Phys. **B583**, 475 (2000).

[11] C. Mudry, S. Ryu, and A. Furusaki, Phys. Rev. B **67**, 064202 (2003).