

Phase coexistence and relaxation of the spherical frustrated Blume-Emery-Griffiths model with attractive particles coupling

A. Caiazzo, A. Coniglio, M. Nicodemi

Dipartimento di Scienze Fisiche, INFN, Unità di Napoli,
Monte Sant'Angelo, I-80126 Napoli, Italy

PACS.75.10.Nr { Spin glass and other random models.

PACS.05.50.+q { Lattice theory and statistics.

PACS.64.60.Ht { Dynamic critical phenomena.

Abstract. { We study the equilibrium and dynamical properties of a spherical version of the frustrated Blume-Emery-Griffiths model at mean field level for attractive particle-particle coupling ($K > 0$). Beyond a second order transition line from a paramagnetic to a (replica symmetric) spin glass phase, the density-temperature phase diagram is characterized by a tricritical point from which, interestingly, a first order transition line starts with coexistence of the two phases. In the Langevin dynamics the paramagnetic/spin glass discontinuous transition line is found to be dependent on the initial density; close to this line, on the paramagnetic side, the correlation-response plot displays interrupted aging.

Various mean field spin glass models have been introduced and investigated in the last years; recently, a few models have been introduced [1,4] which combine features of spin glasses and lattice gas, and thus allow to describe important physical quantities, such as density. The frustrated Blume-Emery-Griffiths (BEG) model [3,4], that we discuss here, is a general framework of such kind. Essentially it consists of a lattice gas ($n_i = 0;1$) where the particles interact through their internal degree of freedom ($s_i = \pm 1$) via quenched random couplings and also through a potential depending on a nonrandom coupling constant, K . A complete mean field equilibrium solution of the frustrated BEG has been recently found [4] by numerically solving the equations of the full replica symmetry breaking scheme through a suitable technique of integration. The solution has given evidence of a new interesting behavior in the spin glass panorama, namely a coexistence line between a paramagnetic (P) and a spin glass (SG) phase; the dynamics along this line has not yet been investigated. In this letter we study a spherical version of the mean field frustrated BEG model in the attractive case ($K > 0$) that can be analytically solved in its equilibrium and Langevin dynamics.

The equilibrium phase diagram we get essentially reproduces that of the Ising version ($s_i = \pm 1; n_i = 0;1$) of the model [4]: the plane density-temperature shows a second order transition line from a P to a (replica symmetric) SG phase, ending in a tricritical point; here, a first order transition line starts where coexistence of the two phases exists. The tricritical point lies on the low density branch of the P spinodal ($K < 1$, Fig. 1a), or on the side of the

high density branch ($K > 1$, Fig. 1b); in this last case both a $P=P$ and a $P=SG$ first order line are obtained. In the Langevin dynamics from a random initial condition (rapid quench from high temperature) the equilibrium correlation functions display a diverging relaxation time at the second order line. Furthermore, the $P=SG$ (and, if present, the $P=P$) discontinuous transition line is found to be dependent on the initial density $d(0)$ (Fig. 3); this critical line does not correspond to any singularity of the equilibrium correlation functions, but rather to a diverging plateau displayed by the density function $d(t)$, when its long time limit jumps from an asymptotic (equilibrium) value to the other (Fig. 2). One can recognize the glassy behavior from the two-time quantities which exhibit an aging regime in the whole glassy phase; furthermore, still in the nonglassy phase, close enough to the $P=SG$ discontinuous transition line, one can explicitly observe the pattern of interrupted aging from the correlation-response diagram (Fig. 4).

Our spherical version of the frustrated mean field BEG model is defined as follows. Let us consider the Ising version of the model, described by the Hamiltonian

$$H = \sum_{i < j} J_{ij} s_i n_j s_j n_j - \frac{K}{N} \sum_{i < j} n_i n_j - \sum_i \mu_i n_i \quad (1)$$

being the chemical potential and J_{ij} quenched Gaussian couplings with zero mean and variance $J_{ij}^2 = 1/N$. We introduce two new spin fields $s_i; s_{2i}$ through the change of variables $s_i = s_{1i}, n_i = (s_{1i}s_{2i} + 1)/2$. Then, we assume $s_{1i}; s_{2i}$ as continuous variables, subjected to spherical constraints ($\sum_i s_{ai}^2 = N$ for $a = 1; 2$). Hereafter we shall use the variables s_{ai} to describe the model. The statics can be solved using the eigenvalue density $\rho(\lambda)$ of a large random matrix [5]. The P and the SG phase are described by:

$$\begin{aligned} \frac{z}{2} \frac{P}{z^2 - 4} &= d & d &= 1 - \frac{1}{z + 2 + 2Kd} & q &= 0 & P \\ z = 2 & & d &= 1 - \frac{1}{2 + 2 + 2Kd} & q &= d - \frac{1}{2} & SG \end{aligned} \quad (2)$$

where $d = [\sum_i (s_{1i}s_{2i} + 1)]/2N$ denotes the density, $q = [\sum_i (s_{1i} + s_{2i})^2]/4N$ the overlap and $z = z_1 + Kd$ with $z_1 = z_2$ the two Lagrange multipliers related to the spherical constraints. The stability analysis yields two critical lines, called spinodal lines, at which the corresponding solution becomes unstable against density fluctuations (and both the compressibility and the specific heat are found to diverge):

$$T = \frac{q}{K + K^2 + (1-d)^2 + d^2} \quad P\text{-spinodal}; \quad T = 2K(1-d)^2 \quad SG\text{-spinodal} \quad (3)$$

A second order transition line is located by $d = T$ in the plane $d-T$ (or by $T = T_c(K)$ in the plane $T-K$); below this line one finds from the first of Eqs. (2) that the P solution becomes unphysical. By increasing the parameter K the spinodal lines move to higher temperatures, while the second order line is K -independent. For $K < 1$ (Fig. 1a) the second order line meets the spinodal lines in a tricritical point A , with temperature T_A , located on the low density branch of the P spinodal:

$$T_c(K) = \frac{2K(3 - 2 + (2K(3 - 2)^2 + 16K(1 + 1))}{4K}; \quad T_A(K) = \frac{4K + 1}{4K} \frac{P}{8K + 1} \quad (4)$$

Below the point A, the system undergoes a $P=SG$ first order transition where the order parameters q and d discontinuously jump and a latent heat is involved; the critical line is numerically determined by imposing the free energy balance between the two solutions. Between the spinodal lines in the plane $d-T$, no pure phase is achievable, not even as a metastable one, and the system is a mixture of the two phases. In the zero K limit, T_A vanishes and the first order line disappears [7]. For $K > 1$ (Fig. 1b) the second order line intersects the P spinodal in its high density branch; in this case a first order transition from the low to the high density P phase is also exhibited. Analytical expressions of the thermodynamical quantities can be derived in terms of the above solutions, Eqs. (2). The linear susceptibility, the compressibility and the specific heat show a cusp by crossing the second order line [7] and a finite jump at the first order ones. The SG susceptibility is infinite in the whole SG phase, while remains finite in the P phase, diverging when the second order line is approached [7].




Fig. 1 { Equilibrium phase diagrams in the plane T and d for $K = 0.5$ (a) and $K = 3$ (b). One can have a spin glass (SG) phase or a paramagnetic (P) phase (which forks at the point B in a low density P and an high density P^+ component). The transition lines are marked in bold; the spinodal lines, at which the relative phase becomes unstable, are also shown as continuous lines; dotted lines represent unphysical analytical continuations of critical lines. Finally, full circles mark the tricritical point A and the bifurcation point B on the P spinodal line. Notice that the P phase becomes unphysical below the second order line in the plane $d-T$; a $P=SG$ first order transition line starts out from the tricritical point A, located on the P spinodal in the case (a) or on the side of the P^+ spinodal in the case (b).

Now we deal with the dynamics. We assume that the two spin fields s_i evolve via usual Langevin equations; using the mean field approximation we can replace the particle-particle interaction with $K d(t) \sum_i (s_{1i} s_{2i} + 1) = 2$. Thus when projected onto the basis where J_{ij} is diagonal, the Langevin equations become:

$$\frac{d}{dt} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{pmatrix} -4 - z_1(t) \\ -4 - z_2(t) \end{pmatrix} + \begin{pmatrix} 4 + (1 + K d(t)) \\ 4 - z_2(t) \end{pmatrix} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} + \begin{pmatrix} h_1(t) + \eta_1(t) \\ h_2(t) + \eta_2(t) \end{pmatrix} \quad (5)$$

where $z_a(t)$ ($a = 1, 2$) are two Lagrange multipliers enforcing the spherical constraints, $h_a(t)$ two external fields and $\eta_a(t)$ the thermal noises with zero mean and correlations

$h_a(t)h_b(t^0) = 2T_{ab}(t-t^0)$. The initial fields $s_a(0)$ are assumed to be Gaussian variables with zero mean and variance $s_a(0)s_b(0) = 1 + 2(1 - \alpha_{ab})d(0)$ where $d(0)$ is the initial density. We study the correlation functions $C_{1a}(t;t^0) = d(t)h_1(t)s_a(t^0)$ ($a = 1, 2$), the response functions $G_{1a}(t;t^0) = d(t)h_1(t)h_2(t^0) = h_2(t^0)$ and the density $d(t) = (C_{12}(t;t) + 1)/2$. Due to its linearity, the Langevin system (5) can be explicitly solved for given noises, thus the dynamical quantities can be evaluated averaging over the noises and the eigenvalues [7,8]. We find for $t \neq t^0$ and zero external fields:

$$C_{1a}(t;t^0) = \frac{d(t+t^0/2)}{d(t)} \frac{d(t+t^0/2)}{d(t^0)} 2T_{1a} \int_{t^0}^{t+t^0/2} du \frac{I_1(t+t^0/2-u)}{t+t^0/2-u} \frac{I_1(u)}{u} \quad (6)$$

$$G_{1a}(t;t^0) = \frac{d(t)}{d(t^0)} \frac{I_1(t-t^0)}{t-t^0} + \frac{a}{2} \frac{d(t^0)}{d(t)}$$

where $a = (1 - \alpha)^{a+1}$, $I_1(t)$ is the modified Bessel function of order 1 and $d(t) = (t)^{2K} \int_0^t d(u) du + 2t$, while $d(t)$ and $d(t)$ satisfy the following coupled integro-differential equations:

$$\begin{aligned} d(t) d(t) &= d(0) \frac{I_1(2t)}{t} + T \int_0^t du \frac{I_1(2(t-u))}{t-u} d(u) \\ \frac{d}{dt} d(t) &= (1 - d(t)) \frac{d}{dt} \ln d(t) + 2 + 2K d(t) \quad T \end{aligned} \quad (7)$$

with the initial conditions $d(0) = 1$ and arbitrary $d(0)$.

The special case $K = 0$ is analytically tractable by standard Laplace techniques [7] and one can find the full time dependence of d and d , but this is not the case for $K > 0$. However, it can be verified that the long time fixed points of the $d(t)$ -equation are described by the P and SG equilibrium equations (2) with $d(t)$ respectively proportional to $e^{(z_P - 1)t}$ and $e^{2t} = \frac{P}{4} t^3$; furthermore local analysis of these fixed points yields the same stability conditions of the equilibrium theory. Moving from high temperature towards the equilibrium second order line, a diverging equilibration time $\tau_{eq}^{2nd} \propto (T - T_c)^{-2}$ is found [7]. In the (both $P=SG$ and $P=P$) first order region, between the spinodal lines in the plane (T, d) , two distinct asymptotic solutions are possible; to know which of them is established for long times, stability analysis only is not sufficient, but an appropriate matching with the initial conditions is required. This has been achieved by numerically solving Eqs. (7). As an example, the different curves in Fig. 2 are obtained by keeping $d(0); K; T$ fixed and taking several values of d slightly above and below the estimated critical threshold d_c at which the system jumps from the P to the SG solution. The approach to the critical threshold of the density function is accompanied by a diverging plateau ($\tau_{eq}^{1st} \propto \ln(d_c/d)$) at an unstable fixed point given by the further root at the SG of Eqs. (2), before falling in one of the two stable asymptotic values. Fig. 3 shows that at fixed K the whole critical line depends on the value of the initial density $d(0)$. Lower values of $d(0)$ favor the P phase and, for very low values, the SG solution is obtained only when the P solution becomes unstable, i.e. at the P spinodal line; in this last case $\tau_{eq}^{1st} \propto (P - P_{sp})^{-1}$. A such behavior is to be essentially related to the competition of the two phases in the first order region; the same features are found for the $P=P$ discontinuous transition, apart from the power law behavior of $d(t) \propto e^{2t}$ visible in Fig. 2.

The equilibrium dynamics in the P phase is formally equivalent to a trivial schematic formulation of the Mode Coupling Theory (MCT) with two modes $a = 1, 2$ (using the notation

of [9] the MCT kernel is F_1); the two-time quantities (6) are functions of $t - t^0$ and satisfy the fluctuation-dissipation theorem (FDT), decaying to zero with relaxation time τ_{eq}^{2nd} [7]. In the SG phase, due to the power law factor in $\chi(t)$, a nonequilibrium long-time dynamics is found; this is characterized by a regime ($t' - t^0$) where FDT holds, followed by an aging regime ($t > \tau_{eq}^{1st}$) where the correlation C_{1a} is a function of t/t^0 and the integrated response $\chi_{1a}(t; t^0) = \int_{t^0}^t G_{1a}(t; u) du$ plotted as a function of the correlation, $\chi_{1a}(C_{1a})$, is constant [7,8]. We notice that a SG-like nonequilibrium regime is present also in the P phase, close enough to the P=SG discontinuous transition line, for waiting times lower than τ_{eq}^{1st} (Fig. 2). In this case one can well distinguish the following three steps in the relaxation process (see the curves $T_{11}(C_{11})$ in Fig. 4): an initial FDT regime ($\tau_{eq}^{1st} > t' - t^0$) followed by an aging regime ($\tau_{eq}^{1st} > t > t^0$), as for the SG phase; finally an interrupted aging regime ($t > \tau_{eq}^{1st} > t^0$).

In conclusion, we have analyzed the equilibrium and dynamical properties of the spherical frustrated BEG model at mean field level for $K > 0$. The introduction of dilution and attraction in a standard spin glass model generates a coexistence line between the P and SG phase that exhibits new interesting features. Future extensions or improvements of our framework can be realized, such as a version of the model with $p > 2$ -particles interactions [10]; this last case could display interesting connections with recent studies on the MCT for glassy systems with attractive interactions [11].




Fig. 2 { G Graphs of density $d(t)$ and $\chi(t)e^{-2t}$ (in logarithmic scale) when the system is close to the discontinuous P=SG transition. The curves are obtained by numerically solving Eqs. (7) for $d = 10^{-x}$ with $x = 2; 4; 6; 8$ from left to right and $d' = 1/27$; the other parameters are fixed at $K = 1, T = 0.2$ and $d(0) = 1$. Before falling in one of the two stable asymptotic values the density function exhibits a plateau at an unstable intermediate fixed point, whose length diverges as d approaches the critical threshold $d(\tau_{eq}^{1st} \ln(d))$.

Fig. 3 { We show the P=SG discontinuous critical lines for three values of the initial density, $d(0) = 1; 0.75; 0.5$ (squares, circles, triangles), in the region of the plane (μ, T) between the spinodal lines ($K = 1$). The equilibrium first order line is also shown (dashed line). For $d(0)$ lower than 0.4 the critical line coincides with the P spinodal line.

Fig. 4 { We plot T_{11} vs. C_{11} for several values of t^0 (indicated in the figure) when the system is in the P phase but very close to the P=SG discontinuous transition line ($K = 1, T = 0.1, d(0) = 1; \tau' = 1.44$) and $t^0 < \tau_{eq}^{1st} \cdot 100$. One can distinguish in the relaxation process an initial FDT regime ($T_{11} = 1 - C_{11}$), followed by an aging regime, as for the SG phase, where the curves approach the dotted line ($T_{11} = 1 + T/d$ where d is the density at the unstable fixed point); finally an interrupted aging regime, due to the finite equilibration time, in which the curves bend upward.

REFERENCES

- [1] S.K. Ghatak and D. Sherrington, J. Phys. C 10 (1977) 3149; E.J.S. Lage and J.R.L. de Almeida, J. Phys. C 15 (1982) L1187; P.Mottishaw and D. Sherrington, J. Phys. C 18 (1985) 5201; F.A. da Costa, C.S.O. Yokoi and S.R. Salinas, J. Phys. A 27 (1994) 3365;
- [2] M. Nicodemi and A. Coniglio, J. Phys. A 30 (1997) L187; J. Arenzon, M. Nicodemi and M. Sellitto, J. Phys. I France 6 (1996) 1143; Coniglio, A. de Candia, A. Fierro and M. Nicodemi, J. Phys. Condens. Matter 11 (1999) A167; A. de Candia and A. Coniglio, Phys. Rev. E 65 (2001) 16132
- [3] M. Sellitto, M. Nicodemi and J. Arenzon, J. Phys. I France 7 (1997) 945; G.R. Schreiber, Eur. Phys. J. B 9 (1999) 479;
- [4] A. Crisanti and L. Leuzzi, Phys. Rev. Lett. 89 (2002) 237204
- [5] This is equivalent to a replica symmetric solution in the replica framework; the replica free energy has to be maximized with respect to the overlap q and minimized with respect to the density d [6,10].
- [6] J.M. Kosterlitz, D.J. Thouless and R.C. Jones, Phys. Rev. Lett. 36 (1976) 1217
- [7] A. Cialazzo, A. Coniglio and M. Nicodemi, Phys. Rev. E 66 (2002) 46101
- [8] S. Ciuchi and F. De Pasquale, Nucl. Phys. B 300 (FS22) (1988) 31; L.F. Cugliandolo and D.S. Dean, J. Phys. A 28 (1995) 4213
- [9] W. Gotze, in Liquids, freezing and glass transition, Hansen J.P., Levesque D., Zinn-Justin editors (North Holland, Amsterdam, 1991)
- [10] A. Cialazzo, A. Coniglio and M. Nicodemi, in preparation
- [11] K. Dawson, G. Fort, M. Fuchs, W. Gotze, F. Sciortino, M. Sperl, P. Tartaglia, Th. Voigtmann and E. Zaccarelli, Phys. Rev. E 63 (2001) 11401; E. Zaccarelli, G. Fort, K. Dawson, F. Sciortino and P. Tartaglia, Phys. Rev. E 63 (2001) 31501; G. Fort, F. Sciortino, P. Tartaglia, E. Zaccarelli, F. Lo Verso, L. Reatto, K. Dawson and C.N. Likos, cond-mat/0305561