

# Isotropic-nematic transition in hard-rod fluids: relation between continuous and restricted-orientation models

Kostya Shundyak and Rene van Roij  
 Institute for Theoretical Physics, Utrecht University,  
 Leuvenlaan 4, 3584 CE Utrecht, The Netherlands  
 (Dated: April 14, 2024)

We explore models of hard-rod fluids with a finite number of allowed orientations, and construct their bulk phase diagram within Onsager's second virial theory. For a one-component fluid, we show that the discretization of the orientations leads to the existence of an artificial (almost) perfectly aligned nematic phase, which coexists with the (physical) nematic phase if the number of orientations is sufficiently large, or with the isotropic phase if the number of orientations is small. Its appearance correlates with the accuracy of sampling the nematic orientation distribution within its typical opening angle. For a binary mixture this artificial phase also exists, and a much larger number of orientations is required to shift it to such high densities that it does not interfere with the physical part of the phase diagram.

PACS numbers: 61.30.Cz, 64.70.Md, 05.70.Fh

## I. INTRODUCTION

Understanding the phase behavior of colloidal suspensions of rodlike particles requires an accurate description of their microscopic properties. Fluids of hard rods may be considered as the simplest systems on which the models incorporating particle orientational degrees of freedom can be tested [1, 2]. One of the exact theoretical results dates back to Onsager [3] who analyzed the transition from a uniform isotropic phase to an orientationally ordered nematic phase in a fluid of monodisperse hard needles. He realized that the average pairwise (rod-rod) excluded volume is reduced in the nematic phase compared to that in the isotropic phase, and argued that the resulting gain of free volume (and hence translational entropy) compensates the loss of orientation entropy (due to the nematic ordering) at sufficiently high concentrations of rods [3]. Onsager derived a nonlinear integral equation for the orientation distribution function, a key quantity of the theory, which is constant in the isotropic phase and peaked about the director in the nematic phase. He circumvented the problem of explicitly calculating the nematic orientation distribution function (ODF) by adopting a variational Ansatz, which was numerically checked to be rather accurate later [1].

The generalization of the Onsager model to binary mixtures of rods showed the possibility of strong fractionation [4, 5] and even nematic-nematic demixing at sufficiently high density, driven by a competition between orientation entropy and ideal mixing entropy [6, 7, 8]. The functional forms of ODF's in these studies were either variational Gaussian [5, 7], truncated expansions in Legendre polynomials [4], or numerically determined on an angular grid [6] or on a scaled angular grid [8]. In all these cases the focus was on describing the system with a continuum of orientations.

An alternative approach is to study models with a finite number  $N$  of allowed orientations while the positions of the centers of mass of the rods remain continuous. The

first such model was proposed by Zwanzig [9], with orientations of a rod to be restricted to  $N = 3$  mutually perpendicular directions  $\{\hat{i}; i = 1; \dots; N\}$ . Despite its inability to resolve the orientational structure of the one-particle distribution function in any detail, it has been successfully applied to explore wetting phenomena near a single hard wall and in a slit [10], or phase diagrams of polydisperse systems [11, 12]. The main advantage of such discrete models in comparison with the continuous ones is their computational simplicity. The combination of spatial inhomogeneity and/or polydispersity with a continuum of orientations is rather demanding numerically, and the computational errors can be reduced significantly by discretizing the orientations [11, 13, 14]. The hope has, of course, always been that with an increase of the number of allowed orientations one would smoothly approach the continuum limit. Here we show that this is not the case.

The possibility of a continuous interpolation between results for the discrete models on the one hand and Onsager-like solutions on the other has first been questioned by Straley [15] in studies of models with dodecahedral ( $N = 6$ ) and icosahedral ( $N = 10$ ) symmetries. He concluded that they do not trend towards the continuum solution due to the single allowed orientation within the typical opening angle ( $\approx 9^\circ$ ) of the nematic distribution at coexistence. Unfortunately, one cannot proceed the sequence of models with  $N = 3; 6; 10$  any further, since a larger fully symmetric set of orientations on the unit sphere does not exist. In order to be able to study the effect of discretizing the allowed orientations we give up part of the symmetry of the set, and this allows us to connect continuous and discrete models. We apply our method not only to a one-component system of rods, but also to binary mixtures, which may be considered as the simplest polydisperse systems.

This paper is organized as follows. In Sec. II we derive the grand potential functional for a model with discrete number of allowed rod orientations from the On-

sager functional. We calculate bulk equations of state for specific orientational sets, and determine the number of orientations required to resemble the continuous Onsager solution. In Sec. III we apply the method to construct bulk phase diagrams of binary mixtures of thin and thick rods. We demonstrate that their structure can be significantly modified due to orientational discretization. A summary and some discussion of our results will be presented in Sec. IV.

## II. MONODISPERSE RODS

Consider a fluid of hard rods of length  $L$  and diameter  $D$  ( $L \gg D$ ) in a macroscopic volume  $V$  at temperature  $T$  and chemical potential  $\mu$ . The "Onsager" grand potential functional  $\Omega$  of the one-particle distribution function  $f(\hat{r})$  can be written, within second virial approximation, as [3]

$$\begin{aligned} \Omega[f] = & \int d\hat{r} f(\hat{r}) \ln[f(\hat{r})] - 1 \\ & + \frac{1}{2} \int d\hat{r} d\hat{r}' E(\hat{r}; \hat{r}') f(\hat{r}) f(\hat{r}'), \end{aligned} \quad (1)$$

where  $\beta = (kT)^{-1}$  is the inverse temperature,  $V$  is the rod's thermal volume, and  $E(\hat{r}; \hat{r}')$  is the excluded volume of rods with orientations  $\hat{r}$  and  $\hat{r}'$ . The function  $f(\hat{r})$  is normalized as  $n = \int d\hat{r} f(\hat{r})$ , with  $n$  the bulk number density (which depends on  $\mu$ ). The minimum condition  $\delta \Omega[f] = 0$  on the functional leads to the nonlinear integrodifferential equation

$$\ln[f(\hat{r})] + \int d\hat{r}' E(\hat{r}; \hat{r}') f(\hat{r}') = 0; \quad (2)$$

to be solved for the equilibrium distribution  $f(\hat{r})$ .

Models with a discrete number  $N$  of allowed rod orientations can be systematically derived from the continuous model (1) by dividing the unit sphere into solid sectors  $\hat{r}_i$  ( $i = 1; \dots; N$ ) around vectors  $\hat{r}_i$ , and fixing the rod density  $f(\hat{r}) = f_i$  within each sector as well as the excluded volume  $E(\hat{r}; \hat{r}') = E(\hat{r}_i; \hat{r}_j)$  for every pair of sectors. The grand potential functional  $\Omega_i$  of such "orientationally discretized" fluid with the density  $n_i = f_i$  and the excluded volumes  $E_{ij} = E(\hat{r}_i; \hat{r}_j)$  is

$$\begin{aligned} \Omega_i = & \sum_{i=1}^N f_i \ln[f_i] - 1 \\ & + \frac{1}{2} \sum_{i,j=1}^N E_{ij} f_i f_j \end{aligned} \quad (3)$$

with normalization  $n = \sum_{i=1}^N f_i$ . The last term in Eq. (3) represents the contribution due to the discretization procedure into the grand potential, i.e. the intrinsic difference between continuous and discrete models.

For a homogeneous distribution of vectors  $\hat{r}_i$  on the unit sphere and  $\hat{r}_i = \hat{r}$  (i.e. for the models with  $N = 3; 6$  and  $10$  [15]), it trivially shifts the chemical potential  $\mu = \mu_0 + \ln f$ , which does not have any consequence for the solutions  $f_i$  at a fixed  $n$ , and for the thermodynamics of the isotropic-nematic transition. However, when  $\hat{r}_i$  is not the same for all  $i$ , it acts as an external orientational field that tends to favor the larger sectors over the smaller ones. This becomes explicit if we consider the Euler-Lagrange equations that corresponds to the discrete functional (or equivalently the analog of Eqs. (2))

$$\ln[f_i] + \sum_{j=1}^N E_{ij} f_j = \mu + \ln f_i; \quad (4)$$

now to be solved for  $f_i$ . Note that the equation of state  $p = p(n; T)$  does not pick up an additional term from discretization,

$$p = n + \frac{1}{2} \sum_{i,j=1}^N E_{ij} f_i f_j; \quad (5)$$

but the distributions  $f_i$  to be inserted into it do depend on the discretization.

Further discussion requires a specification of the set of allowed orientations  $\hat{r}_i$ ;  $i = 1; \dots; N$  and the associated solid sectors  $\hat{r}_i$ . Unfortunately, it is impossible to completely cover a surface of the unit sphere by equal regular spherical  $M$ -polygons, where  $M$  indicates the number of polygon's sides (only 5 Platonic solids exist). But symmetries of the function  $f(\hat{r})$  can be explicitly included into the set of vectors  $\hat{r}_i$  in order to simplify the problem. For the present study we fix the  $z$  axis of the coordinate system to be parallel to the nematic director  $\hat{n}$  and assume uniaxial symmetry of the function  $f(\hat{r}) = f(\theta)$ , with  $\theta = \arccos(\hat{r} \cdot \hat{n})$  the angle between  $\hat{r}$  and  $\hat{n}$ . The azimuthal angle is denoted by  $\phi$ , and hence we characterize a vector  $\hat{r} = (\sin \theta \cos \phi; \sin \theta \sin \phi; \cos \theta)$  by the angles  $\theta$  and  $\phi$ . The "up-down" symmetry of the nematic phase reduces the orientational space to half the upper-hemisphere, i.e.  $\theta \in [0; \pi/2]$  and  $\phi \in [0; 2\pi]$ . As we do not expect any azimuthal symmetry breaking we restrict attention to  $N$  uniformly distributed values for  $\theta$  for every allowed  $\phi$ . We have considered a uniform distribution of  $N$  polar angles  $\theta \in [0; \pi/2]$ , i.e.

$$\begin{aligned} (\theta_k; \phi_l) = & \left( \frac{(k-1)\pi}{2(N-1)}; \frac{(l-1)2\pi}{N-1} \right); \\ k = 1; \dots; N; \quad l = 1; \dots; N; \end{aligned} \quad (6)$$

as well as a uniform distribution of  $N$  values of  $\cos \theta \in [0; 1]$ , i.e.

$$\begin{aligned} (\theta_k; \phi_l) = & \left( \arccos \left( 1 - \frac{k-1}{N-1} \right); \frac{(l-1)2\pi}{N-1} \right); \\ k = 1; \dots; N; \quad l = 1; \dots; N \end{aligned} \quad (7)$$

with the conventional definition of  $\hat{f}_i = \frac{R}{i} \sin \theta_i$ .

Figure 1 shows the dimensionless pressure  $p = pL^2D$  as a function of the dimensionless bulk density  $n = nL^2D$  for the grid (6) with different  $N$  and  $N = 5$ . The plateaux (of the solid lines) correspond to the isotropic-nematic coexistence, obtained by equating pressure and chemical potential in the two phases. For  $N = 9$  the transition occurs between the isotropic phase (I) and an (almost) perfectly aligned nematic phase (A) with the pressure being close to the ideal gas pressure. Note that such grids correspond to a single "within the Onsager" opening angle, i.e.  $0 < \theta_1 = 18 < \theta_2$ . As soon as  $0 < \theta_1 < \theta_2 = 18$ , or  $N > 9$ , the distribution function  $(\hat{f}_i)$  at isotropic-nematic coexistence starts to converge to the continuous solution. These results are in full agreement with the previous explanation of Straley [15]. Equations of state for the model (7) are very similar to Fig. 1 but start to resemble Onsager-like distribution function for  $N > 80$  due to the poor sampling of  $\hat{f}_i$  near the nematic director. Equations of state for the Zwanzig ( $N = 3$ ), dodecahedral ( $N = 6$ ) and icosahedral ( $N = 10$ ) models were calculated using the original formulations [15], and are included for comparison. Our results for  $N = 9$  seem to converge well to these existing results.

For  $N > 20$  the pressure of the high-density nematic phase clearly demonstrates a linear dependence on the bulk density, i.e.  $p(n) \propto n$ . With increasing  $N$  it gradually approaches a limiting scaling behavior  $p(n) = 3n$ , established for the continuous Onsager solution by means of a scaling argument [16].

The discretization of the rod's allowed orientations shows the existence of an "artificial transition" from a less-ordered nematic phase (N) to a near-perfectly aligned phase (A) for  $N > 9$ , as indicated by the dashed horizontal lines in Fig. 1. It occurs due to the same competition between excluded volume and orientational entropy as in the IN transition, and puts an additional constraint on the description of the nematic bulk state by restricted-orientation models. Below we argue that it has important consequences for discrete models of (polydisperse) hard-rod mixtures, where separation into nematic phases with different composition occurs at sufficiently high densities.

### III. BINARY MIXTURES

Consider a binary mixture of thin ( $D_1$ ) and thick ( $D_2$ ) hard rods of equal length  $L$  and the diameter ratio  $d = D_2/D_1 = 4$  in a macroscopic volume  $V$  at temperature  $T$  and chemical potentials  $\mu_1$  and  $\mu_2$ , respectively. The "Onsager" grand potential functional for this system can be written as [1]

$$[\hat{f}(\hat{f})g] = \sum_{i=1}^N \frac{X_i^2}{2} \int d\hat{f}(\hat{f}) \ln[\hat{f}(\hat{f})] - 1$$

$$+ \frac{1}{2} \sum_{i,j=1}^N X_i^2 X_j^2 \int d\hat{f}(\hat{f}) d\hat{f}'(\hat{f}') E_{ij}(\hat{f};\hat{f}') - \sum_{i=1}^N \hat{f}_i \ln[\hat{f}_i] - 1 \quad (8)$$

with normalization  $n = \sum_{i=1}^N \hat{f}_i$ . It is known from previous work [17, 18, 19] that the bulk phase diagram of this system exhibits (i) strong fractionation at isotropic-nematic (I-N<sub>2</sub>) coexistence, (ii) nematic-nematic (N<sub>1</sub>-N<sub>2</sub>) coexistence ending in a consolute N<sub>1</sub>N<sub>2</sub> point at sufficiently high pressure, and (iii) an IN<sub>1</sub>N<sub>2</sub> triple point. The discrete version of this model follows directly from (3) as

$$[\hat{f}(\hat{f})g] = \sum_{i=1}^N \frac{X_i^2}{2} \int d\hat{f}(\hat{f}) \ln[\hat{f}(\hat{f})] - 1 + \sum_{i,j=1}^N \frac{X_i^2 X_j^2}{2} \int d\hat{f}(\hat{f}) d\hat{f}'(\hat{f}') E_{ij}(\hat{f};\hat{f}') - \sum_{i=1}^N \hat{f}_i \ln[\hat{f}_i] \quad (9)$$

with the densities  $\hat{f}_i = (\hat{f}_i) \hat{f}_i$ , the excluded volumes  $E_{ij}(\hat{f};\hat{f}') = E_{ij}(\hat{f}_i;\hat{f}_j)$  and the number densities normalization  $n = \sum_{i=1}^N \hat{f}_i$ . Figure 2 shows the phase diagrams for discrete systems in the  $p$ - $x$  representation with the dimensionless pressure  $p = pL^2D_1$  and the mole fraction of thick rods  $x = n_2/(n_1 + n_2)$ , for several orientational grids (6) with  $N = 11$  (a), 20 (b), 30 (c), 50 (d) and  $N = 10$ . Note that all four discretizations are such that they reproduce the physical "Onsager"-like I-N transition at  $x = 0$  and  $x = 1$ , at pressures  $p = 17.7$  and  $4.4$ , respectively. However, the existence of the artificial aligned nematic phase A gives rise to spurious I-A, N<sub>1</sub>-A and N<sub>2</sub>-A phase equilibria, where I, N<sub>1</sub> and N<sub>2</sub> are the physical isotropic and nematic phases. (For the coarsest discretization with  $N = 11$  (Fig. 2(a)) the N<sub>1</sub> phase is stable in a very narrow region beyond the resolution of the picture.) Upon refining the discretization from  $N = 11$  the IN<sub>2</sub>A triple point ( $r$ ) shifts to higher pressures, and combines with the IN<sub>1</sub>A triple point ( $4$ ) at  $N = 30$  (Fig. 2(c)) to form the physical IN<sub>1</sub>N<sub>2</sub> triple point (now denoted by  $r$ ) and an artificial N<sub>1</sub>N<sub>2</sub>A triple point ( $4$ ) at slightly higher pressure. Further grid refinement to  $N = 50$  yields the physical phase diagram with an IN<sub>1</sub>N<sub>2</sub> triple point and N<sub>1</sub>N<sub>2</sub> consolute point as in Ref. [17], but with a spurious N-A coexistence at pressures beyond the "physical" part of the phase diagram. At these pressures one does not distinguish N<sub>1</sub> and N<sub>2</sub> nematic phases, but the single nematic phase denoted by N here.

### IV. SUMMARY

We have explored the connections between continuous and restricted orientation models of monodisperse and binary hard-rod fluids (in the Onsager "needle" limit  $L \gg D$ ). Our main finding is that a discretization of the orientations leads to existence of a nonphysical almost perfectly aligned nematic phase (A) at high densities. If the discretization is coarse, i.e., the number of

allowed orientations is small, then the A phase can co-exist with the isotropic phase (I), and at sufficiently low discretization with the nematic phase N. We also found that the continuum limit requires a finer orientation grid for a mixture than the one-component fluid.

Clearly, our results are strongly influenced by the adopted limit  $L = D$ , but we would expect similar effects, although weaker, for finite  $L=D$  ratio.

These findings could be relevant for the study of inhomogeneous and/or polydisperse fluids of rods, which are computationally more demanding and hence impose the use of rather coarse grid of orientations in order to be tractable and practical. The present study shows that care must be taken with such rather coarse grids, since

they can give rise to an artificial, discretization-induced aligned nematic phase.

#### Acknowledgments

We would like to thank Prof. J.P. Straley for useful correspondence. This work is part of the research program of the "Stichting voor Fundamenteel Onderzoek der Materie (FOM)", which is financially supported by the "Nederlandse organisatie voor Wetenschappelijk Onderzoek (NWO)".

- 
- [1] G.J. Vroege and H.N.W. Lekkerkerker, *Rep. Prog. Phys.* 55, (1992) 1241.
  - [2] S. Fraden, in *Observation, Prediction and Simulation of Phase Transitions in Complex Fluids*, edited by M. Baus et al. (Kluwer, Dordrecht, 1995), p. 113.
  - [3] L. Onsager, *Ann. N.Y. Acad. Sci.* 51, (1949) 627.
  - [4] H.N.W. Lekkerkerker, Ph. Coulon, R. Van Der Haegen, and R. Deblieck, *J. Chem. Phys.* 80, 3427 (1984).
  - [5] Th. Odijk and H.N.W. Lekkerkerker, *J. Phys. Chem.* 89, 1272 (1985).
  - [6] T.M. Birshtein, B.I. Kolegov, and V.A. Pryamitsyn, *Polym. Sci. U.S.S.R.* 30, 316 (1988).
  - [7] G.J. Vroege and H.N.W. Lekkerkerker, *J. Phys. Chem.* 97, 3601 (1993).
  - [8] R. van Roij and B. Mulder, *J. Chem. Phys.* 105, 11237 (1996); 109, 1584 (1998).
  - [9] R. Zwanzig, *J. Chem. Phys.* 39, 1714 (1963).
  - [10] R. van Roij, M. Dijkstra and R. Evans, *Europhys. Lett.* 49, 350 (2000).
  - [11] N. Clarke, J.A. Cuesta, R. Sear, P. Sollich and A. Speranza, *J. Chem. Phys.* 113, 5817 (2000).
  - [12] Y. Martinez-Raton and J.A. Cuesta, *J. Chem. Phys.* 118, 10164 (2003).
  - [13] Z.Y. Chen and J. Noolandi, *Phys. Rev. A* 45, 2389 (1992).
  - [14] K. Shundyak and R. van Roij, *J. Phys.: Condens. Matter* 13, 4789 (2001).
  - [15] J.P. Straley, *J. Chem. Phys.* 57, 3694 (1972).
  - [16] R. van Roij and B. Mulder, *Europhys. Lett.* 34, 201 (1996).
  - [17] R. van Roij, B. Mulder and M. Dijkstra, *Physica A* 261, 374 (1998).
  - [18] K. Shundyak and R. van Roij, *Phys. Rev. Lett.* 88,

205501 (2002).

- [19] K. Shundyak and R. van Roij, *Phys. Rev. E* 68, 0617XX (2003).

#### FIGURE CAPTIONS

1. Equation of state for models with different number of allowed polar angles  $N$ . Positions of the phase transitions are indicated by the horizontal lines. The continuous Onsager solution (Ons) can be reproduced in the present density interval with  $N = 50$ . Dotted lines correspond to equations of state for the Zwanzig (Z), dodecahedral (6S) and icosahedral (10S) models [15]. The dashed horizontal lines correspond to a spurious nematic-nematic ( $N \rightarrow A$ ) transition due to poor discretization of the allowed orientations.
2. Bulk phase diagrams of a binary thin-thick mixture of hard rods (diameter ratio  $D_2/D_1 = 4:0$ , equal length  $L = D_2$ ), in the pressure-composition representation, with  $p$  the dimensionless pressure and  $x$  the mole fraction of the thicker rods. We distinguish the low-pressure isotropic phase (I), high-pressure nematic phases ( $N_1$  and  $N_2$ ), aligned phase A, upper (4) and lower (r) triple phase co-existence and an  $N_1 - N_2$  critical point (\*). The grey regions, enclosed by the binodals, denote the two-phase regimes, and the tie lines that connect coexisting phases, are horizontal.




FIG .1: K .Shundyak and R .van Rooij




FIG . 2: K . Shundyak and R . van Rooij