

Routes to gelation in a clay suspension

B. Ruzicka¹, L. Zulian¹, G. Ruocco^{1,2}¹ INFN and Dipartimento di Fisica, Università di Roma "La Sapienza", P.z.le A. Moro 2, I-00185 Roma, Italy and² INFN-CRSOFT, Università di Roma "La Sapienza", P.z.le A. Moro 2, I-00185, Roma, Italy.

(Dated: April 14, 2004)

The gelation of water suspension of a synthetic clay (Laponite) has been studied by dynamic light scattering in a wide range of clay weight concentration ($C_w = 0.003 - 0.031$). At variance with previous determination, indicating a stable liquid phase for $C_w < C_w^* = 0.015 - 0.018$, we find that the gelation takes actually place in the whole examined C_w range. More importantly, we find that C_w^* marks the transition between two different routes to gelation. We hypothesize that at low concentration Laponite suspension behaves as an attractive colloid and that the slowing down of the dynamics is attained by the formation of larger and larger clusters while at high concentration the basic units of the arrested phase could be the Debye-Hückel spheres associated to single Laponite plates.

PACS numbers: 64.70.Pf, 82.70.Dd, 78.35.+c, 61.20.Lc

Colloidal systems are ideal benchmarks for studying, by optical microscopy and light scattering, equilibrium slow dynamical processes and the formation of non-ergodic arrested states of matter. In this context sterically stabilized colloidal systems have been often used as a model for hard spheres, to study the fluid-crystal transition and the glass formation [1]. For suspensions of hard spheres the latter transition occurs at relatively high packing fraction ($\phi \approx 0.58$). When screened charged interactions are present, crystal and disorder arrested phases can be formed also at very low volume fraction ($\phi \approx 0.01$). Recent works [2, 3, 4, 5] focusing on short-range attractive colloids have attempted a connection between the gel and the glass arrested states of matter. In these systems, a re-entrant glass line, two kinds of glasses (attractive and repulsive) and a glass-glass line have been predicted [2, 3, 6] and experimentally observed [4, 5]. Correlations between the dynamical behavior of gels and glasses suggest that a common understanding of these two disordered forms of matter may emerge. In the case further complicated where short-ranged attractive interactions are complemented by weak repulsive electrostatic interactions, the gel formation process can be fully modeled as a glass transition phenomenon [7]. In the last case, the gel phase is stabilized by the competition of the short range attraction and the long range repulsion. Despite these recent progresses, a deeper comprehension of the still puzzling liquid-gel transition in colloidal systems is requested.

Aqueous suspensions of charged platelike colloids have been the subject of intense experimental and theoretical investigations. The synthetic Hectorite clay Laponite is a perfect model for these charged platelike colloids. It is in fact composed of nearly monodisperse, rigid, disc-shaped platelets with a well-defined thickness of 1 nm, an average diameter of about 30 nm and a negative surface charge of a few hundred e. In spite of intensive research on Laponite suspensions motivated by its important industrial applications, there is no general agreement about

the mechanism of gelation; both attractive and repulsive forces are claimed to be responsible for the gel formation attributed by different authors to Wigner glass transition [8], frustrated nematic transition [9, 10, 11, 12], micro-segregation [9, 13, 14], gelation [15, 16, 17], etc. Hence, studies of the aggregation in Laponite suspensions will give an important contribution to the understanding of the gel formation process in systems where repulsive long range and attractive short range interactions compete.

According to the phase diagram obtained from Mourchid et al. [9] at ionic strength below $I = 10^{-2}$ M Laponite suspensions can be in two different physical states. Low concentration suspensions ($C_w < C_w^*(I)$) form stable, equilibrium fluid phase. Higher concentration suspensions ($C_w \geq C_w^*(I)$) are initially fluids but experience aging and pass into a gel phase after a time that depends on the clay amount. Several dynamical scattering studies have investigated the aging process of samples belonging to this high C_w phase region [8, 18, 19, 20, 21], while only two recent papers [16, 21] report about the lower concentration region. While the first paper [16] reports about the formation of a gel that does not flow when tilted at low Laponite concentration ($C_w = 1\%$, $I = 5 \text{ mM NaCl}$), the authors of the second paper [21] find that samples at low concentrations ($C_w = (0.1 - 1)\%$, $I = 5 - 10^4 \text{ M}$) are stable liquid suspensions. For this reason and to try to understand the gelation mechanism for this charged colloidal system we have performed accurate and systematic measurements on several samples in the low concentration region and at higher clay concentrations.

In this letter we present a dynamic light scattering study at increasing concentrations from 0.3 to 3.1 wt % at ionic strength between $I \approx 10^{-4} \text{ M}$ and $I \approx 10^{-3} \text{ M}$ where $C_w^* = 1.5 - 1.8\%$ [9]. A new gel region, at $C_w < C_w^*$, has been identified and the gelation processes in the two gel phases (at $C_w < C_w^*$ and at $C_w > C_w^*$) have been accurately studied. Two different routes to gelation

for the two regions have been found and a possible explanation for the processes involved in the building of the arrested phases has been proposed.

Laponite is a hectorite synthetic clay manufactured by Laporte Ltd, who kindly supplied us the material. Particular attention has been devoted to samples preparation to avoid dissolution of Laponite platelets that occurs if the samples are exposed to air contamination due to the presence of atmospheric CO_2 [22, 23] and can seriously affect the measurements. For this reason the whole procedure has been performed in a glove box under N_2 flux and the samples have been always kept in safe atmosphere during and after sample preparation. The powder was firstly dried in an oven at $T = 400 \text{ K}$ for 4 hours (up to 20% of the as received powder weight is due to adsorbed water), then the powder is dispersed in deionized ($\text{pH} = 7$) water, stirred vigorously until the suspensions were cleared and then filtered through $0.45 \mu\text{m}$ pore size Millipore filters. For some of the samples the pH has been measured by a Crison Glp 22 pH meter, the final value, reached after 10–20 min from the laponite dispersion in water, was in the range $\text{pH} = 9.8$ – 10.0 . The final value of $\text{pH} = 10$, neglecting the contribution of Na^+ counterions released from some of the platelets to the solution, would fix the ionic strength around 10^{-4} M . Assuming that 50 counterions are released by each laponite platelet [15], the ionic strength never exceeds 10^{-3} M for all the investigated samples. The fact that laponite dissolution does not take place in our samples is confirmed by NMR spectroscopy [24] that checked continuously the amount of Na^+ and Mg^{2+} ions present in the solutions. These measurements show in fact that, both in the low and in the high concentration ranges, the fraction of solvated Na^+ ions does not change with time, and that the amount of solvated Mg^{2+} ions are below the instrumental sensibility. These observations indicate that the chemical reaction of Laponite dissolution [22, 23] is not affecting our samples. Furthermore, in none of the samples studied we observed any evidence of sedimentation also several months after their preparation as has been checked by direct sample inspection and by performing test measurements of the correlation function at different heights in the cells. The starting aging time ($t_w = 0$) is defined as the time when the suspension is filtered. This sample preparation procedure is similar, but not identical, to that already used in previous works [16, 17, 20, 21]. Some authors do not report information on the drying of the powder, others allow the suspension to get in contact with atmosphere, and different details can lead to small differences in the sample behavior on aging. In addition, the determination of the sample concentration is also problematic, as the filtering procedure can alter the actual concentration if large clusters are present in the suspension after the stirring phase. To minimize this effect, we have prepared a large amount of concentrated solution, then –after the filtering– we have diluted this parent sample at the de-

sired concentrations. In this way, in the present measurements, the relative concentrations are well defined, even if the absolute value in C_w can be affected by a systematic error, that we estimate in the range $C_w = C_w \pm 0.1$.

Dynamic light scattering measurements were performed using an ALV-5000 logarithmic correlator in combination with a standard optical set-up based on a He-Ne ($\lambda = 632.8 \text{ nm}$) 10 mW laser and a photomultiplier detector. The intensity correlation function was directly obtained as $g_2(q; t) = \langle I(q; t)I(q; 0) \rangle = \langle I(q; 0) \rangle^2$, where q is the modulus of the scattering wave vector defined as $q = (4\pi n) \sin(\theta/2)$ ($\theta = 90^\circ$ in the present experiment). For the low concentration samples ($C_w = 1.5\%$) the measurements have been repeated once a week for a long period of time (up to four months for the lowest concentration), while for $C_w > 1.5\%$ the sample was left in the scattering system and photocorrelation spectra were continuously acquired until the gelation was reached.




FIG. 1: Evolution of the measured intensity correlation functions (symbols) and corresponding fits with Eq. (1) (continuous lines) for two different Laponite suspensions at the indicated concentrations at different waiting times t_w . The curves are measured at increasing waiting times that (from left to right) are $t_w = 288, 792, 1128, 1464, 1632, 1800, 1968, 2136, 2328, 2640$ hours for sample (A) and $t_w = 6, 30, 54, 78, 102, 126, 150$ hours for sample (B) (for which only data in the ergodic phase are reported).

As an example, correlation functions at increasing aging times t_w for two different samples at low- (Fig. 1A - $C_w = 1.1 \text{ wt } \% < C_w$) and high-concentration (Fig. 1B $C_w = 2.5 \text{ wt } \% > C_w$), are reported in Figure 1. As it is evident from the figure both the samples are doing aging, the dynamics is in fact becoming slow and slow for increasing waiting time t_w . This behavior was expected for the higher concentration sample of Fig. 1B, several measurements [8, 18, 19, 20, 21], indeed, have shown the existence of aging and a sol-gel transition for long enough waiting times. The aging process was instead unexpected for the lower concentration sample of Fig. 1A where the liquid state is predicted from the phase diagram [9], and confirmed by recent dynamic light scattering measurements [21] where, perhaps, the authors did not wait for a time long enough to observe significant aging. From Fig. 1A it is instead evident that the system is aging and for the longest waiting time reported ($t_w = 2640$ hours) there is a qualitative change in the correlation function: a crossover between a complete and an incomplete decay. This behavior is the indication of a strong ergodicity breaking, signature of a sol-gel transition, as already observed in [8, 18] for samples in the higher concentration region and in [16] for a sample at $C_w = 1\%$ with $I = 5 \cdot 10^3 \text{ M}$. Our dynamic light scattering measurements show that for all the samples studied, down to the lowest concentrated one ($C_w = 0.3\%$), there is a typical waiting time, t_w^g , increasing with decreasing clay concentration C_w , at which the system undergoes the gelation. Depending from the initial concentration this gelation time can vary from hours to several months.

Figure 1 shows also that the correlation functions decay following a two steps behavior, i.e. there are two different relaxation processes, a fast and a slow ones. For this reason the fitting expression should contain two contributions. In this case the squared sum of an exponential and a stretched exponential function is used, as already reported in Ref.[20]:

$$g_2(q;t) - 1 = b a e^{-(t/\tau_1)} + (1-a)e^{-(t/\tau_2)^\beta} \quad (1)$$

where b represents the coherence factor. The stretched exponential, instead than another simple exponential term, is used since it has been found to give a good description of the slow relaxation process in glassy systems. The fitting expression well describes the photocorrelation spectra for all the aging times in the liquid (ergodic) phase and for all the investigated concentrations with the relaxation time τ_1 associated to the fast dynamics and the relaxation time τ_2 and the stretching parameter β that describe the slow part of the autocorrelation function. The fits are shown as full lines in Fig. 1 and the resulting chi square is always within its standard deviation.

Another important peculiarity of the correlation func-

tions, that can be directly observed in the raw data reported in Fig. 1, is that the aging process evolves differently for the lower and higher concentration samples. While in fact the initial decay of the correlation functions - fast dynamics, characterized by the correlation time τ_1 - remains constant with waiting time for the higher concentration sample (Fig. 1B) it seems to become slower with increasing waiting time for the lower concentration sample (Fig. 1A). This different behavior is confirmed from the results of the fits, that also show another major difference between the samples at low and high concentrations.

In the following we will report the t_w and C_w dependence of the fitting parameters, in particular we focus on the behavior of the parameters of the slow decay: the relaxation time τ_2 , the stretching exponent β and the "mean" relaxation time τ_m :

$$\tau_m = \frac{1}{2} - \left(\frac{1}{\beta}\right): \quad (2)$$

where Γ is the usual Euler gamma function.




FIG. 2: Waiting time dependence of τ_m (see Eq. 2), i.e. of the average relaxation time of the slow decay of the correlation functions as those reported in Fig. 1. As an example, the t_w dependence of τ_m is reported for the three indicated concentrations. Continuous lines are fits of the data with Eq. 3.

As an example, the values of τ_m obtained for three different concentrations are reported in Fig. 2. It is evident the common behavior of τ_m that seems to diverge at a given $t_w = t_w^g$ i.e. when the gelation occurs. This clearly shows that, as already seen from the raw correlation functions, also the samples at concentrations lower/equal to 1.5 wt % are actually doing aging and undergo a gelation transition. To have more information about this gelation process we represent the aging time (t_w) dependence of mean relaxation time (τ_m) with the law:




FIG. 3: Concentration dependence of the divergence time t_w^1 (A) and of the B parameter (B), entering in Eq. 3. The dashed lines are guides to the eyes. The shadow area indicates the region where the liquid-gel transition was supposed to be according to Ref. [9]

$$m = m_0 \exp \left[B \frac{t_w}{t_w^1} \right] \quad (3)$$

We do not have any rationale behind such a law, it can be considered a generalization to long waiting time of the exponential growth with t_w observed for τ_2 by Abou et al. [20] in the high concentration samples. Equation 3—reported in Fig. 2 as full lines—well describe the measurements. The most significative parameters of the fits, t_w^1 and B , for all the studied concentrations are shown in Fig. 3. Here the vertical dotted region indicates the range of concentrations that, in the phase diagram [9], would mark the transition from the "liquid" to the gel phase in the range of $1 \cdot 10^{-3}$ – 10^{-4} M. The results of the fits indicate that t_w^1 —that can be considered as the time at which the gelation actually occurs—is continuously decreasing with increasing clay concentration, without any evident discontinuity in correspondence of the "transition" region. The concentration dependence of the parameter B , which measures how fast m approaches the divergence, is shown in Fig. 3B. This parameter is almost constant for all the samples in the low concentration region, while shows a large discontinuity on passing in the higher concentration region. It is important to note that

this jump takes place in a region that encompasses the supposed "liquid-gel transition" region.




FIG. 4: The waiting time dependence of the m (A) and (B) are reported as a function of the scaled variable t_w/t_w^1 for some of the investigated concentrations: $\square = 0.3\%$, $\circ = 0.5\%$, $\times = 0.6\%$, $\triangle = 0.8\%$, $\diamond = 0.9\%$, $+$ = 0.9% , $\square = 1.0\%$, $\circ = 1.1\%$, $\times = 1.2\%$, $\triangle = 1.3\%$, $\diamond = 1.4\%$, $\circ = 1.5\%$, $\times = 2.2\%$, $\triangle = 2.5\%$, $\diamond = 2.8\%$. The data for $C_w = 1.5\%$ collapse on a single master curve.

The fact that the value of the B parameter is almost constant for all the low concentration samples is an indication of the existence of a scaling law. Indeed, if we plot the quantities $m(t_w)$ and (t_w) as a function of t_w/t_w^1 all the different C_w data with almost constant B should collapse on a single master curve. The m and the β parameters in function of the normalized waiting time are reported in Fig. 4A and 4B respectively. As expected, all the data for both the m and the β parameters of lower concentrations collapse, within their statistical uncertainties, on a single curve while the data at higher clay concentrations have a different behavior. We already observed from the direct comparison of Fig. 1A with Fig. 1B that the aging process is qualitatively different for the low and high concentration samples. Figures 3 and 4 quantify this difference in the physical properties characterizing the aging phenomenon in the two different concentration regions.

In conclusion the present observations indicate that the stable phase of Laponite suspensions in pure water at $C_w = 1.5\%$ —expected to be liquid according to previ-

ous studies [9]— actually is an arrested phase. Recently Nicolai et al. [15] proposed a revisitation of the phase diagram and suggested that the equilibrium state in the low concentration region is not liquid but a very fragile gel that takes a long time to appear [17]. Probably this "long time" is the reason why previous measurements indicated this phase as liquid. Our measurements clearly show not only that at all the investigated concentrations the stable phase is an arrested phase but also that the aggregation process is basically different for the samples at $C_w = C_w$ and those at $C_w < C_w$. This is evident from the behavior of η and reported in Fig. 4 and from the discontinuity of the B parameter across the $C_w = C_w$ region as shown in Fig. 3. In this sense the "liquid-gel" transition line of the phase diagram [9] is not a real liquid-gel transition but rather seems to indicate a sort of "fragile gel-gel" transition.

The origin of the different routes towards an arrested phase observed for low- and high-concentration Laponite suspensions calls for an explanation that goes beyond the aim of the present work. At the level of speculation, it is worth to recall that the microscopic interaction between Laponite plates is due to a screened Coulomb interaction, which can be modeled by a Yukawa-like repulsion at long distances and a quadrupolar electric and/or a van der Waals terms at short distances [25, 26]. This competition of short-range attraction and long range repulsion resembles that recently proposed [7] to describe the phenomenology of colloidal gels. In Ref. [7] it has been suggested that the gelation taking place in attractive colloidal suspensions at very low concentration involves the growth of larger and larger clusters (driven by the short range attraction). The glass transition of these clusters, driven by the long range repulsion, is responsible for the arrest of the dynamics and the formation of a gel phase. The authors prove that the gel is essentially a Wigner glass, composed by clusters of colloidal particles. We can speculate that low concentrations Laponite suspensions behave as the attractive colloids in Ref. [7]. This scenario is in agreement with the fact that the long time needed to form the arrested phase in the low concentration range is spent by the system to build up the clusters. Also at high Laponite concentration the arrested state could be a Wigner glass but in this case the packing fraction of the

Debye-Hückel sphere associated to each platelets reaches values as high as 0.43 [8] and the single Laponite platelet would be the elementary constituent of the Wigner glass.

-
- [1] P. N. Pusey, and W. van Megen, *Nature* 320, 340 (1986); *Phys. Rev. Lett.* 59, 2083 (1987).
 - [2] J. Bergenholz and M. Fuchs, *Phys. Rev. E* 59, 5706 (1999).
 - [3] L. Fabbian et al., *Phys. Rev. E* 59, R1347 (1999).
 - [4] K. N. Pham et al., *Science* 296, 104 (2002).
 - [5] T. Eckert and E. Bartsch, *Phys. Rev. Lett.* 89, 125701 (2002).
 - [6] E. Zaccarelli et al., *Phys. Rev. E* 66, 41402 (2002).
 - [7] F. Sciortino, S. Mossa, E. Zaccarelli, and P. Tartaglia, preprint cond-mat/0312161.
 - [8] D. Bonn et al., *Europhys. Lett.* 45, 52 (1998).
 - [9] A. M. Ourchid et al., *Langmuir* 11, 1942 (1995).
 - [10] J. C. P. Gabriel, C. Sanchez, and P. Davidson, *J. Phys. Chem.* 100, 11139 (1996).
 - [11] M. Kroon, W. L. Vos, and G. H. Wegdam, *Phys. Rev. E* 57, 1962 (1998).
 - [12] A. M. Ourchid et al., *Langmuir* 14, 4718 (1998).
 - [13] F. Pignon et al., *Phys. Rev. E* 56, 3281 (1997).
 - [14] C. Martin et al., *Phys. Rev. E* 66, 21401 (2002).
 - [15] P. Mongondry, J. F. Tassin, and T. Nicolai, submitted to *J. Colloid Interface Sci.*
 - [16] T. Nicolai and S. Coccard, *J. Colloid Interface Sci.* 244, 51 (2001).
 - [17] T. Nicolai and S. Coccard, *Eur. Phys. J. E* 5, 221 (2001).
 - [18] M. Kroon, G. H. Wegdam and R. Sprik, *Phys. Rev. E* 54, 6541 (1996).
 - [19] A. Knaebe et al., *Europhys. Lett.* 52, 73 (2000).
 - [20] B. Abou, D. Bonn, and J. Meunier, *Phys. Rev. E* 64, 021510 (2001).
 - [21] M. Bellur et al., *Phys. Rev. E* 67, 031405 (2003).
 - [22] D. W. Thompson and J. T. Butterworth, *J. Colloid Interface Sci.* 151, 236 (1992).
 - [23] A. M. Ourchid and P. Levitz, *Phys. Rev. E* 57, R4887 (1998).
 - [24] S. Capuani, T. Gili, and B. Maraviglia, preprint.
 - [25] M. Dijkstra, J. P. Hansen, and P. A. M. Adden, *Phys. Rev. Lett.* 75, 2236 (1995).
 - [26] E. Trizac et al., *J. Phys.: Condens. Matter.* 14, 9339 (2002).