

On the zero-resistance states generated by radiation in GaAs/AlGaAs

S. Fujita* and K. Ito†

University at Buffalo, SUNY, Buffalo, NY 14260, USA

(Dated: March 22, 2022)

The applied radiation excites “holes”. The condensed composite (c)-bosons formed in the excited channel create a superconducting state with an energy gap. The supercondensate suppresses the non-condensed c-bosons at low temperatures, but it cannot suppress the c-fermions in the base channel, and the small normal current accompanied by the Hall field yields a B -linear Hall resistivity.

PACS numbers: 71.10.Pm, 71.15.Qe, 73.43.Cd

Recently Mani *et al.* [1] observed a set of zero-resistance (superconducting) states in GaAs/AlGaAs heterojunction subjected to radiations at the low temperatures (~ 1.5 K) and the magnetic fields (~ 0.2 T). The (diagonal) resistance $R_{xx} \equiv R$ rises exponentially and symmetrically on both sides of the fields centered at $B = 4/5 B_f$, $4/9 B_f$, $B_f = \omega m^*/e$, ω = radiation frequency, m^* = effective mass, e = electron charge, indicating an energy gap ϵ_g in the elementary excitation spectrum. The phenomenon is similar to that observed in the same system in the traditional quantum Hall effect (QHE) regime ($T \sim 0.5$ K, $B \sim 10$ T) [2], with the main difference that the superconducting states are not accompanied by the Hall resistivity plateaus for the system subjected to radiation. We call the temperature below which the superconducting state appears the critical temperature T_c . The T_c (1.5 K) at $B = 4/5 B_f$ is considerably higher than the traditional QHE critical temperature (0.5 K). Zudov *et al.* [3] reported similar magnetotransport properties for the system subjected to radiation with slightly different experimental conditions [3]. They suggested that the principal resistivity minima occur at $B = 2/(2j+1)B_f$, $j = 1, 2, \dots$, rather than $B = 4/(4j+1)B_f$ (Mani’s case). They also noted a noticeable side resistivity minimum besides the principal set of the minima.

In finer experiments Mani *et al.* [4] observed (a) that the deviation in the Hall resistance $\Delta R_H \equiv R_H - R_{H,\text{dark}}$, correlates with the resistance R such that ΔR_H nearly vanishes when $R = 0$, and (b) that ΔR_H is negative, and it is antisymmetric with respect to small B -fields: $\Delta R_H = \alpha B$, α = constant, for small B . The second property means that there is a current due to “hole”-like particles having the charge of the opposite sign to that of the main current carrier charge. We suspect that the applied radiation generates the “holes”. This may be checked by applying the circularly polarized lasers. The small slope $d\Delta R_H/dB = \alpha$ means that the “hole”-like particle density is considerable higher than

the “electron”-like particle density. Mani *et al.* [1] suggested for the cause of the spectral gap the electron pairing due to the excitations induced by radiation. Other mechanisms were proposed by several theoretical groups [5]. But none of them are conclusive. In particular no explanation is given to the question why the superconducting state can occur without the Hall resistivity plateau.

Earlier Fujita *et al.* [6] developed a microscopic theory of the QHE based on the electron(fluxon)-phonon interaction. In this theory the composite (c-) particles (bosons or fermions) having a conduction electron and a number of flux quanta (fluxons) are bound by the phonon-exchange attraction. The composites move as bosons (fermions) depending on the odd (even) number of fluxons in them. At the Landau level (LL) occupation number (filling factor) $\nu = 1/Q$, Q odd, the c-bosons with Q fluxons are generated, and condense below the critical temperature T_c . The Hall resistivity plateau is due to the Meissner effect, see below.

In the present work we shall extend our theory to the excited channel and show that the main series represent the integer QHE with the $\nu = 1$ state corresponding to the superconducting state $B = 4/5 B_f$ in Mani’s case (the state $B = 2/3 B_f$ in Zudov’s case). The state in the base channel at $\nu = 1$ remains fermionic, accounting for the small current and the B -linear behavior in the Hall resistance. If we write $\nu = P/Q$, the Mani-series $4/(4j+1)$ can be recovered by setting $j = P$, $Q = 1$. We predict that the fractional QHE should exist for $Q = 3, 5, \dots$. The aforementioned side dip observed by Zudov *et al.* [3] should correspond to the state at $\nu = 4/3$. This dip is missing in Mani *et al.* experiments. We believe that this is because the experimental temperature $T = 0.5$ is so low that the $\nu = 1$ state overshadowed it. All of the superconducting states $\nu = P/Q$, $P = 2, 3, \dots$, disappear at 0 K.

GaAs forms a zinc blende lattice. We assume that the interface is in the plane (001). The Ga^{3+} ions form a square lattice with the sides directed in $[110]$ and $[\bar{1}\bar{1}0]$. The “electron” (wave packet) will then move isotropically with an effective mass m_1 . The As^{3-} ions also form a square lattice at a different height in $[001]$. The “holes”, each having a positive charge, will move similarly with an effective mass m_2 . A longitudinal phonon moving in $[110]$ or in $[\bar{1}\bar{1}0]$ can generate a charge (current) density variations, establishing an interaction between the phonon and

*Electronic address: Fujita@buffalo.edu

†Electronic address: Keiito@buffalo.edu; Also at Research Division, National Center for University Entrance Examinations, 2-19-23 Komaba, Meguro-ku, Tokyo 153-8501, Japan; Electronic address: ito@rd.dnc.ac.jp

the electron (phonon). If one phonon exchange is considered between the electron and the fluxon, a second-order perturbation calculation establishes an effective electron-fluxon interaction

$$|V_q V'_q| \frac{\hbar \omega_q}{(\epsilon_{|\mathbf{p}+\mathbf{q}|} - \epsilon_p)^2 - (\hbar \omega_q)^2}, \quad (1)$$

where \mathbf{q} ($\hbar \omega_q$) is the phonon momentum (energy), V_q (V'_q) the interaction strength between the electron (fluxon) and the phonon. If the energies ($\epsilon_{\mathbf{p}+\mathbf{q}}$, ϵ_p) of the final and initial electron states are equal, the effective interaction is attractive.

Following Bardeen-Cooper-Schrieffer (BCS) [7], we start with a Hamiltonian H with the phonon variables eliminated:

$$H = \sum_{\mathbf{k}} \sum_s \epsilon_k^{(1)} n_{\mathbf{k}s}^{(1)} + \sum_{\mathbf{k}} \sum_s \epsilon_k^{(2)} n_{\mathbf{k}s}^{(2)} + \sum_{\mathbf{k}} \sum_s \epsilon_k^{(3)} n_{\mathbf{k}s}^{(3)} - v_0 \sum_{\mathbf{q}} \sum'_{\mathbf{k}} \sum'_{\mathbf{k}'} \sum_s \left[B_{\mathbf{k}'\mathbf{q}s}^{(1)\dagger} B_{\mathbf{k}\mathbf{q}s}^{(1)} + B_{\mathbf{k}'\mathbf{q}s}^{(1)\dagger} B_{\mathbf{k}\mathbf{q}s}^{(2)\dagger} + B_{\mathbf{k}'\mathbf{q}s}^{(2)} B_{\mathbf{k}\mathbf{q}s}^{(1)} + B_{\mathbf{k}'\mathbf{q}s}^{(2)} B_{\mathbf{k}\mathbf{q}s}^{(2)\dagger} \right], \quad (2)$$

where $n_{\mathbf{k}s}^{(j)}$ is the number operator for the “electron” (1) [“hole” (2), fluxon (3)] at momentum \mathbf{k} and spin s with the energy $\epsilon_{\mathbf{k}s}^{(j)}$. We represent the “electron” (“hole”) number $n_{\mathbf{k}s}^{(j)}$ by $c_{\mathbf{k}s}^{(j)\dagger} c_{\mathbf{k}s}^{(j)}$, where c (c^\dagger) are annihilation (creation) operators satisfying the Fermi anticommutation rules: $\{c_{\mathbf{k}s}^{(i)}, c_{\mathbf{k}'s'}^{(j)\dagger}\} \equiv c_{\mathbf{k}s}^{(i)} c_{\mathbf{k}'s'}^{(j)\dagger} + c_{\mathbf{k}'s'}^{(j)\dagger} c_{\mathbf{k}s}^{(i)} = \delta_{\mathbf{k},\mathbf{k}'} \delta_{s,s'} \delta_{i,j}$, $\{c_{\mathbf{k}s}^{(i)}, c_{\mathbf{k}'s'}^{(j)}\} = 0$. We represent the fluxon number $n_{\mathbf{k}s}^{(3)}$ by $a_{\mathbf{k}s}^\dagger a_{\mathbf{k}s}$, with $a(a^\dagger)$, satisfying the anticommutation rules. $B_{\mathbf{k}\mathbf{q}s}^{(1)\dagger} \equiv c_{\mathbf{k}+\mathbf{q}/2,s}^{(1)\dagger} a_{-\mathbf{k}+\mathbf{q}/2,-s}^\dagger$, $B_{\mathbf{k}\mathbf{q}s}^{(2)} \equiv c_{\mathbf{k}+\mathbf{q}/2,s}^{(2)} a_{-\mathbf{k}+\mathbf{q}/2,-s}$. The prime on the summation means the restriction: $0 < \epsilon_{\mathbf{k}s}^{(j)} < \hbar \omega_D$, ω_D = Debye frequency. If the fluxons are replaced by the conduction electrons (“electrons”, “holes”) our Hamiltonian H is reduced to the original BCS Hamiltonian, Eq. (24) of ref. 7. The “electron” and “hole” are generated, depending on the energy contour curvature sign [8]. For example only “electrons” (“holes”), are generated for a circular Fermi surface with the negative (positive) curvature whose inside (outside) is filled with electrons. Since the phonon has no charge, the phonon exchange cannot change the net charge. The pairing interaction terms in Eq. (2) conserve the charge. The term $-v_0 B_{\mathbf{k}'\mathbf{q}s}^{(1)\dagger} B_{\mathbf{k}\mathbf{q}s}^{(1)}$, where $v_0 \equiv |V_q V'_q| (\hbar \omega_0 A)^{-1}$, A = sample area, is the pairing strength, generates the transition in the “electron” states. Similarly, the exchange of a phonon generates a transition in the “hole” states, represented by $-v_0 B_{\mathbf{k}'\mathbf{q}s}^{(2)} B_{\mathbf{k}\mathbf{q}s}^{(2)\dagger}$. The phonon exchange can also pair-create and pair-annihilate “electron” (“hole”)-fluxon composites, represented by $-v_0 B_{\mathbf{k}'\mathbf{q}s}^{(1)\dagger} B_{\mathbf{k}\mathbf{q}s}^{(2)\dagger}$, $-v_0 B_{\mathbf{k}'\mathbf{q}s}^{(2)} B_{\mathbf{k}\mathbf{q}s}^{(1)}$. At 0 K the system can have equal numbers of $-(+)$ c-bosons, “electrons” (“holes”) composites, generated by $-v_0 B_{\mathbf{k}'\mathbf{q}s}^{(1)\dagger} B_{\mathbf{k}\mathbf{q}s}^{(2)\dagger}$.

The c-bosons, each with one fluxon, will be called the fundamental (f) c-bosons. Their energies $w_q^{(j)}$ are obtained from [8]

$$w_q^{(j)} \Psi(\mathbf{k}, \mathbf{q}) = \epsilon_{|\mathbf{k}+\mathbf{q}|}^{(j)} \Psi(\mathbf{k}, \mathbf{q}) - (2\pi\hbar)^{-2} v_0^* \int' d^2 k' \Psi(\mathbf{k}', \mathbf{q}), \quad (3)$$

where $\Psi(\mathbf{k}, \mathbf{q})$ is the reduced wavefunction for the fc-boson; we neglected the fluxon energy. The v_0^* denotes the strength after the ladder diagram binding, see below. For small q , we obtain

$$w_q^{(j)} = w_0 + (2/\pi) v_F^{(j)} q, \quad w_0 = \frac{-\hbar \omega_D}{\exp(v_0^* D_0)^{-1} - 1}, \quad (4)$$

where $v_F^{(j)} \equiv (2\epsilon_F/m_j)^{1/2}$ is the Fermi velocity and $D_0 \equiv D(\epsilon_F)$ the density of states per spin. Note that the energy $w_q^{(j)}$ depends *linearly* on the momentum q .

The system of free fc-bosons undergoes a Bose-Einstein condensation (BEC) in 2D at the critical temperature [8]

$$k_B T_c = 1.24 \hbar v_F n_0^{1/2}. \quad (5)$$

The interboson distance $R_0 \equiv n_0^{-1/2}$ calculated from this expression is $1.24 \hbar v_F (k_B T_c)^{-1}$. The boson size r_0 calculated from Eq. (4), using the uncertainty relation ($q_{\max} r_0 \sim \hbar$) and $|w_0| \sim k_B T_c$, is $(2/\pi) \hbar v_F (k_B T_c)^{-1}$, which is a few times smaller than R_0 . Hence, the bosons do not overlap in space, and the model of free bosons is justified. For GaAs/AlGaAs, $m^* = 0.067 m_c$, m_c = electron mass. For the 2D electron density 10^{11} cm^{-2} , we have $v_F = 1.36 \times 10^6 \text{ cm s}^{-1}$. Not all electrons are bound with fluxons since the simultaneous generations of \pm fc-bosons is required. The minority carrier (“hole”) density controls the fc-boson density. For $n_0 = 10^{10} \text{ cm}^{-2}$, $T_c = 1.29 \text{ K}$, which is reasonable.

In the presence of Bose condensate below T_c the unfluxed electron carries the energy $E_k^{(j)} = \sqrt{\epsilon_k^{(j)2} + \Delta^2}$, where the quasi-electron energy gap Δ is the solution of

$$1 = v_0 D_0 \int_0^{\hbar \omega_D} d\epsilon \frac{1}{(\epsilon^2 + \Delta^2)^{1/2}} \times \left\{ 1 + \exp \left[-\beta (\epsilon^2 + \Delta^2)^{1/2} \right] \right\}^{-1}, \quad \beta \equiv \frac{1}{k_B T}. \quad (6)$$

Note that the gap Δ depends on T . At T_c , there is no condensate and hence Δ vanishes.

Now the moving fc-boson below T_c has the energy \tilde{w}_q obtained from

$$\tilde{w}_q^{(j)} \Psi(\mathbf{k}, \mathbf{q}) = E_{|\mathbf{k}+\mathbf{q}|}^{(j)} \Psi(\mathbf{k}, \mathbf{q}) - (2\pi\hbar)^{-2} v_0^* \int' d^2 k' \Psi(\mathbf{k}', \mathbf{q}), \quad (7)$$

where $E^{(j)}$ replaced $\epsilon^{(j)}$ in Eq. (3). We obtain

$$\tilde{w}_q^{(j)} = \tilde{w}_0 + (2/\pi)v_F^{(j)}q \equiv w_0 + \epsilon_g + (2/\pi)v_F^{(j)}q, \quad (8)$$

where $\tilde{w}_0(T)$ is determined from $1 = D_0 v_0 \int_0^{\hbar\omega_D} d\epsilon [|\tilde{w}_0| + (\epsilon^2 + \Delta^2)^{1/2}]^{-1}$. The energy difference: $\tilde{w}_0(T) - w_0 \equiv \epsilon_g(T)$ represents the T -dependent *energy gap*. The energy \tilde{w}_g is negative. Otherwise, the fc-boson should break up. This limits $\epsilon_g(T)$ to be $|w_0|$ at 0 K. The ϵ_g declines to zero as the temperature approaches T_c from below.

The fc-boson, having the linear dispersion (8), can move in all directions in the plane with the constant speed $(2/\pi)v_F^{(j)}$. The supercurrent is generated by the \pm fc-bosons condensed monochromatically at the momentum directed along the sample length. The supercurrent density (magnitude) J , calculated by the rule: (charge e^*) \times (carrier density n_0) \times (drift velocity v_d), is

$$J \equiv e^* n_0 v_d = e^* n_0 (2/\pi) |v_F^{(1)} - v_F^{(2)}|. \quad (9)$$

The induced Hall field (magnitude) E_H equals $v_d B$. The magnetic flux is quantized $B = n_\phi (h/e)$, $n_\phi =$ fluxon density. Hence we obtain

$$\rho_H \equiv \frac{E_H}{J} = \frac{v_d B}{e^* n_0 v_d} = \frac{1}{e^* n_0} n_\phi \left(\frac{h}{e} \right). \quad (10)$$

If $e^* = e$, $n_\phi = n_0$, we obtain $\rho_H = h/e^2$ in agreement with the plateau value observed.

The model can be extended to the integer QHE at $\nu = P$ ($Q = 1$). The field magnitude is less. The LL degeneracy (eBA/h) is linear in B , and hence the lowest P LL's must be considered. The fc-boson density n_0 per LL is the electron density n_e over P and the fluxon density n_ϕ is the boson density n_0 over P :

$$n_0 = n_e/P, \quad n_\phi = n_0/P. \quad (11)$$

At $\nu = 1/2$ there are c-bosons, each with two fluxons. The c-fermions have a Fermi energy. The \pm c-fermions have effective masses. The Hall resistivity ρ_H has a B -linear behavior while the resistivity ρ is finite.

Let us now take a general case $\nu = P/Q$, odd Q . Assume that there are P sets of c-fermions with $Q - 1$ fluxons, which occupy the lowest P LL's. The c-fermions subject to the available B -field form c-bosons with Q fluxons. In this configuration the c-boson density n_0 and the fluxon density n_ϕ are given by Eqs. (11). Using Eqs. (10) and (11) and assuming the fractional charge [9]

$$e^* = e/Q, \quad (12)$$

we obtain

$$\rho_H \equiv \frac{E_H}{J} = \frac{v_d}{e^* n_0 v_d} n_\phi \left(\frac{h}{e} \right) = \frac{Q}{P} \frac{h}{e^2}, \quad (13)$$

as observed. In our theory the integer Q denotes the number of fluxons in the c-boson and the integer P the

number of the LL's occupied by the parental c-fermions, each with $Q - 1$ fluxons.

Our Hamiltonian in Eq. (2) can generate and stabilize the c-particles with an arbitrary number of fluxons. For example a c-fermion with two fluxons is generated by two sets of the ladder diagram bindings, each between the electron and the fluxon. The ladder diagram binding arises as follows. Consider a hydrogen atom. The Hamiltonian contains kinetic energies of the electron and the proton, and the attractive Coulomb interaction. If we regard the Coulomb interaction as a perturbation and use a perturbation theory, we can represent the interaction process by an infinite set of ladder diagrams, each ladder step connecting the electron and the proton. The energy eigenvalues of this system is not obtained by using the perturbation theory but they are obtained by solving the Schrödinger equation directly. This example indicates that a two-body bound state is represented by an infinite set of ladder diagrams and that the binding energy (the negative of the ground-state energy) is calculated by a non-perturbative method.

Jain introduced the effective magnetic field [10]

$$B^* \equiv B - B_\nu = B - (1/\nu)n_e(h/e) \quad (14)$$

relative to the standard field for the composite (c -) fermion at the even-denominator fraction. We extend this to the bosonic (odd-denominator) fraction. This means that the c -particle moves field-free at the exact fraction. The c -particle is viewed as the quasiparticle containing an electron circulating around Q fluxons. The jumping of the guiding centers (the CM of the c -particle) can occur as if they are subject to no B -field at the exact fraction. The excess (or deficit) of the magnetic field is simply the effective magnetic field B^* . The plateau in ρ_H is formed due to the Meissner effect. Consider the case of zero temperature near $\nu = 1$. Only the energy E matters. The fc-bosons are condensed with the ground-state energy w_0 , and hence the system energy E at $\nu = 1$ is $2N_0 w_0$, where N_0 is the number of $-$ fc-bosons (or $+$ fc-bosons). The factor 2 arises since there are \pm fc-bosons. Away from $\nu = 1$ we must add the magnetic field energy $(2\mu_0)^{-1} A(B^*)^2$, so that

$$E = 2N_0 w_0 + (2\mu_0)^{-1} A(B^*)^2. \quad (15)$$

When the field is reduced, the system tries to keep the same number N_0 by sucking in the flux lines. Thus the magnetic field becomes inhomogeneous outside the sample, generating the magnetic field energy $(2\mu_0)^{-1} A(B^*)^2$. If the field is raised, the system tries to keep the same number N_0 by expelling out the flux lines. The inhomogeneous fields outside raise the field energy as well. There is a critical field $B_c^* = (4\mu_0|w_0|)^{1/2}$. Beyond this value, the superconducting state is destroyed, generating a symmetric exponential rise in R . In our discussion of the Hall resistivity plateau we used the fact that the ground-state energy w_0 of the fc-boson is negative, that is, the c -boson is bound. Only then the critical field $B_c^* = (4\mu_0|w_0|)^{1/2}$

can be defined. Here the phonon exchange attraction played an important role. The repulsive Coulomb interaction, which is the departure point of the prevalent theories [2,11], cannot generate a bound state.

In the presence of the supercondensate the non-condensed c-boson has an energy gap ϵ_g . Hence the non-condensed c-boson density has the activation energy type exponential temperature-dependence:

$$\exp[-\epsilon_g/(k_B T)] . \quad (16)$$

In the prevalent theories the energy gap for the fractional QHE is identified as the sum of the creation energies of a quasi-electron and a quasi-hole [11]. With this view it is difficult to explain why the activation-energy type temperature dependence shows up in the steady-state quantum transport. Some authors argue that the energy gap ϵ_g for the integer QHE is due to the LL separation $= \hbar\omega_0$. But the separation $\hbar\omega_0$ is much greater than the observed ϵ_g . Besides from this view one cannot obtain the activation-type energy dependence.

The BEC occurs at each LL, and therefore the c-boson density n_0 is less for high P , see Eq. (11), and the strengths become weaker as P increases.

We are now ready to discuss the QHE under radiation. The experiments by Mani *et al.* [4] indicate that the applied radiation excites a large number of “holes” in the system. Using these “holes” and the preexisting “electrons” the phonon exchange can pair-create \pm c-bosons, which condense below T_c in the excited channel. The condensed c-bosons in motion are responsible for the supercurrent. In the presence of the condensed c-bosons, the non-condensed c-bosons have an energy gap ϵ_g , and therefore they are absent at 0 K. The c-fermions in the base channel will have the energies $E_p = (\epsilon_p^2 + \Delta^2)^{1/2}$ but

their energy spectra have no gap. Hence they are not completely suppressed at the lowest temperatures. They contribute a small normal current, and are responsible for the B -linear behavior of the Hall resistivity observed.

Mani *et al.* experiments [1], Fig. 2, show that the strength of the superconducting state does not change much for the radiation frequency ω in the range (0.47, 110) GHz. This feature may come as follows. The 2D density of states for the conduction electrons associated with the circular Fermi surface is independent of the electron energy, and hence the number of the excited electrons is roughly independent of the radiation energy (frequency). The “hole”-like excitations are absent with no radiation. We suspect that the “hole”-band edge is a distance ϵ_0 away from the system’s Fermi level. This means that if the radiation energy $\hbar\omega$ is less than ϵ_0 , the radiation can generate no superconducting state. This feature can be checked by applying radiation of frequencies lower than 0.47 GHz.

In summary the QHE under radiation is the QHE at the upper channel. The condensed c-bosons generate a superconducting state with a gap ϵ_g in the c-boson energy spectrum. The supercondensate changes the c-fermion energy from ϵ_k to $(\epsilon_k^2 + \Delta^2)^{1/2}$ in the base channel. This energy spectrum has no gap, and hence the c-fermions cannot be suppressed completely at the lowest temperatures, and generate a finite resistive current accompanied by the Hall field. This explains the B -linear Hall resistivity. Our microscopic theory can be tested experimentally by the detection of (a) the “hole”-like excitations by a circularly polarized laser, (b) the bosonic state at $\nu = 4/3$ and $1/3$, (c) the “hole” band edge.

We thank Dr. Mani for fruitful discussion.

-
- [1] R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayana-murti, W. B. Johnson, and V. Umansky, *Nature* **420**, 646 (2002).
 - [2] D. C. Tsui, H. L. Störmer and A. C. Gossard, *Phys. Rev. Lett.* **48**, 1559 (1982).
 - [3] M. A. Zudov, R. R. Du, L. N. Pfeiffer and K. W. West, *Phys. Rev. Lett.* **90**, 046807 (2003).
 - [4] R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayana-murti, W. B. Johnson and V. Umansky, *cond-mat/0310474*.
 - [5] J. C. Phillips, *Sol. St. Comm.* **127**, 233 (2003); A. C. Durst, S. Sachdev, N. Read, and S. M. Girvin, *Phys. Rev. Lett.* **91**, 086803 (2003); A. V. Andreev, I. L. Aleiner, and A. J. Millis, *Phys. Rev. Lett.* **91**, 056803 (2003); J. Shi and X. C. Xie, *Phys. Rev. Lett.* **91**, 086801 (2003); F. S. Bergeret, B. Huckestein, and A. F. Volkov, *Phys. Rev. B* **67**, 241303 (2003).
 - [6] S. Fujita, S. Godoy and D. Nguyen, *Found. Phys.* **25**, 1209 (1995).
 - [7] J. Bardeen, L. N. Cooper and J. R. Schrieffer, *Phys. Rev.* **108**, 1175 (1957).
 - [8] S. Fujita, Y. Tamura and A. Suzuki, *Mod. Phys. Lett. B* **15**, 817 (2001).
 - [9] R. B. Laughlin, *Phys. Rev. Lett.* **50**, 1395 (1983); F. D. M. Haldane, *Phys. Rev. Lett.* **51**, 605 (1983).
 - [10] J. K. Jain, *Phys. Rev. Lett.* **63**, 199 (1989); *Phys. Rev. B* **40**, 8079 (1989); *ibid.* **41**, 7653 (1990).
 - [11] R. E. Prange and S. M. Girvin, (eds) *The Quantum Hall Effect*, 2nd Ed., (Springer-Verlag, New York, 1990).