Quantum theory of photonic crystal polaritons

D. Gerace, M. Agio, and L. C. Andreani

Istituto Nazionale per la Fisica della Materia and Dipartimento di Fisica \Alessandro Volta",

Universita di Pavia, via Bassi 6, 27100 Pavia, Italy

We form ulate a fullquantum mechanical theory of the interaction between electrom agnetic modes in photonic crystal slabs and quantum well excitons embedded in the photonic structure. We apply the form alism to a high index dielectric layer with a periodic patterning suspended in air. The strong coupling between electrom agnetic modes lying above the cladding light line and exciton center of mass eigenfunctions manifests itself with the typical anticrossing behavior. The resulting band dispersion corresponds to the quasi-particles coming from the mixing of electrom agnetic and material excitations, which we call photonic crystal polaritons. We compare the results obtained by using the quantum theory to variable angle relative agreement.

PACS num bers: 42.50 Ct, 42.70 Qs, 71.36.+ c, 73.20 Mf, 78.20 Bh

Introduction. Since the pioneering works of more than fieen years ago by E.Yablonovitch and S.John [1], a great deal of research in physics has been dedicated to the study of the optical properties of photonic crystals [2]. In the last few years much attention has been devoted to photonic crystal (PC) structures embedded in planar dielectric waveguides, which are also called photonic crystal slabs. These are systems with a periodic dielectric response in the plane of the waveguide, while a dielectric mism atch is used to con ne the electrom agnetic eld in the vertical direction.

O ne of the main issues of PC slabs is the existence of the light line problem : only photonic modes lying below the cladding light line are truly guided and with zero intrinsic linew idth, while the modes lying above the light line are radiative. A method to calculate the photonic band dispersion and the intrinsic linew idth (com plex energies) both below and above the light line has been recently proposed [3, 4]. Until now, how ever, the physics of PCs has been considered mostly for what concerns the \optical" point of view, that is studying the propagation of light in periodic dielectric media.

In this work we analyze the radiation-matter interaction in PC slabs, by considering the elects of the interplay between the electrom agnetic eld and sem iconductor quantum well (QW) excitons. In particular, we show that polaritonic elects are present in PC slabs when the exciton-photon coupling is larger than the intrinsic radiative linew idth of a photonic mode above the cladding light line. The band dispersion is greatly modi ed in the vicinity of the excitonic resonance, leading to the formation of mixed states which are analogous to exciton-polaritons in bulk sem iconductors [5] and microcavities [6{8] and which we call photonic crystal polaritons (PCPs). Experimental results showing the form ation of PCPs in organic-based systems were previously reported in Ref. 9.

Theory. Here we brie y describe the quantum theory of PCPs, starting with the second quantized total ham iltonian of the system, which is given by

$$\hat{H} = \frac{X}{\underset{k_{1},m_{1}}{\overset{k_{1}}{x_{1}}}} \frac{X}{a_{k_{1}}} + \frac{X}{\underset{k_{2}}{x_{1}}} \frac{X}{a_{k_{1}}} + \frac{X}{a_{k_{1}}} \frac{X}{a_{k_{1}}} + \frac{X}{a_{k_{1}}} \frac{X}{a_{k_{1}}} + \frac{X}{a_{k_{1}}} (\hat{a}_{k_{1}} + \hat{a}_{k_{1}}^{y}) (\hat{b}_{k}^{y} - \hat{b}_{k_{1}})$$

$$+ \frac{X}{\underset{k_{2},m_{1},m_{2}}{\overset{C}{x_{1}}}} \frac{C_{k_{1}} C_{k_{1}}}{a_{k_{1}}} (\hat{a}_{k_{1}} + \hat{a}_{k_{1}}^{y}) (\hat{a}_{k_{1}$$

In Eq. (1), the rst term indicates the photonic band dispersion (real part of the complex eigenenergies), \hat{a}_{kn} (\hat{a}_{kn}^{y}) being the destruction (creation) operators of a photon with wave vector k and band number n. These energies are obtained by expanding the magnetic eld in term softhe guided modes of the e ective waveguide, that is the one with an average dielectric constant, and then by solving the Maxwell equation as a linear eigenvalue problem $[\beta]$. The imaginary part of the photonic modes is calculated by using a perturbative approach [4]. If we restrict our considerations to the system schematically shown in the inset of Fig. 1a, the direction of periodicity is x and thus the wave vector is given by the component k_x . The modes in a PC slab suspended in air can be classied as even (odd) with respect to specular re ection through the plane xy, and even (odd) with respect to the plane of incidence (that is xz in the case considered here). Throughout this paper we consider only modes which are spatially even with respect to xy plane, and odd with respect to the vertical mirror plane (even TE modes). These modes interact with transverse QW excitons. The excitonic problem, whose solution gives the energies \sim_k , is treated by solving the Schrödinger equation for the exciton center of mass envelope function in a periodic piecew ise constant potential having the same patterning as the photonic crystal structure. The in-plane wave vector and an integer are good quantum num bers for the exciton wavefunctions, due to the periodic potential and to the spatial dispersion of the center



FIG.1: (a) Realand (b) in aginary parts of the complex photonic energies of the even TE m ode as a function of the wave vector in the rst Brillouin zone, for the system shown in the inset, without excitonic resonance (no QW); (c) realand (d) in aginary parts for the same structure with a QW at the center of the dielectric core, with an excitonic resonance at $\sim_0 = 1.4 \text{ eV}$. The dashed lines in (a) and (c) represent the light line in air.

of m ass; the corresponding destruction (creation) operators are \hat{b}_k (\hat{b}_k^v). The exciton-photon coupling m atrix elements, C $_{kn}$, are calculated in term s of the m icroscopic physical quantities as

$$C_{kn} = \frac{2 e^{2} \sim \frac{2}{k}}{!_{kn}} h_{k}^{(exc)} j E_{kn}(r_{j}) f^{j}$$
(2)

In Eq. (2), ${}_{k}^{(exc)}$ is the all-electron exciton wavefunction, E_{kn} is the electric eld prole for the photonicm ode at frequency $!_{kn}$ in the PC slab, and the sum is over all the QW electrons. In particular, C_{kn} is found to depend on the overlap between the exciton envelope function and the transverse electric eld (for what concerns TE modes). It is proportional to $(f=S)^{1=2}$, where f=S is the oscillator strength per unit area which depends on the QW thickness [8]. The ham iltonian is diagonalized by using a generalized H op eld transform ation to expand new destruction (creation) operators \hat{P}_{k} (\hat{P}_{k}^{Y}) as a linear combination of \hat{a}_{kn} (\hat{a}_{kn}^{Y}) and \hat{b}_{k} (\hat{b}_{k}^{Y}), with the condition $[\hat{P}_{k};\hat{H}] = E_{k}\hat{P}_{k}$ [10]. The new eigenenergies E_{k}

correspond to m ixed excitations of radiation and m atter, i.e. the PCPs.

All the results presented in this paper refer to a model structure, namely a high index dielectric core suspended in air with a one dimensional patterning, and a QW placed at the center of it (see inset of Fig. 1a). The dielectric constant of the sem iconductor-based core layer is set to the value " = 12, the dielectric constant of the air being sim ply "air = 1. The slab thickness and the air fraction are set to the values d=a = 0.2 and r=a = 0.3 respectively, the lattice constant being a = 350 nm . A QW of thickness $L_{QW} = 8$ nm is considered, and the oscillator strength per unit area is set to the typical value $f=S = 8.4 \quad 10^{12}$ cm⁻². The intrinsic radiative exciton linew idth is assumed to be = 0.1 meV.

R esults and D iscussion. In Fig. 1a we show the photonic band dispersion in the energy range 1.1-1.7 eV. In Fig. 1b we display the corresponding in aginary part as a function of the wave vector, which shows a maximum at about 1.4 eV. The in aginary part goes to zero at $k_x = 0$, that is at norm al incidence, and at $k_x = 0.74$; this last behavior is due to the crossing of the light line,



FIG.2: (a) Calculated surface re extance of a TE polarized plane wave incident on the structure of the inset of Fig.1a. The excitonic resonance is set at $\sim_0 = 1.58 \text{ eV}$; the angle of incidence is varied from = 0 to = 20, with a step of 1. (b) Real and (c) in aginary parts of the com plex energies of the photonic crystal polaritons, calculated with the quantum theory. The squared points in (b) are extracted from the calculated re extrace curves shown in (a), and the polariton splitting is indicated by an arrow.

as shown in Fig. 1a, corresponding to the photonic mode becoming truly guided and stationary. If the photonic in aginary part is larger than the exciton-photon coupling matrix element (which is of the order of a few meV), a QW placed at the center of the PC slab does not produce any important change in the photonic band dispersion. Indeed, this result is shown in Figs. 1c and 1d, in which we display the real and in aginary parts of the complex eigenenergies com ing from the diagonalization of Eq. (1)with an excitonic resonance at $\sim _0 = 1.4 \text{ eV}$. In this weak coupling regime the photon and the exciton are almost uncoupled, as it can be seen from the crossing of the two dispersion relations in Fig. 1c. The imaginary part of the exciton increases by an order of m agnitude correspondingly to the crossing point, but this has a negligible e ect on the photonic radiative linew idth, which is still an order of m agnitude larger than the excitonic one (see Fig.1d).

In order to observe the strong coupling regime, the energy of the excitonic resonance has to lie where the in aginary part of the photonic mode is smaller than the coupling matrix element. From Fig. 1a and 1b we see that for $\sim_0 = 1.58$ eV the imaginary part of the corresponding photonic band is about Im (\sim !) = 10⁻³ eV. In Fig. 2a we show the results of a variable angle reectance calculation, which is done by a scattering matrix approach [11]. The dielectric function of the QW layer is frequency dependent, with a resonance at $\sim_0 = 1.58$ eV. It is known that the sharp features appearing in the reectance spectrum correspond to the excitation of pho-

to the excitation of photonic modes above the light line [9, 12], thus giving a

point (k;!) of the corresponding photonic band dispersion; the wave vector component parallel to the surface is given by $k = (!=c) \sin \cdot The(k;!)$ points extracted from the results of Fig. 2a are compared to the dispersion of PCPs in Fig. 2b, in which the squares are the scattering matrix results. The quantum theory is in excellent agreement with the classical approach. In particular, both theories con m the anticrossing behavior of exciton and photon modes, which is a clear effect of the strong coupling regime. With the parameters used in this work, the polariton splitting is as high as 10 m eV at the anticrossing point $k_x = 0.14$, as shown in Fig. 2b. This splitting is found to be slightly larger than in sem iconductor-based m icrocavities [7, 8]. In Fig. 2c, nally, the imaginary part of the PCPs complex eigenenergies is shown. At $k_x = 0.14$ the imaginary parts of the upper and lower polariton branches become equal to the same value Im (E) = 10^{-3} eV, thereby indicating that m ixed states of radiation and m atter form in the PC slab. The dispersionless curve at Im (E) = 10^{4} eV in Fig. 2c corresponds to the uncoupled excitonic modes.

C on clusions. We have shown that the quantum theory of the interaction between electrom agnetic modes in photonic crystal slabs and QW excitons can describe both the weak and strong coupling regimes. Moreover, we have found a very good agreem ent between the quantum theory and a classical approach based on a scattering matrix method. We thus conclude that new mixed states of radiation and matter excitations can be experimentally measured in sem iconductor-based photonic crystal slabs by using variable angle rejectance or transmittance techniques, provided that the experim ental conditions required for being in the strong coupling regime could be satis ed.

A cknow ledgem ents. This paper is dedicated to the memory of G iovanna Panzarini, whose research on exciton polaritons in 3D m icrocavities was of great inspiration for the present developments. This work was supported by M IUR through Co n 2002, and by INFM through PRA PHOTONIC.

- [1] E. Yablonovitch, Phys. Rev. Lett. 58, 2059 (1987);
 S. John, Phys. Rev. Lett. 58, 2486 (1987).
- [2] For recent reviews, see e.g. papers in IEEE J.Quantum Electron. 38, Feature Section on Photonic Crystal Structures and Applications, ed. by T.F.K rauss and T.Baba, pp. 724-963 (2002).
- [3] L.C.Andreaniand M.Agio, IEEE J.Quantum Electron. 38,891 (2002).

- [4] L.C.Andreani, phys. stat. sol. (b) 234, 139 (2002).
- [5] J. J. Hop eld, Phys. Rev. 112, 1555 (1958);
 V. M. Agranovich, J. Exp. Theor. Phys. 37, 430 (1959) [Sov.Phys.JETP 37, 307 (1960)].
- [6] C.W eisbuch, M.N ishioka, A. Ishikawa, and Y.Arakawa, Phys. Rev. Lett. 69, 3314 (1992).
- [7] M.S.Skolnick, T.A.Fisher, and D.M.W hittaker, Sem icond.Sci.Technol. 13, 645 (1998).
- [8] L. C. Andreani, in Electron and Photon Connement in Semiconductor Nanostructures, ed. by B. Deveaud, A. Quattropani and P. Schwendimann, IOS Press (Amsterdam, 2003).
- [9] T. Fujita, Y. Sato, T. Kuitani, and T. Ishihara, Phys. Rev. B 57, 12428 (1998).
- [10] G.Panzariniand L.C.Andreani, Phys. Rev. B 60, 16799 (1999).
- [11] D.M.W hittaker and I.S.Culshaw, Phys. Rev. B 60, 2610 (1999).
- [12] V. N. Astratov, D. M. W hittaker, I. S. Culshaw, R. M. Stevenson, M. S. Skolnick, T. F. Krauss, and R. M. De La Rue, Phys. Rev. B 60, 16255 (1999).