PREPRINT

Length-dependent oscillations of the conductance through atom ic chains: The importance of electronic correlations

Rafael A.Molina¹, Dietmar W einmann² and Jean-Louis Pichard^{1;3}

¹ CEA/DSM, Service de Physique de l'EtatCondense, Centre d'Etudes de Sachy, 91191 Gif-sur-Yvette, France

² Institut de Physique et Chim ie des Materiaux de Strasbourg, UMR 7504 (CNRS-ULP),
23 rue du Loess, BP 43, 67034 Strasbourg Cedex 2, France

³ Laboratoire de Physique Theorique et Modelisation, Universite de Cergy-Pontoise, 95031 Cergy-Pontoise Cedex, France

PACS.73.23.-b { Electronic transport in m esoscopic systems. PACS.05.60.Gg { Quantum transport. PACS.73.63.Nm { Quantum wires.

A bstract. { We calculate the conductance of atom ic chains as a function of their length. U sing the Density M atrix R enormalization G roup algorithm for a many-body model which takes into account electron-electron interactions and the shape of the contacts between the chain and the leads, we show that length-dependent oscillations of the conductance whose period depends on the electron density in the chain can result from electron-electron scattering alone. The amplitude of these oscillations can increase with the length of the chain, in contrast to the result from approaches which neglect the interactions.

Q uantum wires of atom ic dimension are among the basic components of molecular electronics. The knowledge of the transport properties of such structures will be crucial for the design of devices, and the understanding of their properties is a major challenge in nanoelectronics. Experimentally, chains of atom s between two electrodes have been formed and investigated [1{3], and the conductance of molecules like H_2 has been measured [4]. Besides the possible applications in molecular devices, this is of great general interest since electronic correlations alter considerably the properties of one-dimensional system s. Ferm i liquid theory completely looses its applicability and Luttinger liquid [5] behavior appears. This has been observed for the case of metallic single wall carbon nanotubes [6]. A recent measurement on atom ic Au, Pt and Ir wires between break junction contacts [7] has revealed oscillations of the conductance has maxim a while it is reduced when the number of atom s is even. This behavior is obtained after averaging over many realizations of the experiment in which the precise form of the contacts between the electrodes and the atom ic chain might be di erent.

Since a few years, the conductance of chains of atom shas been the subject of an increasing theoretical activity (see Sect. 11.5.1 of Ref. [3] for an overview), and parity-oscillations had

c EDP Sciences

EUROPHYSICS LETTERS

been expected for chains of m on ovalent atom s like Au. The experim ental observation of evenodd oscillations in chains of the polyvalent atom sPt and Ir [7] how ever calls form ore profound studies.

For a non-interacting half- lled tight-binding chain coupled to leads, Fabry-Perot like interferences lead to size-dependent parity oscillations in the conductance [8]. Taking into account the electron-electron interactions perturbatively [9] within the Hubbard model and using density functional theory for chains of N a atom s [10] indicate that these parity oscillations of the conductance persist once interactions are included in the model. Local charge neutrality has been identified [10{12] as the main cause of the even-odd oscillations in chains of monovalent atom swith a half- lled valence band. How ever, the phase of the parity oscillations depends on details of the model used to describe the coupling of the chain to the leads [11]. A four-atom period was obtained from density functional calculations for the conductance of atom ic A 1 chains [13]. The analysis of the phase of these oscillations lead to the conclusion that the electron density in the chain is determined by the Fermi energy in the leads, and that local charge neutrality is not satis ed inside the chains.

Taking into account the full electronic correlations within a model of spinless ferm ions with next-neighbor interactions at half lling we showed that the even-odd oscillations can simply be the result of the presence of strong electronic correlations [14]. We even found that the interactions could enhance the amplitude of the parity oscillations in the conductance.

In this work we present a study of the length dependent oscillations in the conductance of atom ic wires, taking fully into account the electronic correlations inside the chain. A fier presenting our num erical results for spinless correlated ferm ions in a one-dimensional chain, we will discuss the role of the shape of the contacts to the leads and the dependence of the conductance-oscillations on the particle density. These results will be contrasted with the theoretical predictions for m odels which neglect the interactions. This allows to identify signatures of the electronic correlations in the length dependent conductance oscillations which could be observed experimentally.

To calculate the conductance of a chain where electronic correlations m ay be strong, and which is coupled to leads, we consider the m odel H am iltonian

$$H = t \begin{pmatrix} X & X^{L} \\ C_{i}^{y}C_{i} + C_{i}^{y}C_{i} \end{pmatrix} + U_{i} [h_{i} + V_{i}] [h_{i} + V_{i}] :$$
(1)

The rst term describes the kinetic energy in the atom ic chain (from site 1 to L) and in the leads. The operators c_i^y create spinless ferm ions on the sites i of an in nite one-dimensional tight binding model, and the hopping element t = 1 is taken constant and sets the energy scale. The second term accounts for the interactions of strength U_i between particles on adjacent sites i 1 and i inside the atom ic chain. Here, $n_i = c_i^y c_i$ is the density operator and the potential V_+ allows to adjust the particle density in the chain. A density = 1=2 corresponding to half lling is obtained by $V_+ = 1=2$ which leads to particle-hole symmetry. To get other values for the density, V_+ is chosen in a self-consistent iterative way as described in R ef. [15]. Though the one-dimensional leads are of course not realistic, the suppression of the electronic correlations by the screening in realm assive electrodes is nevertheless accounted for by the absence of interactions. The shrinking diam eter of the contacts between the leads and the chain is modeled by a linear increase of the interaction strength U_i from zero to its full value $U_i = U$ over L_c sites [14].

To determ ine the conductance of the atom ic chain in the presence of electronic correlations, we exploit the connection between its zero-tem perature linear conductance and the persistent current or the charge sti ness of a non-interacting ring in which the interacting system is

2



Fig.1 { Conductance through interacting atom ic chains at half lling (= 1=2), as a function of their length L for the case of abrupt contacts ($L_c = 1$) and di event values of the interaction strength.

embedded [14,16,17]. We use the density matrix renorm alization group (DMRG) algorithm [18] to obtain the charge sti ness of the model (1) after closing it to a ring. For a detailed discussion of the method and the veri cation of its validity see Ref. [15] and references therein.

Results for the length-dependence of the dimensionless conductance g in units of the conductance quantum e^2 =h are shown in Fig. 1 for the case of half lling. The presence of interactions leads to clear parity oscillations in the conductance through the atom ic chain, with perfect conductance g = 1 for chains containing an odd number of sites, independent of the interaction strength. For even chains, the conductance is reduced when the interaction strength U increases. This reduction becomes more pronounced in longer chains.

In the limit of very strong interactions, the parity oscillations at half lling can be understood by considering the one-body hopping terms in (1) as a perturbation to the interaction term (for simplicity, we take abrupt contacts and $U_i = U$ for all i = f2;:::Lg). In a chain of odd length L, the ground state of the interaction term alone is a superposition of two degenerate components. One of them corresponds to N = (L + 1)=2 particles occupying the odd sites i = f1;3;:::;Lg, for the other N 1 particles occupy the even sites of the chain. For in nitesim al hopping t=U 1, the two degenerate components become e coupled through sequences of N one-particle hopping processes. At half lling, when the Ferm i energy in the non-interacting leads vanishes, this corresponds to a one-particle resonant tunneling situation. This results in perfect transm ission and g = 1, independent of the weakness of the coupling between the e ective bound state and the leads which is of the order of t(t=U)^N ¹.

For the case of an interacting chain with even length L, the ground state in the absence of the hopping term of (1) is also a superposition of two components, $j_{ri} > = \sum_{n=1}^{N} c_{2n}^{V} \mathcal{D}i$ $\sum_{n=1}^{N} c_{2n-1}^{y}$ Di. They correspond to the same number N = L=2 of particles, and $j_{1}i =$ occupying the even and the odd sites of the chain, respectively. Even though these degenerate components are coupled by sequences of N one-particle hopping processes as in the odd case, these sequences do not involve particles hopping between the interacting chain and the leads. Therefore, these processes do not contribute directly to the transport through the chain. The process which dom inates zero-tem perature transport through the chain in the even case is coherent cotunneling. Taking into account the lowest order processes in t=U, involving transitions via N -particle states with modi ed charge con guration and states having N 1 U $^{\rm L}$ of the conductance with the interaction particles in the chain leads to a decrease strength U for even chain length L. This is a signature of the Mott insulating behavior occurring for in nitely long chains at U > 2. The strong suppression of the even conductance with the length at U = 4 in Fig. 1 can be understood as a precursor of the insulating behavior. In the lim it of weak interactions, we can calculate the conductance as $g = 4 g_{L_{11}}(0) j$,



Fig.2 { Conductance as a function of the number of sites in the interacting region for = 1=2 and symmetric contacts of length $L_c = 3$ (left), and asymmetric contacts of length $L_{c1} = 4$, $L_{cr} = 2$ (right). The geometries we have in m ind are sketched above the graphs.

where $G_{L,1}(0)$ is the G reen's function for the propagation of a particle from the st to the last site of the chain at the Ferm i energy of the leads $E_F = 0$, within the full many-body H am iltonian [9,19], evaluated at zero tem perature. We can easily calculate this G reen's function in second order perturbation theory in U. The conductance of odd chains is not a ected by this perturbation, but in the case of even chains it is reduced by an am ount proportional to U², consistent with the num erical results for sm all chains and weak interaction [14]. Such an approach has been used to show that weakly interacting Hubbard chains connected to non-interacting leads exhibit parity oscillations [9].

We now address the role of the shape of the contacts between the atom ic chain and the leads in order to consider the more realistic case of smooth contacts. The conductance for a symmetric setup with a chain connected to the leads through two contacts of length $L_C = 3$ is shown in Fig.2 (left), at half lling. The even-odd oscillations as a function of the chain length persist in the presence of such smooth contacts, and the perfect transmission of odd chains is not a ected. However, the reduction of the conductance in even chains is less pronounced when the contacts are smooth and the amplitude of the oscillations is smaller. Nevertheless, the dependence of the oscillations on the interaction strength and on the length of the chain are similar to the situation of abrupt contacts. Only in the unrealistic limit of in nitely smooth contacts one expects perfect transmission for even chains as for the odd ones [14] and the parity oscillations disappear.

W hile it seems reasonable to assume symmetric contacts when describing certain experimental situations like break junctions, it is dicult to exclude the case of a slightly asymmetric situation where the two ends of the atom ic chain are connected to the electrodes in dicerent ways. This breaking of the rection symmetry of the model has consequences that can be seen in Fig. 2 (right). Here, the left and right contacts are assumed to be of length $L_{Cl} = 4$ and $L_{Cr} = 2$, respectively. Most strikingly, the conductance obtained for odd chains no longer reaches the unitary limit g = 1 and decreases with increasing interaction strength and chain length. A s compared to the symmetric situation, the conductance values are reduced for the even chains as well. However, the parity oscillations persist and the phase of the oscillations is robust as well: one has maxim a of the conductance for odd and minim a for even chains.

The parity oscillations that we have studied so far occur in the case of half lling which corresponds to chains of monovalent atom s with a half-lled valence band like N a and A u. A t other values of the lling, the period of the length-dependent oscillations in the conductance is



Fig.3 { Conductance as a function of the length of an atom ic chain connected to the leads by abrupt contacts for llings = 1=3 (left), = 1=4 (center), and = 1=5 (right).

di erent. To illustrate its dependence on the lling, we show in Fig.3 results for the densities = 1=3, 1/4, and 1/5. For sim plicity, we use abrupt and symmetric contacts. In this case, perfect transmission appears for chain lengths of L = 1 + q = w ith integer q = f0;1;:::q, and the period of the length-dependent oscillations is 1= sites for < 1=2. At larger llings, the density of holes 1 is relevant for the periodicity of the conductance. Furtherm ore, it can be seen that the am plitude of the oscillations is smaller when the lling is lower. This can be understood from the fact that the conductance oscillations in our model are caused by the short range electron-electron interactions. At low lling, this interaction becomes less relevant since the probability to not two particles on adjacent sites of the chain decreases.

Setting the interactions to zero (U = 0) in our model leads to a perfect hom ogeneous chain which exhibits perfect transmission and conductance g = 1 for all llings and lengths L, and the length-dependence of the conductance disappears. This shows that electronic correlations alone can be the origin of length-dependent conductance oscillations in atom ic chains. In a non-interacting model, another scattering mechanism is necessary for reproducing the oscillating behavior of the conductance. A possibility is to suppose that the contacts between the atom ic chain and the electrodes are a source of rejections.

This can be achieved by adding the contact term

$$H_{c} = (t \quad \underline{v}_{1}) \quad c_{1}^{y}c_{0} + c_{0}^{y}c_{1} + (t \quad \underline{v}_{2}) \quad c_{1,+1}^{y}c_{1,+1} + c_{1,-1}^{y}c_{1,+1}$$
(2)

to our H am iltonian (1). This replaces the hopping matrix elements t between the chain and the left and right leads by the couplings v_1 and v_r , respectively. Choosing jv_1j ; $jv_rj < t$, this describes a chain of L atom s coupled to one-dimensional leads by weak links which give rise to Fabry-Perot like interferences. In this model, the particle density in the chain is the same as the one in the leads which is determined by the Fermi intergry. This situation is dimensional assuming local charge neutrality as discussed in R ef. [13]. U sing a transferm atrix form alism we can calculate the transmission at the Fermi intergry, and using Landauer's form ula $g = j_1(k_F)^2$ the conductance

$$g = 16\sin^{4}k_{F} v_{1} v_{1}^{1} v_{r} v_{r}^{1} e^{2ik_{F}L} v_{r}e^{2ik_{F}} v_{1}^{1} v_{1} [Me^{2ik_{F}}]^{1}$$
(3)

of the non-interacting system with the Ferm iwave number k_F . This result is symmetric with respect to the exchange of v_r and v_l .

The only L-dependence of g in (3) is through the factor $e^{2ik_F L}$. Therefore, the result is periodic in L with period $=k_F$. In dimension one, the Ferm is avenumber is related to the

lling through $k_{\rm F} = 0$, and therefore the period of the oscillations is given by 1 = 0 sites as in the interacting case discussed above. However, the phase is dimensional event, except for half-lling,

EUROPHYSICS LETTERS



Fig.4 { Conductance calculated from Eq. (3) as a function of the number of sites in a non-interacting chain which is connected to one-dimensional leads through weak links of strength $v_1 = v_r = 0.75t$. The lling is = 1=3 (left), 1=4 (center), and 1=5 (right).

where one obtains always even-odd oscillations with the maxim a for odd N. This can be seen from Fig. 4, where the analytical result of Eq. (3) is shown for the example of $v_1 = v_r = 0.75$ and di erent llings. A ssum ing local charge neutrality and using a tight-binding model with a lling which is independent of the lling of the leads, the same periods have been found in the

conditions for resonant transport [12]. How ever, the phase of the oscillations is di erent [13].

As compared to the interacting case, the behavior of the non-interacting m odel presents striking di erences. First of all, without interactions the length-dependence is perfectly periodic and the conductance satis esg(L + 1 =) = g(L). Thus, the amplitude of the oscillations is independent of L. This is because the motion of the particles between the contacts is ballistic. In contrast, for interacting particles, the length-dependent oscillations we obtained (see Fig. 1) are superimposed with a decrease of the conductance m inim a and an increase of the amplitude of the oscillations with the length of the chain. For asymmetric contacts, the amplitude remains m ore or less constant, but the conductance decreases with the length for both, m axim a and m inim a of the oscillations (see Fig. 2). W hile such a decrease is not seen in the experimental data for Au chains, Pt and Ir do show [7] such a behavior which is reminiscent of our results for correlated chains, but di cult to explain within non-interacting m odels.

In addition, while the maxima in our correlated model always reach perfect conductance g = 1 in the case of symmetric coupling to the leads, without interactions this is the case at half lling and only for special parameter combinations outside half lling [12]. In the case of asymmetric contacts $v_1 \notin v_r$ the amplitudes of the length-dependent oscillations obtained from the conductance of Eq. 3 are somewhat reduced, but remain clearly observable, similar to the case of weakly correlated chains in Fig. 2. This is consistent with the fact that the oscillations have been observed experimentally [7] in a situation where the contacts were very probably not exactly symmetric.

The period of the length-dependent oscillations predicted by both non-interacting approaches agrees with the result of our numerical calculations for a correlated chain. This indicates that the period is indeed determined by the lling of the conductance band alone, and given by 1=, independent of details like the shape of the contacts between the chain and the leads. How ever, outside half lling, the phase of the oscillations is di erent for all three mentioned approaches. Therefore, we can expect that, in contrast to the period, the phase depends on ne details of the experimental situation. As a consequence, a more re ned analysis than averaging over m any realizations of the preparation of the chain as done in Ref. [7] could be necessary for their observation.

In conclusion, we have studied the e ect of electronic correlations on the length-dependent oscillations observed in atom ic wires. In the absence of interactions, these oscillations can

be understood as an interference e ect between two scatterers connecting the chain to the leads. However, the electrons of an atom ic chain form a correlated system which can by itself be an important source of scattering, and hence of resistance. We have studied the consequences of this additional scattering mechanism within a non-perturbative many-body approach, and have shown that it leads to very sim ilar conductance oscillations. This suggests that electronic correlations are an additional and independent reason for their occurrence. In the presence of re ection symmetry, this mechanism leads to perfect conductance at the maxima of the oscillations, for all llings. The length-dependent oscillations persist in the presence of asym metric contacts between the chain and the leads. The period of the oscillations is given by the inverse of the conductance band lling, independently of the model and the mechanism leading to the oscillations. But the phase of the oscillations depends on details of the model and whether or not interactions are present. In non-interacting models, the electrons are mainly scattered in the contacts, and the conductance oscillations are periodic. Electronic correlations lead to the occurrence of scattering processes in the whole chain and result in a decrease of the conductance with the length, in addition to the oscillations. This is a striking di erence between our correlated chain and non-interacting models. As a consequence, the amplitude of the oscillations depends on the length, and can even increase with the length of the chain when electronic correlations are present. These ndings should be relevant for distinguishing in the interpretation of recent and future experiments on the length-dependence of the conductance of atom ic chains and molecular wires the many body e ects com ing from electron correlations from the e ects describable by simpler one body theories.

W e thank G.-L. Ingold and R A. Jalabert for very helpful discussions and P. Schm itteckert for his DMRG program. RAM acknow ledges nancial support from the European Union through the Hum an Potential Program (contract HPRN-CT-2000-00144).

REFERENCES

- [1] H.Ohnishi, Y.Kondo, and K.Takayanagi, Nature 395, 780 (1998).
- [2] A J. Yanson et al., Nature 395, 783 (1998).
- [3] N.Agrat, AL.Yeyati, and JM. van Ruitenbeek, Physics Reports 377, 81 (2003).
- [4] R H M . Sm it et al, Nature 419, 906 (2002).
- [5] J.M. Luttinger, J.M ath. Phys. 4, 1154 (1963).
- [6] B.Gao et al., arX iv cond-m at/0311645.
- [7] R H M .Sm it et al, Phys. Rev. Lett. 91, 076805 (2003).
- [8] P.L. Pernas, A. Martin-Rodero, and F. Flores, Phys. Rev. B 41, 8553 (1990).
- [9] A.Oguri, Phys. Rev. B 59, 12240 (1999).
- [10] H.S.Sim, H.W. Lee, and K.J.Chang, Phys. Rev. Lett. 87, 096803 (2001).
- [11] P.Havu et al., Phys. Rev. B 66, 075401 (2002).
- [12] Z.Y. Zeng and F.C laro, Phys. Rev. B 65, 193405 (2002).
- [13] K S. Thygesen and K W .Jacobsen, Phys. Rev. Lett. 91, 146801 (2003).
- [14] R.A.Molina et al, Phys. Rev. B 67, 235306 (2003).
- [15] R.A.Molina et al., arX iv cond-m at/0402250.
- [16] J.Favand, F.M ila, Eur. Phys. J. B 2, 293 (1998).
- [17] O P.Sushkov, Phys. Rev.B 64, 155319 (2001).
- [18] P.Schmitteckert, Ph.D. thesis (Univ. Augsburg, 1996).
- [19] S.D atta, Electronic Transport in M esoscopic System s, C am bridge University P ress (1995).