Screening by composite charged particles: the case of quantum well trions

C. Ciuti and G. Bastard

Laboratoire Pierre Aigrain, Ecole Norm ale Superieure, 24, rue Lhom ond, 75005 Paris, France

(D ated: M arch 22, 2022)

W e study the screening of an external potential produced by a two-dimensional gas of charged excitons (trions). We determ ine the contribution to the dielectric function induced by these composite charged particles within a random phase approximation. In mixtures of free electrons and trions, the trion response is found dominant. In the long wave-length limit, trions behave as point charges with mass equal to the sum of the three particle components. For nite wave-vectors, we show how the dielectric response is sensitive to the composite nature of trions and the internal degrees of freedom. Predictions are presented for the screening of a C oulom b potential, the scattering by charged impurities and the properties of trionic plasm ons.

A major breakthrough in our understanding of the optical response of doped sem iconductor quantum wells (QW s) was achieved when the existence of trions was demonstrated experimentally both in II-VI and III-V sem iconductors [1, 2, 3]. These com posite particles appear as the weakly bound state of an exciton (a bound electron-hole pair) to an electron or a hole depending on the n-type or p-type doping of the QW. For the sake of clarity, in the following we will deal with negatively charged excitons (X). The physics of these composite charged particles in the low excitation regime has attracted a considerable interest, especially concerning the m etal-insulator transition [2, 4], the intrinsic radiative recombination e ciency [5, 6], the singlet-triplet crossover under strong magnetic eld [7], the role of phonons in the di usion properties [8], and the rem arkable drift transport induced by an applied electric bias[9, 10].

M ore recently, the st investigations of a dense trion gas have been performed by resonant optical pumpprobe[11]. In principle, by optical pum ping of the trion resonance, it should be possible to "convert" partially or even completely the background electron gas into a trion gas. Hence, it can be possible to study the many-body properties of a gas of com posite charged particles. One interesting issue to be addressed is the screening response of such a peculiar gas to an external potential, such as the Coulomb eld generated by a charged im purity. In this respect, the description of a trion gas as point particles (like electrons) should be only a long wave-length approach, while at shorter wave-length the trion granularity should show up. Since the impurity-induced scattering, the dc m obility and the collective excitations are sensitive to di erent wave-lengths, it is desirable to determ ine the complete response of a dense trion gas to external disturbances and to check whether and when the internal structure can be important. In this letter, we will present a Random Phase Approximation (RPA) treatm ent of the trion gas response. W e will show the role of the composite nature of trions in the determ ination of the dielectric response.

To simplify our description, we shall present results for a purely two-dimensional system, om itting the form factors due to the nite extension of the quantum wellwavefunctions of electrons and holes along the QW grow th direction. An X trion state is represented by a three-body wave-function, in which the center of mass, the relative m otion and the spin part can be factorized, namely

$$k_{; ;S_{e};S_{hz}} = \frac{e^{ik R}}{P} ; s_{e} (1; 2) s_{e}; s_{hz} (s_{1z}; s_{2z}; s_{hz});$$
(1)

where k and R are the center of m ass wave-vector and position respectively, A is the sample area, is the relative motion quantum number, while $S_{\rm e}$ and $S_{\rm hz}$ denote the spin state. The relative motion part depends on the variables $_1 = _1$ _h and $_2 = _2$ h, where 1, 2, h are the in-plane positions of the rst, second electron and the hole respectively, while s_{1z} , s_{2z} , s_{hz} are the spin components along the perpendicular direction z. In sem iconductor quantum wells, heavy-hole trions have eight spin states, where Se 2 f0;1g represents the total spin of the two electrons and S_{hz} 2 f 3=2g is the heavy-hole band angularm om entum projection. The two states with $S_e = 0$ are called singlet trions, while the six states with S_e = 1 are triplet trions. To ful 1 the Pauli exclusion principle, for a singlet (triplet) spin state, the ;se is symmetric (antirelative motion wave-function symmetric) under exchange of the positions of the two electrons. The X charge density operator reads

$$\hat{\mathbf{n}}(\mathbf{n}) = \mathbf{e}^{(2)}(\mathbf{n}_{1}) + \mathbf{n}^{(2)}(\mathbf{n}_{2})^{(2)}(\mathbf{n}_{h}) :$$
(2)

The matrix elements of \hat{n} () on the basis of the trion states are diagonal with respect to the spin indexes. They can be written as

$$hk^{0}; {}^{0}; S_{e}; S_{hz} \dot{f} \dot{f} \dot{f}; ; S_{e}; S_{hz} \dot{i} = \frac{e}{A} e^{i(k k^{0})} T_{k k^{0}}^{0}; ; S_{e}; ;$$
(3)

where the trion "granularity" factor T_q "; ;^{Se} accounting for the composite nature is

$$T_{q}^{\circ; ;S_{e}} = e_{;q}^{\circ; ;S_{e}} + e_{;q}^{\circ; ;S_{e}} + h_{;q}^{\circ; ;S_{e}}$$
(4)

where the electron contribution reads

$${}^{0}; ; {}^{S_{e}} = d^{2} {}_{1}d^{2} {}_{2} {}^{?}_{;S_{e}} ; {}^{S_{e}} e^{iq \frac{m_{e}}{M} 2} {}^{\frac{m_{e}+m_{h}}{M} 1};$$
(5)

and the hole part is

$$\int_{h_{q}}^{0} ; ; S_{e} = d^{2} d^{2} d^{2} 2 \int_{S_{e}}^{2} ; S_{e} e^{iq \left(\frac{m}{M}e\left(1+2\right)\right)}; (6)$$

with m_e , m_h , $M = 2m_e + m_h$ the electron, hole and trion masses respectively. If an external potential V_{ext} (;t) = $\Pr_{q \frac{1}{2}V_{ext}}(q;!)e^{i(q !t)} + hc$: acts on the system, then the trionic wave-functions are perturbed. The perturbation of the trion wave-function generates a non-hom ogeneous charge density, which creates a local Hartree potential and an exchange-correlation correction (we neglect it here). Hence, the screened potential is given by the self-consistent equation $V_s = V_{ext} + V_{loc}$, where V_{loc} (;t) = $d^2 \ 0 - \frac{e}{j} \ n (\ 0; t)$, being n (;t) the induced charge density and is the static dielectric constant of the sem iconductor quantum well. The self-consistent potential energy felt by a trion is

$$U_{s}^{tr}(_{1};_{2};_{3};t) = e(V_{s}(_{1};t) + V_{s}(_{2};t) V_{s}(_{h};t)):$$
(7)

The matrix elements of the perturbation energy on the trion states at t = 0 are

$$hk^{0}; {}^{0}; S_{e}; S_{hz} \mathcal{Y}_{s}^{tr} \mathcal{K}; ; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{hz} i = eV_{s}(k k^{0}; !) T_{k k^{0}}^{0}; S_{e}; S_{e};$$

By calculating the lowest-order perturbation theory for the trion wave-functions and sum m ing incoherently the contribution from all the populated trion states (RPA), we get the the screened potential $V_s(q;!) = V_{ext}(q;!) = _X$ (q;!), where the trion-induced dielectric function reads

$$_{X}$$
 (q;!) = 1 $\frac{2 e^{2}}{qA}$ $_{X}$ (q;!): (9)

The trion RPA -polarization contribution is

$$_{X} (q;!) = \frac{X}{E_{k+q; 0;S_{e};S_{hz}}} \frac{(f_{k+q; 0;S_{e};S_{hz}} f_{k;;S_{e};S_{hz}}) f_{q}; 0;S_{e} f_{k}}{E_{k+q; 0;S_{e}} E_{k;;S_{e}} h! i0^{+}};$$

where the sum is meant over k, , 0 , $S_e,\,S_{\rm hz}$. The quantity $f_k,\,;_{S_e},\!s_{\rm hz}$ is the trion occupation number (not necessarily at equilibrium) and E $_k,\,;_{S_e}$ the orbital energy of the unperturbed trion state. Note that for a (spin-unpolarized) electron gas

$${}_{e} (q;!) = 2 \binom{X}{k} \frac{f_{k+q}^{(e)} f_{k}^{(e)}}{\frac{h^{2}jk+qj^{2}}{2m_{e}} \frac{h^{2}k^{2}}{2m_{e}} h! i0^{+}}$$
(11)

and $_{\rm e}$ (q;!) = 1 $\frac{2 e^2}{qA} e^2$ (q;!). In presence of a mixed gas of trions and electrons, the linear susceptibilities of the two components add up, i.e. the polarization



FIG.1: Solid line: normalized static screening wave-vector $q_s(q;0)=q_s(0;0)$ versus dimensionless wave-vector q_a for a pure gas of trions. Dashed line: same quantity, but without accounting for the trion granularity. Stars: the granularity factor $f(q)f = f_e(q) = h(q)f$. Parameters: T = 10 K, a = 8 nm, $m_e = 0.1 \text{ m}_0$, $m_h = 0.2 \text{ m}_0$, $n_{trions} = 1 = 10^{10} \text{ cm}^2$, = 9.

of the mixture is $_{m ix} (q;!) = _{X} (q;!) + _{e} (q;!)$. Considering only the contribution of the ground trion state, the general expression in Eq. (10) can be considerably simplied. This approximation holds when only the ground state is populated and in the limit of small h!. In the case of zero magnetic eld, the ground state is a singlet 1s-like state, which is twice degenerate (due to the hole spin degree of freedom). By considering only the contribution of the 1s-state, we get

$$_{X} (q;!) ' 2 ^{X} (\underbrace{f_{k+q;1s} f_{k;1s}) \mathbf{T} (q)^{2}}_{k} : \frac{\frac{h^{2} \mathbf{k} + q \mathbf{f}}{2M} \frac{h^{2} \mathbf{k}^{2}}{2M} h! i0^{+}}{2M} : (12)$$

In the follow ing, we will consider the follow ing trial function

$$_{1s}(_{1};_{2}) = \frac{1}{2(a^{2})^{2}} \exp \left(\frac{1+a^{2}}{2a^{2}} \right); \quad (13)$$

where $a^{?}$ is the elective trion radius. A courate numerical solutions for the internal motion of quantum well trions are reported in the literature (see e.g. Refs. 5, 13). With our model wave-function, the granularity factor is T (q) = $2_{e}(q)_{h}(q)$ with

$$e(q) = \frac{1}{nh + \frac{m_e + m_h}{M} qa^2} \frac{1}{1 + \frac{m_e}{M} qa^2} \frac{1}{1 + \frac{m_e}{M} qa^2} (14)$$

and

:

$$_{h}(q) = \frac{1}{\pi h} \frac{1}{1 + \frac{m_{e}}{M} qa^{2}} (15)$$

In the long wave-length lim it (q! 0), jr (q) f! 1, i.e. the granularity factor does not play any role. Hence, for smallwave-vectorsq,X trionsbehave as point particles with charge e and m assM = $2m_e + m_h$, as expected. As shown in Fig. 1, the granularity factor decreases m onotonically with increasing q and then has a node for the wave-vector q such that $2_{e}(q) = h_{h}(q)$. This node of the granularity factor is due to the compensation between the contributions of two electrons and the hole within the bound X . This kind of cancellation e ect is absent in a plasm a of uncorrelated electrons and holes, where the screening contributions of the two components add up. Note that the value of qa? depends on the mass ratio $m_e = m_h$ and on the shape of the internal motion wavefunction (for our model wave-function qa? 1:1 when $m_e = m_h = 0.5$). Finally, in the short wave-length lim it $(q >> 1=a^{?})$, the granularity asymptotically vanishes, because the perturbing potential oscillates too quickly in the length scale of the trion internal motion wavefunction.

As a rst illustrative example of our theory, we consider the trion response to a Coulomb potential induced by a charged impurity, that is $V_{ext}(q) = 2 \text{ exp}(qd)=(Aq)$, where d is the distance between the remote impurity donor and the quantum well plane (d \notin 0 for a modulation-doped sample). The screened potential can be written as

$$V_{s}(q;0) = \frac{2 eexp(qd)}{A(q+q_{s}(q;0))};$$
(16)

being $q_s(q;0) = (m_{ix}(q;0))$ 1) q and m_{ix} the dielectric function for the mixed gas of trions and electrons. In Fig. 1, we show the relative screening parameter $q_s(q;0)=q_s(0;0)$ versus q (in units of the trion effective radius a?) for a pure trion gas, which can be obtained by resonant optical pum ping. For sim plicity, we have taken an equilibrium Ferm i-D irac distribution function with temperature T = 10K and with density $n_{tr} = 10^{10}$ cm². We have used typical parameters for a CdTe-based sem iconductor[1], namely $m_e = 0:1 m_{0}$, $m_h = 0.2 m_0$, = 9 (working in the CGS system) and a trion radius $a^{?} = 8 \text{ nm}$. The thick solid line represents $q_s(q;0)=q_s(0;0)$ due to the trion gas, while the dashed line shows the same quantity without including the granularity factor $\mathcal{T}(q)$ f. Indeed, the composite nature of trions has a major in pact in their screening response, dram atically quenching the response at nite wave-vectors.

O ne important issue to verify is the behavior of the static screening in presence of a mixed population of trions and electrons. In Fig. 2, we show a surface plot of $q_s(q;0)a^2$ as a function of the trion density fraction f, which is de ned by the relation $n_{tr} = f(n_e + n_{tr}) = fn_{tot}$. Passing from a pure electron gas (f = 0) to a pure trion gas (f = 1), the static screening changes considerably. For q ! 0, the screening wave-vector increases for increasing trion fraction due to the heavier



FIG.2: Contours of the screening wave-vector $q_{\rm b}(q;0)$ (units of 1=a[?]) as a function of qa and the trion density fraction f (f = 1 corresponds to a pure trion gas, f = 0 is for a pure electron gas). Parameters: total density (electrons + trions) $n_{\rm tot} = 10^{10}$ cm². O ther parameters as in Fig.1.

m ass of the trion. In fact, in the low tem perature lim it $q_s(0;0) / m_e$ for a pure 2D electron gas[12] and therefore $q_s(0;0) / (2m_e + m_h)$ for a pure 2D trion gas. Fig. 2 shows that even for sm all trion fractions, the screening of the m ixture is dom inated by the trion component. Hence, the trion granularity e ects at nite wave-vectors shown in Fig. 1 are important also in the m ixture case.

The scattering induced by charged in purities is the main interaction process a ecting the transport properties of charge carriers at low temperatures. W ithin the Ferm i's golden rule, the wave-vector dependent lifetime of trions $_{\rm tr}$ (k) in presence of the m ixed gas of trions and electrons is given by the expression

$$\frac{1}{tr(k)} = \frac{N_{imp}}{A} - \frac{2 M e^4}{2h^3} \int_{0}^{Z_2} d \frac{exp(2dq)}{(q_{mix}(q;0))^2}; \quad (17)$$

with q = 2k jsin (=2) j and $N_{imp} = A$ is the density of in purities per unit area. Notice that the electron lifetime is $e(k) = (M = m_e)_{tr}(k)$. The velocity lifetime $_{tr}^{v}$ (k) is given by same expression in Eq. (17), but with an additional factor (1 cos) within the integral. In Fig. 3, we plot the scattering integral $I_{scatt}(k) = R_0$ $\int_{0}^{\infty_{2}} d \exp(2dq) = (q_{m ix}(q; 0))^{2}$ (in units of a ²). In the case of a pure trion gas (f = 1, thick solid line), the k-dependence of the scattering integral has a resonant structure. The peak is due to the node of the granularity factor f(q)f (see Fig. 1). Indeed, around the nodal wave-vector, the trion screening is dram atically quenched and the scattering e ciency enhanced. This e ect is absent if the granularity of the trion is not taken into account (thick dashed line) and in the case of a pure electron gas (thin solid line). Note that in the case of a



FIG. 3: Impurity scattering rate integral I_{int} (units of a^{2}) as a function of qa. Thick solid line: pure trion gas (f = 1). Thick dashed line: the same, without granularity. Dotted line: mixture of trions and electrons (f = 0:3). This solid line: pure electron gas (f = 0). Distance between impurity and carrier planes: d = 4nm. O ther parameters as in Fig. 2.

m ixture oftrions and electrons, the contribution oftrions is dom inant, as shown by the dotted line, corresponding to a m ixture where the trion fraction is only 30%. The results for the velocity lifetim es (not shown) are qualitatively analogous.

Sim ilarly to the case of electrons, a pure gas of trions has longitudinal collective excitations (trionic plasmons), which are given by the solution of the equation X (q; ! (q)) = 0. In the long wave-length lim it, the trionic plasm ons behave exactly as the electronic plasm ons [12], once the electron m ass m $_{\rm e}$ is replaced by the trion m ass M = $2m_e + m_h$. For nite wave-vectors, as a result of the trion granularity, the plasm on frequency is decreased, due to the quenching of the screening e ciency. For a nite wave-vector q, when only the trion ground state is involved, the behavior is that of an electron gas, once the electron charge e is replaced by the e ective charge $e^{0}(q) = e_{T}(q) \leq e_{r}$, as it can be deduced from Eqs. (9) and (12). For excitation energies of the order of the trion binding energy, the excited states will give extra plasm on branches corresponding to transition between internal states. For a mixed gas of trions and electrons, the plasm on branches are the solutions of $m_{ix}(q; ! (q)) = X(q; ! (q)) + e(q; ! (q))$ 1 = 0. Hence, a coupling is present between the trionic and the electronic branches of plasm ons. The com plex phenom enology of the collective excitations in presence of a m ixture of electrons and trions will be addressed in detail in a future publication.

In conclusion, we have investigated the screening response of trions, taking into account their com posite nature. We have obtained a RPA-dielectric function for a m ixed gas of trions and electrons. This interesting physical regim e is experim entally achievable by resonant opticalpum ping of trions in doped QW s. In the static regim e, we have shown the major impact of the trion granularity in determ ining the dielectric response at nite wavevectors. W e have calculated the scattering rates of electrons and trions due to the interaction with the charged impurities. The internal motion of trions is responsible for a quenching of the screening response for wavelengths comparable or smaller than the trion e ective radius. M oreover, the granularity produces resonant features in the wave-vector dependence, due to the com pensation which can occur between the contribution of the two electrons and the hole within the X . We hope that our study will stimulate the research in the fundam ental properties of trions in the dense regime. We expect that the e ect of trion granularity in the screening response will have an impact on the transport properties of these charged particles, whose investigation is underway [9, 10]. In particular, the impurity-induced localization and the onset of the metal-insulator transition [4] should be considerably modi ed by the conversion of electrons into trions through resonant optical pumping.

LPA-ENS (form er LPMC-ENS) is "Unite Mixte de Recherche Associe au CNRS (UMR 8551) et aux Universites Paris 6 et 7".

- [1] K.Kheng et al, Phys.Rev.Lett. 71, 1752 (1993).
- [2] G.Finkelstein, H.Shtrikm an, I.Bar-Joseph, Phys. Rev. Lett. 74, 976 (1995).
- [3] A.J.Shields et al, Phys. Rev. B 51, R18049 (1995).
- [4] G.Eytan et al, Phys.Rev.Lett. 81, 1666 (1998).
- [5] A. Esser, E. Runge, R. Zimmermann, W. Langbein, Phys. Rev. B 62, 8232 (2000).
- [6] V.Ciulin et al, Phys. Rev. B 62, R16310 (2000).
- [7] G. Yusa, H. Shtrikman, I. Bar-Joseph, Phys. Rev. Lett. 87, 216402 (2001).
- [8] M. T. Portella-Oberli et al., Phys. Rev. B 66, 155305 (2002).
- [9] D. Sanvitto et al., Science 294, 837 (2001).
- [10] F.Pulizziet al, Phys.Rev.B 68, 205304 (2003).
- [11] M. T. Portella-Oberli et al., Phys. Stat. Sol. 238, 513 (2003).
- [12] T. Ando, A. Fow ler, F. Stern, Rev. M od. Phys. 54, 437 (1982) and references therein.
- [13] L C O. Dacal, JA. Brum, Phys. Rev. B 65, 115324 (2002).