

Particle density distributions in Fermi gas superfluids: Differences between one and two channel models

Jelena Stajic¹, Qijin Chen², and K. Levin¹

¹ *James Franck Institute and Department of Physics, University of Chicago, Chicago, Illinois 60637 and*

² *Department of Physics and Astronomy, Johns Hopkins University, Baltimore, Maryland 21218*

We show how to describe the $T \neq 0$ behavior associated with the usual BCS- Bose Einstein condensation (BEC) crossover ground state. We confine our attention here to the BEC and near-BEC regime where analytical calculations are possible. At finite T , non-condensed fermion pairs must be included, although they have been generally ignored in the literature. Within this BEC regime we compute the equations of state for the one and two channel models; these two cases correspond to whether Feshbach resonance effects are omitted or included. Differences between these two cases can be traced to differences between the nature of a Cooper pair and bosonic condensate. Our results are also compared with the Gross Pitaevskii equations of state for true bosons. Differences found here are associated with the underlying fermionic character of the system. Finally, the particle density distribution functions for a trap containing superfluid fermionic atoms are computed using a Thomas-Fermi approach. The one and two channel behavior is found to be very different; we find a narrowing of the density profile as a result of Feshbach resonance effects. Importantly, we infer that the ratio between bosonic and fermionic scattering lengths depends on the magnetic detuning and is generally smaller than 2. Future experiments will be required to determine to what extent this ratio varies with magnetic fields.

I. INTRODUCTION

The recent observations [1, 2, 3, 4, 5, 6, 7, 8] of Bose-Einstein condensation (BEC) of molecules formed from fermionic atoms are extremely exciting. Because of a Feshbach resonance, in which fermions couple to molecular bosons, it is possible [9, 10], via application of magnetic fields, to obtain an arbitrarily strong attraction between fermions, and to probe the crossover [11, 12] from BEC to BCS. The resultant superfluidity of pre-formed pairs has a natural counterpart in some theories [13, 14, 15, 16] of high T_c superconductors. Indeed, a striking, and important feature of the cuprate superconductors is their pronounced precursor superconductivity, or “pseudogap” effects, the origin of which is still under active debate.

A second, very important motivation for these experiments is based on the theoretical observation that a BCS-like ground state wavefunction is capable [17] of describing both fermionic and bosonic-based superconductors, provided that the chemical potential of the fermions, μ , is determined self consistently. Given the vast success of weak coupling or BCS theory, it is extremely important to formulate this extended theory at all T , and confront it with controlled experiments.

In this paper we explore the implications of this specific ground state wavefunction [17, 18] in detail to address both $T = 0$ and $T = T_c$, with particular emphasis on the “near-BEC” regime. Our goals are (i) to emphasize how to include finite temperature effects in a manner consistent with the ground state; this necessitates the introduction of non-condensed fermion pairs which lead to “pseudogap” effects [19], (ii) to provide analytic calculations and insights by working in a (near-BEC) regime where calculations are more tractable, (iii) to discuss in some detail the differences between the “one channel” and “two channel” models for the Feshbach resonance, and (iv) to compare with well established theories of weakly interacting Bose superfluid [20], as well as (v) with the measured density distributions in a trap. We begin

with the homogeneous case and then consider the trap configuration; from this we infer the ratio between the effective bosonic and fermionic scattering lengths, which is found to be strongly dependent on the magnetic detuning ν_0 , and generally less than 2, as a consequence of many-body combined with Feshbach resonance effects.

II. T MATRIX FORMALISM: BEYOND NOZIERES AND SCHMITT-RINK

The Hamiltonian used in the cold atom and high T_c crossover studies consists, in its most general form, of two types of interaction effects: those associated with the direct interaction between fermions parameterized by U , and those associated with “fermion-boson” interactions, whose strength is governed by g .

$$\begin{aligned} H - \mu N = & \sum_{\mathbf{k}, \sigma} (\epsilon_{\mathbf{k}} - \mu) a_{\mathbf{k}, \sigma}^\dagger a_{\mathbf{k}, \sigma} + \sum_{\mathbf{q}} (\epsilon_{\mathbf{q}}^{mb} + \nu - 2\mu) b_{\mathbf{q}}^\dagger b_{\mathbf{q}} \\ & + \sum_{\mathbf{q}, \mathbf{k}, \mathbf{k}'} U(\mathbf{k}, \mathbf{k}') a_{\mathbf{q}/2+\mathbf{k}, \uparrow}^\dagger a_{\mathbf{q}/2-\mathbf{k}, \downarrow}^\dagger a_{\mathbf{q}/2-\mathbf{k}', \downarrow} a_{\mathbf{q}/2+\mathbf{k}', \uparrow} \\ & + \sum_{\mathbf{q}, \mathbf{k}} \left(g(\mathbf{k}) b_{\mathbf{q}}^\dagger a_{\mathbf{q}/2-\mathbf{k}, \downarrow} a_{\mathbf{q}/2+\mathbf{k}, \uparrow} + h.c. \right) \end{aligned} \quad (1)$$

Here the fermion and boson kinetic energies are given by $\epsilon_{\mathbf{k}} \equiv \hbar^2 k^2 / 2m$ and $\epsilon_{\mathbf{q}}^{mb} \equiv \hbar^2 q^2 / 2M$, respectively, and ν is an important parameter which represents the “detuning”. The variational ground state which we will consider here is a product of both fermionic and bosonic contributions

$$\bar{\Psi}_0 = \Psi_0 \otimes \Psi_0^B \quad (2)$$

where the normalized fermionic wave function is the standard crossover state [17, 18]

$$\Psi_0 = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}}^\dagger c_{-\mathbf{k}}^\dagger) |0\rangle \quad (3)$$

and the normalized molecular or Feshbach boson contribution Ψ_0^B is represented by

$$\Psi_0^B = e^{-\lambda^2/2 + \lambda b_0^\dagger} |0\rangle. \quad (4)$$

The variational parameters are, thus, $u_{\mathbf{k}}$, $v_{\mathbf{k}}$ and λ . Applying standard variational techniques on the ground state wavefunction leads to Eqs. (5) and (40), and the $T = 0$ limit of Eq. (26). along with the result that $\lambda = \phi_m \equiv \langle b_{\mathbf{q}=0} \rangle$.

Whether both forms of interactions (U and g) in the Hamiltonian are needed in either system is still under debate. The bosons ($b_{\mathbf{k}}^\dagger$) of the cold atom problem [9, 10] will be referred to as Feshbach bosons. These represent a separate species, not to be confused with the fermion pair ($a_{\mathbf{k}}^\dagger a_{-\mathbf{k}}^\dagger$) operators. Thus we call this a “two channel” model.

For the sake of clarity we begin by ignoring the Feshbach resonance-induced interactions. The variationally induced constraints on the fermionic degrees of freedom [17] are given by

$$0 = 1 + U \sum_{\mathbf{k}} \frac{1}{2E_{\mathbf{k}}} \quad (5)$$

with the fermion density $n = 2 \sum_{\mathbf{k}} v_{\mathbf{k}}^2$, where $v_{\mathbf{k}}^2 \equiv [1 - (\epsilon_{\mathbf{k}} - \mu)/E_{\mathbf{k}}]/2$. We define the quasiparticle dispersion, $E_{\mathbf{k}}$, below.

One can anticipate that the excited states of this system are associated with their natural extension [15, 16, 19] following BCS theory. For $T \leq T_c$, the gap equation and number equations now involve Fermi functions $f(E_{\mathbf{k}})$ and are given by

$$\frac{m}{4\pi\hbar^2 a_s} = \sum_{\mathbf{k}} \left[\frac{1}{2\epsilon_{\mathbf{k}}} - \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} \right], \quad (6)$$

$$n = \sum_{\mathbf{k}} \left[1 - \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} + 2 \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} f(E_{\mathbf{k}}) \right], \quad (7)$$

where μ is now evaluated at finite T and $E_{\mathbf{k}} = \sqrt{(\epsilon_{\mathbf{k}} - \mu)^2 + \Delta^2}$. Here we assume a contact potential where a_s is the isolated two body scattering length, which is simply related to U , (see, for example, Eq. (A9)).

Two important observations need to be made. Here Δ is associated with the excitation gap for fermions. *This excitation gap reflects both condensed and non-condensed pairs and is not the same as the order parameter Δ_{sc} , except at $T = 0$.* Eqs. (6) and (7) are widely cited in the literature, but with an important difference. Throughout the literature Δ is taken to be the order parameter. By contrast, to determine Δ_{sc} here we will first need to characterize the contribution of non-condensed pairs to Δ .

Our second important observation is that for $T \leq T_c$, the Fermi factors in the above equations are essentially negligible [21]. It follows that *both $\Delta(T)$ and $\mu(T)$ are temperature independent in this near-BEC regime.* Indeed, this is consistent with the physical picture of well established, pre-formed pairs in the BEC limit, so that the fermionic energy scales are unaffected by T below T_c .

The simple physics in Eq. (6) may be schematically represented by plots of Δ versus temperature. Figure 1 contrasts




FIG. 1: Contrasting temperature dependences of Δ in the BCS and BEC regimes. Similarly, in the BEC regime μ is a constant, so that all fermionic energy scales are T independent, as expected.

the behavior in the weak coupling or BCS and strong coupling or BEC regimes. In the BCS limit $\Delta(T)$ follows the behavior of the order parameter, whereas in the BEC regime, pairs are pre-formed and there is no temperature dependence in $\Delta(T)$ on the scale of T_c . We now extend these qualitative observations to a more quantitative level.

A. Extending Conventional Crossover Theory to $T \neq 0$: BEC Limit Without Feshbach Bosons

The self consistent equations in the BEC limit for general temperature T can then be written as

$$\frac{m}{4\pi\hbar^2 a_s} = \sum_{\mathbf{k}} \left[\frac{1}{2\epsilon_{\mathbf{k}}} - \frac{1}{2E_{\mathbf{k}}} \right], \quad (8)$$

$$n = \sum_{\mathbf{k}} \left[1 - \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} \right], \quad T \leq T_c. \quad (9)$$

It follows from the above equations that, just as in the $T = 0$ limit [22, 23], we have for general $T \leq T_c$,

$$n_{pairs} = \frac{n}{2} = Z_0 \Delta^2 \quad (10)$$

where the coefficient of proportionality

$$Z_0 \approx \frac{m^2 a_s}{8\pi\hbar^4}. \quad (11)$$

We arrive at an important physical interpretation. Even though Δ or n_{pairs} is a constant in T , this constant must be the sum of two temperature dependent terms. Just as in the usual theory of BEC these two contributions correspond to condensed and non-condensed components

$$n_{pairs} = n_{pairs}^{condensed}(T) + n_{pairs}^{non-condensed}(T), \quad (12)$$

so that we may decompose the excitation gap into two contributions

$$\Delta^2 = \Delta_{sc}^2(T) + \Delta_{pg}^2(T), \quad (13)$$

where $\Delta_{sc}(T)$ corresponds to condensed and $\Delta_{pg}(T)$ to the non-condensed (or pseudo) gap component. Each of these are

proportional to the respective number of condensed and non-condensed pairs with proportionality constant Z_0 . Just as in BEC, at T_c ,

$$n_{pairs}^{non-condensed}(T_c) = \frac{n}{2} = \sum_{\mathbf{q}} b(\Omega_{\mathbf{q}}, T_c), \quad (14)$$

where $b(x)$ is the usual Bose-Einstein function and $\Omega_{\mathbf{q}}$ is the dispersion of the non-condensed pairs, which will be self consistently determined below. Thus

$$\Delta^2(T_c) = \Delta_{pg}^2(T_c) = Z_0^{-1} \sum b(\Omega_{\mathbf{q}}, T_c) = \frac{n}{2} Z_0^{-1}. \quad (15)$$

We may deduce directly from Eq. (15) that $\Delta_{pg}^2 = -\sum_Q t(Q)$, if we presume that below T_c , the non-condensed pairs have propagator

$$t(Q) = \frac{Z_0^{-1}}{\Omega - \Omega_{\mathbf{q}}}. \quad (16)$$

This lead to a key question: how can one deduce the contribution from *non-condensed* pairs? We now work backwards to infer the dispersion $\Omega_{\mathbf{q}}$ for these pairs. A fundamental requirement on non-condensed pairs in equilibrium with a Bose condensate is that their effective chemical potential satisfy

$$\mu_{pair}(T) = 0, \quad T \leq T_c. \quad (17)$$

Equation (6) can be shown to be consistent with Eq. (17) provided that the propagator for non-condensed pairs is given by

$$t(Q) = \frac{U}{1 + U\chi(Q)}, \quad (18)$$

where

$$\chi(Q) \equiv \sum_K G(K)G_0(Q - K), \quad (19)$$

and G represents the fermionic Green's function which has a self energy $\Sigma(K) = -\Delta^2 G_0(-K)$. Here G_0 is the bare propagator. The details of this analysis are presented in Appendix A.

Another important point should be noted. This pair propagator or T-matrix differs from that first introduced by Nozieres and Schmitt-Rink [24] because here there is one dressed and one bare Green's function. In the approach of Ref. 24 both are taken as bare Green's functions. By contrast there are other schemes in the literature [25, 26, 27] where both Green's functions are dressed. We end by noting that at small four-vector Q (and moderately strong coupling) we may expand Eq. (18) to obtain

$$t(Q) = \frac{Z_0^{-1}}{\Omega - \Omega_{\mathbf{q}} + \mu_{pair} + i\Gamma_Q}. \quad (20)$$

Now we can deduce directly from Eq. (18) that the dispersion of non-condensed pairs is of the form

$$\Omega_{\mathbf{q}} = \hbar^2 q^2 / 2M_0^*. \quad (21)$$

In summary, this quadratic dispersion can be derived from the pair susceptibility $\chi(Q)$. In turn, the particular form for $\chi(Q)$ shown in Eq. (19) is chosen in order to be consistent with Eqs. (6) and (7). In this sense the usual BEC constraint expressed by Eq. (17) is intimately connected to the BCS-like gap equation of Eq. (6). The details of this analysis are shown in Appendix A.

B. Differences Between One and Two Channel Models

We now extend this analysis to include Feshbach bosons [11, 12]. For this situation we can write down an equation [19] equivalent to Eq. (6) with the direct fermion interaction U replaced by $U_{eff} \equiv U + g^2/(2\mu - \nu)$ and effective scattering length $a_s \rightarrow a_s^* = mU^*/(4\pi\hbar^2)$. Here we define

$$U^* \equiv U_0 - \frac{g_0^2}{(\nu_0 - 2\mu)} \equiv \frac{4\pi\hbar^2 a_s^*}{m}, \quad (22)$$

where a_s^* is dependent on μ . We thus have

$$\Delta(T) = -U_{eff} \sum_{\mathbf{k}} \Delta(T) \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}}, \quad (23)$$

$$n = \sum_{\mathbf{k}} \left[1 - \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} + 2 \frac{\epsilon_{\mathbf{k}} - \mu}{E_{\mathbf{k}}} f(E_{\mathbf{k}}) \right], \quad (24)$$

where $E_{\mathbf{k}} = \sqrt{(\epsilon_{\mathbf{k}} - \mu)^2 + \Delta^2(T)}$, and again Δ is to be distinguished from the order parameter [19]. Alternatively one can rewrite Eq. (23) as

$$\frac{m}{4\pi\hbar^2 a_s^*} = \sum_{\mathbf{k}} \left[\frac{1}{2\epsilon_{\mathbf{k}}} - \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} \right]. \quad (25)$$

In Eq. (24) n represents the number of fermions, but the number equation for the total number of particles involves both condensed and uncondensed bosons as well

$$n + 2n_b + 2n_b^0 = n^{tot}, \quad (26)$$

where $n_b^0 = \phi_m^2$ is the number of molecular bosons in the condensate. The number of noncondensed molecular bosons is given by

$$n_b(T) = - \sum_{Q \neq 0} D(Q) \quad (27)$$

where the Bose propagator is

$$D(Q) \equiv \frac{1}{i\Omega_n - \epsilon_q^{mb} - \nu + 2\mu - \Sigma_B(Q)}. \quad (28)$$

and we choose the self energy [19]

$$\Sigma_B(Q) \equiv -g^2 \chi(Q) / [1 + U\chi(Q)] \quad (29)$$

to be consistent with the Hugenholtz-Pines condition that bosons in equilibrium with a condensate must necessarily have zero chemical potential. This is equivalent to

$$\mu_{boson}(T) = 0, \quad T \leq T_c, \quad (30)$$

where $\mu_{boson} = 2\mu - \nu + \Sigma_B(0)$. It follows after some simple algebra that Eq. (30) is then consistent with Eq. (23).

C. Parameter Choices and Renormalization Scheme

We have introduced quantities (U_0, g_0, ν_0) which characterize the two body scattering in vacuum and correspond to the scattering lengths a_s or a_s^* . How are these related [28] to the parameters (U, g, ν) which enter into the Hamiltonian? To begin we ignore Feshbach effects. The way in which the two body interaction U enters to characterize the scattering (in vacuum) is different from the way in which it enters to characterize the N-body processes leading to superfluidity. In each case, however, one uses a T-matrix formulation to sum an appropriately selected but infinite series of terms in U . For the two-body problem in vacuum, there is a measurable characteristic of the scattering, the scattering length, a_s ,

$$\frac{m}{4\pi\hbar^2 a_s} \equiv \frac{1}{U_0} \quad (31)$$

which is related to U via the Lippmann-Schwinger equation

$$\frac{m}{4\pi\hbar^2 a_s} \equiv \frac{1}{U} + \sum_{\mathbf{k}} \frac{1}{2\epsilon_{\mathbf{k}}} \quad (32)$$

In this way one may solve for the unknown U in terms of a_s or U_0 . The two-body T-matrix equation (Eq.(32)) can be rewritten as

$$U = \Gamma U_0, \quad \Gamma = \left(1 + \frac{U_0}{U_c}\right)^{-1} \quad (33)$$

where we define the quantity U_c as

$$\frac{1}{U_c} = - \sum_{\mathbf{k}} \frac{1}{2\epsilon_{\mathbf{k}}} \quad (34)$$

Here U_c is the critical value of the potential associated with the binding of a two particle state in vacuum. Specific evaluation of U_c requires that there be a cut-off imposed on the above summation, associated with the range of the potential.

Now, for the Feshbach problem, the calculation proceeds in a similar fashion and one finds [28]

$$U = \Gamma U_0, \quad g = \Gamma g_0, \quad \nu - \nu_0 = -\Gamma \frac{g_0^2}{U_c}, \quad (35)$$

Here we define the parameter ν_0 which is directly related to the difference in the applied magnetic field B and B_0

$$\nu_0 = (B - B_0)\Delta\mu^0, \quad (36)$$

where $\Delta\mu^0$ is the difference in the magnetic moment of the two paired hyperfine states. To connect the various energy scales, which appear in the problem, typically 1Gauss $\approx 60E_F$ for ^{40}K .

D. Characterizing the Condensate

There is an important difference in the nature of the condensate for the two cases, with and without Feshbach bosons.

In the latter case there are two components to the condensate associated with Cooper pairs (Δ_{sc}) and condensed molecular bosons. Moreover, the Cooper condensate enters only into the gap equation, whereas the molecular Bose condensate enters also into the number equation. As long as the bosonic condensate n_b^0 is non-negligible, this difference leads to essentially different physics between the one and two channel problem, as will be demonstrated below.

In this section we enumerate the different relationships between the Cooper condensate and bosonic condensate terms. The order parameter associated with Eq. (1) represents a linear combination of both paired fermions (Cooper condensate) and condensed molecules. It is given by [11, 12]

$$\tilde{\Delta}_{sc} = \Delta_{sc} - g\phi_m \quad (37)$$

where, the boson order parameter $\phi_m = \langle b_{\mathbf{q}=0} \rangle$. We have

$$\Delta_{sc} = -U \sum_{\mathbf{k}} \langle a_{-\mathbf{k}\downarrow} a_{\mathbf{k}\uparrow} \rangle, \quad (38)$$

therefore, $\tilde{\Delta}_{sc}$ can be written as

$$\tilde{\Delta}_{sc} = - \left[U + \frac{g^2}{(2\mu - \nu)} \right] \sum_{\mathbf{k}} \langle a_{-\mathbf{k}\downarrow} a_{\mathbf{k}\uparrow} \rangle \quad (39)$$

where we have used the fact [28] that

$$\phi_m = \frac{g\Delta_{sc}}{(\nu - 2\mu)U}. \quad (40)$$

The number of condensed Feshbach bosons which enters the number equation [Eq. (26)] is given by $n_b^0 = \phi_m^2$. Thus we have

$$n_b^0 = \frac{g^2 \Delta_{sc}^2}{[(\nu - 2\mu)U]^2} = \frac{g^2 \tilde{\Delta}_{sc}^2}{[(2\mu - \nu)U + g^2]^2} \quad (41)$$

Using Eqs. (35) we can also write

$$n_b^0 = \left(1 - \frac{U_0}{U^*}\right)^2 \frac{\tilde{\Delta}_{sc}^2}{g_0^2}, \quad (42)$$

where U^* is defined in Eq. (22).

III. EQUATIONS OF STATE AT $T = 0$

A. One Channel Model

We now rewrite our central equations (6), (7) and (15) in the near-BEC limit to compare more directly with the case of a weakly interacting Bose gas, described by Gross-Pitaevskii (GP) theory. It can be shown that (in the absence of FB)

$$n = \Delta^2 \frac{m^2}{4\pi\sqrt{2m|\mu|}\hbar^3}, \quad (43)$$

which, in conjunction with the expansion of Eq. (6),

$$\frac{m}{4\pi\hbar^2 a_s} = \left(\frac{2m}{\hbar^2}\right)^{3/2} \frac{\sqrt{|\mu|}}{8\pi} \left[1 + \frac{1}{16} \frac{\Delta^2}{\mu^2}\right], \quad (44)$$

yields

$$\mu = -\frac{\hbar^2}{2ma_s^2} + \frac{a_s\pi n\hbar^2}{m}. \quad (45)$$

These equations hold at all $T \leq T_c$. At $T = 0$, these equations have been shown [22, 23] to be equivalent to the results of GP theory where one identifies an effective inter-pair scattering length $a_B = 2a_s$ via $n_B = \frac{\mu_B}{4\pi a_B \hbar^2 / M_B}$. Here $n_B = n/2$ represents the number density of pairs, $\mu_B = 2\mu + \hbar^2 / ma_s^2$ is the “bare” chemical potential of the pairs, and $M_B \approx 2m$ the pair mass.

B. Effects of Feshbach bosons

We now show that in the presence of Feshbach bosons this equation of state is no longer that of GP theory and, moreover, there are important differences in the ratio of the bosonic to fermionic scattering lengths. As a result of the Bose condensate n_b^0 in Eq. (26) one finds an extra term in the number equation

$$n^{tot} = \Delta^2 \left[\frac{m^2}{4\pi\sqrt{2m|\mu|}\hbar^3} + 2\frac{(1-U_0/U^*)^2}{g_0^2} \right]. \quad (46)$$

Combining the gap and number equation yields

$$\begin{aligned} \frac{m}{4\pi\hbar^2 a_s^*} &= \left(\frac{2m}{\hbar^2} \right)^{\frac{3}{2}} \frac{\sqrt{|\mu|}}{8\pi} \\ &\times \left[1 + \frac{1}{16\mu^2} \frac{n^{tot}}{\frac{1}{16\pi\sqrt{|\mu|}} \left(\frac{2m}{\hbar^2} \right)^{\frac{3}{2}} + 2\frac{(1-U_0/U^*)^2}{g_0^2}} \right]. \end{aligned} \quad (47)$$

Solving for μ in terms of a_s^* one finds a new equation of state

$$\mu \approx -\frac{\hbar^2}{2ma_s^{*2}} + \frac{2\pi^2}{m} \frac{g_0^2 \hbar^2}{U_0^2} n^{tot} a_s^{*4} \quad (48)$$

to lowest order in a_s^* . The second term in the above equation derives from the Bose condensate term. The first term in Eq. (46) contributes a term of order a_s^{*7} to this correction [29]. This behavior should be contrasted with the situation when FB are absent, where $a_B = 2a_s$. It should also be emphasized that the fermionic scattering length in a model without FB is an independent experimental parameter, while here a_s^* depends on μ , and must be obtained self-consistently.

IV. CALCULATIONS AT T_c

We turn next to a calculation of T_c , which requires that we determine Ω_q [via the T-matrix of Eq. (18)] as a function of the scattering length a_s . We address the one-channel case first. The general expression for $1/M_0^*$ in the near BEC limit is given by

$$\frac{1}{M_0^*} = \frac{1}{Z_0 \Delta^2} \sum_{\mathbf{k}} \left[\frac{1}{m} v_{\mathbf{k}}^2 - \frac{4E_{\mathbf{k}} \hbar^2 k^2}{3m^2 \Delta^2} v_{\mathbf{k}}^4 \right], \quad (49)$$

where we have used Eqs. (21) and (20). After expanding to lowest order in na_s^3

$$M_0^* \approx 2m \left(1 + \frac{\pi a_s^3 n}{2} \right). \quad (50)$$

From Eq. (15) we then conclude that, at T_c ,

$$\frac{n}{2} = \sum_{\mathbf{q}} b(\Omega_{\mathbf{q}}, T_c). \quad (51)$$

Equation (51) reflects the fact that, in the near-BEC limit, and at T_c , all fermions are constituents of uncondensed pairs. From the above equation it follows that $(M_0^* T_c)^{3/2} \propto n = \text{const.}$ which, in conjunction with Eq. (50) implies

$$\frac{T_c - T_c^0}{T_c^0} = -\frac{\pi a_s^3 n}{2}. \quad (52)$$

Here T_c^0 is the transition temperature of the ideal Bose gas with $M_0 = 2m$. This downward shift of T_c follows the effective mass renormalization, much as expected in a Hartree treatment of GP theory at T_c . Here, however, in contrast to GP theory for a homogeneous system with a contact potential [20], there is a non-vanishing renormalization of the effective mass.

We turn now to the analysis of the behavior of T_c in the presence of Feshbach bosons. The (inverse) residue in the T -matrix is replaced by

$$Z_1 = Z_0 + \frac{g^2}{[U(2\mu - \nu) + g^2]^2} \approx \frac{n}{2\Delta^2} \quad (53)$$

and the $\hbar^2 q^2$ coefficient $B_1 \equiv 1/(2M_1^*)$ in $\Omega_{\mathbf{q}}$ is such that

$$B_1 = \frac{B_0 Z_0 + \frac{1}{2M} \frac{g^2}{[U(2\mu - \nu) + g^2]^2}}{Z_1} \approx \frac{1}{4m} \left[1 - \frac{2g_0^4}{U_0^4} \pi^3 n a_s^{*9} \right]. \quad (54)$$

Here Z_0 and B_0 are the appropriate counterparts when FB are absent. Since Z_0 is proportional to the fermionic contribution to the density, it is very small in the BEC limit. Using the same reasoning as in the previous case, we conclude that the ratio T_c/B_1 is constant with varying coupling. *Thus, again, T_c follows the behavior of the inverse effective mass* with, to leading order, $(T_c^0 - T_c)/T_c^0 \propto a_s^{*9}$.

V. PARTICLE DENSITY PROFILES IN TRAPS

The differences between the equations of state for the one and two channel models will have physical implications in the density profiles of particles in a trap. We now introduce the harmonic trapping potential $V(r) = \frac{1}{2}m\omega^2 r^2$, which is treated in the Thomas-Fermi (TF) approximation. In this approximation, one replaces μ with $\mu(r) = \mu - V(r)$. In contrast to the uniform case, here $\mu_{pair}(r, T)$ becomes non-zero beyond a critical radius $R_c(T)$, where $R_c(T_c) = 0$. In this way at T_c , only the center of the trap is superfluid, while at




FIG. 2: Density profiles at (a) $\nu_0 = -1260$ and (b) $\nu_0 = -250$. Solid and dashed lines are at $T = 0$ with and without Feshbach bosons (FB), respectively, and the dotted lines are at $T = T_c$. Inset to (a) plots the FB weight in the condensate. Inset to (b) shows the ratio of the inter-boson to inter-fermion scattering lengths at trap center. Units are chosen such that $k_F \equiv 1$, and $E_F = (3N)^{1/3} \hbar\omega = 1$. Hence the units for U_0 , E_F/k_F^3 , and for g_0 , $E_F/k_F^{3/2}$, are both set to unity. For $\nu_0 = -250$ the interaction parameter $k_F a_s^* \approx 0.22$.

$T = 0$ all of the trap contains condensed states. To obtain the $T = 0$ density profile, $n(r)$, we insert $\mu(r)$ into Eq. (47) and solve for $n(r)$ [Here $n(r)$ refers to the sum of both fermion and molecular boson contributions]. The solution is

$$n(r) = 16\mu^2(r) \left[\frac{\hbar}{\sqrt{2m}|\mu(r)|a_s^*} - 1 \right] \times \left[\frac{1}{16\pi\sqrt{|\mu(r)|}} \left(\frac{2m}{\hbar^2} \right)^{3/2} + 2 \frac{(1 - \frac{U_0}{U^*})^2}{g_0^2} \right], \quad (55)$$

where we use $N = \int n(r) d^3r$ to self-consistently determine μ . Here it should be noted that a_s^* is itself a function of $\mu(r)$.

In Fig. 2(a), we plot $n(r)$ for the parameters $U_0 = -0.89$, $g_0 = -35$, $\nu_0 = -1260$ and $N = 10^5$. For this value of ν_0 we are somewhat away from the deepest BEC regime as is necessary to ensure the validity of the TF approximation and the fermionic contribution to the density is no longer negligible compared to its bosonic counterpart. Indeed, the percentage weight of the molecular boson condensate (shown in the inset) indicates that by $\nu_0 \approx -400$ the Cooper pair condensate is beginning to dominate. In this figure we also show the density profile as computed in the absence of FB (dashed line), as well as the behavior at $T = T_c$ (dotted line). Our $T = T_c$ curves were computed in the absence of FB for both panels. We were unable, thus far to find important differences between this distribution and that of an ideal Bose gas. Here one has to solve self consistently for $\mu_{pair}(r)$, as well.

To arrive at a meaningful comparison of the two $T = 0$ cases (with and without Feshbach bosons) we used the same value for the effective two-body scattering length a_s^* , obtained from the self-consistent calculations of $n(r)$ in the presence of FB. The profile without FB bosons is then calculated using the familiar TF result near the BEC limit [23] at a fixed fermionic scattering [30] length a_s^* . The same plots are presented with $\nu_0 = -250$ in Fig. 2(b), which is further from the BEC limit.

An important consequence of the one vs two channel problems is that for the latter, as the extreme BEC limit is approached, the number of fermions is diminished in favor of FB. This occurs due to the self consistent adjustment of the fermionic chemical potential. Thus the interaction between the bosons mediated by the fermions is weaker than what might have been expected without consideration of FB. This weakened interaction is reflected in Fig. 2 through a comparison between the solid and dashed lines, which shows that the trap profiles are narrower in the presence of Feshbach effects. Indeed, this could have been anticipated from the above calculations in the homogeneous case, at $T = 0$ and T_c , which deduced a very weak dependence of the bosonic scattering length and effective pair mass on the fermionic scattering length a_s^* . This finding mirrors the result in a trapped atomic Bose gas [20] where one sees that a weaker repulsive inter-boson interaction leads to a narrowed density profile. Comparison between the lower and upper panels of Fig. 2 shows that in both cases (with and without FB), the profiles become narrower as the BEC limit is approached.

We may use the results in Fig. 2 to obtain a semi-quantitative estimate of the bosonic scattering length a_B , based on a phenomenology used in experimental analysis [1, 2, 3]. We compare our results to the Thomas-Fermi approximated GP equation at $T = 0$ which yields $n_B(r) = \frac{\mu_B - V(r)}{U_B}$ where the inter-boson interaction U_B is connected to the scattering length a_B via $U_B = \frac{4\pi\hbar^2 a_B}{M_B}$. If we fit the profiles of Fig. 2 to the inverted parabola $(C_1 - C_2 r^2)$, we may infer that $a_B = \frac{M_B^2 \omega^2}{4\pi\hbar^2 C_2}$. The ratio a_B/a_s^* is plotted vs ν_0 in the inset to the lower panel of Fig. 2b. The same analysis applied to the profile without FB yields the familiar 2:1 ratio of the bosonic to fermionic scattering lengths. The parameter a_B is an important quantity which appears in experiment to be considerably less than a factor of 2 times its fermionic counterpart.

VI. CONCLUSIONS

There are very detailed calculations [31] of four-body atomic scattering processes which yield a fixed length ratio of 0.6 for the bosonic to fermionic scattering lengths. Presumably in the asymptotic BEC regime this constraint should be imposed on a more exact theory of the ground state. Indeed, a weakness of the mean field approach taken here (based on a generalized BCS ground state) is that the “inter-boson” scattering length is treated only approximately. This can be viewed as related to the inclusion of only one fermion and two fermion propagators (*i.e.*, T-matrix) with no higher order

terms.

A strength of this approach, over the counterpart atomic calculations is that many body physics associated with the broken symmetry or superfluid state is included. *A comparison of the one and two channel calculations presented here shows that, through many body effects, the nature of the condensate enters in an important way to determine the equation of state and, thereby, the ratio of the scattering lengths.* It should also be stressed that the many body context with which the number 0.6 is to be compared is different from 2.0, which is the number generally assumed in the literature. This difference appears once one includes Feshbach bosons. Indeed, while it depends upon the magnetic field or detuning, in the near-BEC regime this ratio is found to be significantly less than 2.0. This is a central point of the present paper. Future experiments will be required to determine, whether, as some have presumed, the ratio of 0.6 applies to all detunings, or whether this number is variable as argued here.

In this paper we have shown that superfluidity of fermionic atoms in the near-BEC limit is in general different from Bose superfluidity (as described by GP theory). We have compared one and two channel models and find that differences from the GP picture for each are associated with the underlying fermionic character of the system. This shows up at $T \neq 0$ for the one channel system and at all T for a Hamiltonian which includes Feshbach bosons.

Essential to our approach is the presence (at all $T \neq 0$) of non-condensed pairs which contribute separately to the fermionic gap parameter Δ , in addition to the condensate term. Noncondensed pairs or “pseudogap effects” have been generally ignored in the literature [24] and we have shown elsewhere [19] that they are negligible only in the strict BCS regime. Our inclusion of these effects corresponds [32] to a self-consistent Hartree treatment of pairing fluctuations in BCS theory, in which condensation naturally occurs in the presence of a finite excitation gap (“pre-formed pairs”). Our approach also makes evident the strong analogies between this generalization of BCS theory and Bose condensation, where both condensed and non-condensed bosons must be properly characterized. *In the two channel model, non-condensed bosons should be viewed as consisting of a strongly hybridized mixture of fermion pairs and Feshbach bosons. In the one channel model they result exclusively from fermion pairs.*

A key aspect of this work is in the comparison we have presented between one and two channel models. Differences arise because of the nature of the condensate. A fermionic or Cooper pair condensate (Δ_{sc}) enters into the gap equation while a bosonic condensate (n_b^0) enters also into the number equation. When there is an appreciable fraction of bosonic condensate (as in the near-BEC and BEC regimes) these differences will be apparent. A distinction between the one and two channel models is relatively unimportant once the bosonic condensate is negligible (usually when the scattering length is large and positive or alternatively negative). These differences are associated with physical properties such as the equations of state and the density profiles in a trap. Feshbach bosons can, in effect, collapse much more completely to the center of a trap than can fermion pairs. This effect can be inferred from

the ratio of the scattering lengths, and is related to the fact that in this limit, fermions are essentially absent, since the number equation constraint can be entirely satisfied by populating the bosonic state. As a result, these bosons are close to ideal.

For the relatively broad Feshbach resonances currently under study in lithium and potassium, differences between the one and two channel models are presumably only important in the BEC and near-BEC limits studied here, where there is an appreciable bosonic condensate. It will be interesting to study narrower Feshbach resonances where these differences may persist into the unitary scattering regime.

Acknowledgments

We acknowledge useful discussions with R. Hulet, D. Jin and Z. Tesaonovic. This work was supported by NSF-MRSEC Grant No. DMR-0213765 (JS and KL), NSF Grant No. DMR0094981 and JHU-TIPAC (QC).

APPENDIX A: CONNECTION BETWEEN $\chi(Q)$ AND THE STANDARD GAP EQUATION

The expansion of the T-matrix at small four-vector $Q = (\Omega, \mathbf{q})$ can be written in the following form:

$$t^{-1}(Q) = Z_0(\Omega - \Omega_{\mathbf{q}} + \mu_{pair} + i\Gamma_{\mathbf{q}}) \quad (\text{A1})$$

Therefore, the condition for the divergence of the T-matrix at zero Q is equivalent to

$$\mu_{pair} = 0 \quad (\text{A2})$$

or

$$U^{-1} + \chi(0) = 0. \quad (\text{A3})$$

We now show that for a proper choice of the pair susceptibility

$$\chi(Q) = \sum_K G(K)G_0(Q - K) \quad (\text{A4})$$

Eq. (A2) is equivalent to the BCS-like gap equation of Eq. (6), provided also the Greens function is of the BCS form

$$G(K) = \frac{u_{\mathbf{k}}^2}{i\omega_n - E_{\mathbf{k}}} + \frac{v_{\mathbf{k}}^2}{i\omega_n + E_{\mathbf{k}}} = \frac{i\omega_n + \xi_{\mathbf{k}}}{(i\omega_n)^2 - E_{\mathbf{k}}^2}. \quad (\text{A5})$$

Just as in BCS theory, we associate this dressed Green's function with self-energy

$$\Sigma(K) = -\Delta^2 G_0(-K) \quad (\text{A6})$$

We calculate $\chi(Q = (i\Omega_m, \mathbf{q}))$ by performing the appropriate Matsubara sums following standard procedure [33]. At $Q = 0$, we have

$$\begin{aligned} \chi(0) &\equiv \sum_K \frac{i\omega_n + \xi_{\mathbf{k}}}{(i\omega_n)^2 - E_{\mathbf{k}}^2} \frac{1}{-i\omega_n - \xi_{\mathbf{k}}} \\ &= \sum_{\mathbf{k}} \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} \end{aligned} \quad (\text{A7})$$

and Eq. (A3) becomes

$$U^{-1} + \sum_{\mathbf{k}} \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} = 0, \quad (\text{A8})$$

which, in conjunction with the two-body scattering equation

$$\frac{m}{4\pi\hbar^2 a_s} = U^{-1} + \sum_{\mathbf{k}} \frac{1}{2E_{\mathbf{k}}}, \quad (\text{A9})$$

gives

$$\frac{m}{4\pi\hbar^2 a_s} = \sum_{\mathbf{k}} \left[\frac{1}{2\epsilon_{\mathbf{k}}} - \frac{1 - 2f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} \right] \quad (\text{A10})$$

Thus, we have demonstrated that, if we presume Eqs. (A4) and (A6), then Eqs. (A2) and (A10) are equivalent.

-
- [1] M. Greiner, C. A. Regal, and D. S. Jin, *Nature* **426**, 537 (2003).
 - [2] S. Jochim *et al.*, *Science* **302**, 2101 (2003).
 - [3] M. W. Zwierlein *et al.*, *Phys. Rev. Lett.* **91**, 250401 (2003).
 - [4] T. Bourdel *et al.*, preprint, cond-mat/0403091 (unpublished).
 - [5] C. A. Regal, M. Greiner, and D. S. Jin, *Phys. Rev. Lett.* **92**, 040403 (2004).
 - [6] M. W. Zwierlein *et al.*, *Phys. Rev. Lett.* **92**, 120403 (2004).
 - [7] J. Kinast *et al.*, preprint, cond-mat/0403540 (unpublished).
 - [8] M. Bartenstein *et al.*, preprint, cond-mat/0403716 (unpublished).
 - [9] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiofalo, and R. Walser, *Phys. Rev. Lett.* **87**, 120406 (2001).
 - [10] E. Timmermans, K. Furuya, P. W. Milonni, and A. K. Kerman, *Phys. Lett. A* **285**, 228 (2001).
 - [11] Y. Ohashi and A. Griffin, *Phys. Rev. Lett.* **89**, 130402 (2002).
 - [12] J. N. Milstein, S. J. J. M. F. Kokkelmans, and M. J. Holland, *Phys. Rev. A* **66**, 043604 (2002).
 - [13] T. Kostyrko and J. Ranninger, *Phys. Rev. B* **54**, 13105 (1996).
 - [14] R. Micnas, J. Ranninger, and S. Robaszkiewicz, *Rev. Mod. Phys.* **62**, 113 (1990).
 - [15] Q. J. Chen, I. Kosztin, B. Jankó, and K. Levin, *Phys. Rev. Lett.* **81**, 4708 (1998).
 - [16] Q. J. Chen, K. Levin, and I. Kosztin, *Phys. Rev. B* **63**, 184519 (2001).
 - [17] A. J. Leggett, in *Modern Trends in the Theory of Condensed Matter* (Springer-Verlag, Berlin, 1980), pp. 13–27.
 - [18] D. M. Eagles, *Phys. Rev.* **186**, 456 (1969).
 - [19] J. Stajic *et al.*, preprint, cond-mat/0309329; accepted for publication in *Phys. Rev. A*. (unpublished).
 - [20] F. Dalfovo *et al.*, *Rev. Mod. Phys.* **71**, 463 (1999).
 - [21] These Fermi functions can be set to zero, since their arguments are of the order of $|\mu|/T_c \gg 1$.
 - [22] A. Perali, P. Pieri, and G. C. Strinati, *Phys. Rev. A* **68**, 031601R (2003).
 - [23] L. Viverit, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Phys. Rev. A* **69**, 013607 (2004).
 - [24] P. Nozières and S. Schmitt-Rink, *J. Low Temp. Phys.* **59**, 195 (1985).
 - [25] R. Haussmann, *Z. Phys. B* **91**, 291 (1993).
 - [26] O. Tchernyshyov, *Phys. Rev. B* **56**, 3372 (1997).
 - [27] Y. Yanase *et al.*, *Phys. Rep.* **387**, 1 (2003).
 - [28] S. J. J. M. F. Kokkelmans *et al.*, *Phys. Rev. A* **65**, 053617 (2002).
 - [29] If one expands the inverse T -matrix in the form of a GP equation, it can be seen that $a_B \propto \Delta^4 a_s^{*3} / n^2 \propto a_s^{*7}$.
 - [30] This scattering length should be distinguished from the isolated two-body scattering parameter which unlike a_s^* does not reflect the interactions responsible for superfluidity. Indeed this isolated two-body parameter appears to be inappropriate, in that it becomes negative deeper into the BEC regime ($\nu_0 \leq \frac{g_0^2}{U_0}$). This sign change should occur quite generally BCS-like behavior at large positive ν_0 .
 - [31] D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, preprint, cond-mat/0309010 (unpublished).
 - [32] J. Stajic, A. Iyengar, Q. J. Chen, and K. Levin, *Phys. Rev. B* **68**, 174517 (2003).
 - [33] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, San Francisco, 1971).