

Critical Ising Model and Financial Market

Takeshi Inagaki

Yamato Laboratory, IBM Japan

Shimotsuruma, Yamato-shi, Kanagawa, Japan

We investigate Ising model description of dynamics of stock price. The model is defined in near 2 dimensions, one dimension is time and another represents ensemble of stocks, and strength of response of investors to price change corresponds to inverse temperature of the system. At critical temperature, infinitely long correlation among number of trades along time is observed and power-law tail in distribution of price fluctuation appears.

1. INTRODUCTION

Emergence of power-law tail in variety of distribution function [1] in economic system gets much attention of physicists because it suggests existence of underlying critical phenomena. Especially, there exist amount of data that record stock trades over decades and fluctuation of stock price is one of most intensively studied topic. In this paper we study change of stock price

$$G(t) = \ln S(t+1) - \ln S(t) \quad (1.1)$$

where S is stock price. We try to explain observed power-law tail of the distribution function $P(G)$ with power-law exponent $-(g+1)$ where 1.5 for small g ($0.5 < g < 3$) and -3 for large g ($3 < g$) where g is price change at individual trade [2].

2. REVIEW OF FINDINGS IN EMPIRICAL STUDIES

Here is summary of finding in empirical studies [3,4,5,6] of statistical properties of financial time series in stock market. Denote number of trades in a certain interval t at time t by $N_t(t) = N_0 + N_t(t)$ where N_0 is average value of N_t over long term. Number of trades is strongly correlated along time as

$$\langle N_t(t+T) N_t(t) \rangle / T \quad (2.1)$$

where observed value of $\langle N_t(t+T) N_t(t) \rangle / T$ is around 0.3 . Another quantity measures activity in a market is share volume. Define amount of share volume of n trades

$$Q_n = \sum_{i=1}^n q_i \quad (2.2)$$

where q_i is share volume at i th transaction. It was found that each q_i is weakly correlated and distribution $P(Q_n)$ of Q_n is Levy stable distribution with a characteristic function

$$\langle k \rangle = e^{-m |k|^j} \quad (2.3)$$

where $j \approx 1.45$. Naively thinking, price changes of stocks are related to share volume $Q_{N_t(t)}$ share traded during t . More preciously, it is related to imbalance of share volume traded seller initiated and traded buyer initiated. That is defined by

$$X_t(t) = \sum_{i=1}^{N_t(t)} a_i q_i \quad (2.4)$$

where $a_i = \pm 1$ is sign for seller initiated (+1) and buyer initiated (-1) trade. The function form of ΔG varies by the time scale of interval t . But the function form of ΔG is stable where

$$\Delta G_t(t) = \sum_{i=1}^{N_t} a_i \quad (2:5)$$

(difference between number of buyer initiated and seller initiated trades in interval $[t, t+\Delta t]$) and it is approximated by

$$hG_t(t) / \tanh(A \Delta t) \quad (2:6)$$

where A is a constant. Distribution function $P(\Delta G)$ has a single peak around $\Delta G = 0$ if we pick trades with small local deviation of share volume

$$P(\Delta G) = hq_i a_i - hq_{-i} a_i i: \quad (2:7)$$

When ΔG exceeds a critical value ΔG_c , peak of $P(\Delta G)$ splits into two peaks with separation

$$\Delta G_c = \Delta G_c^+ - \Delta G_c^- \quad (2:8)$$

Because q_i is iid process, it is expected that this behavior is inherited to $N_t(t)$ as well.

3. CORRELATION IN TIME

Existence of correlation (2.1) implies existence of interaction between $N_t(t)$ and $N_t(t+1)$. This interaction might be mediated by $\Delta G_t(t) \Delta G_{t+1}(t)$. Because q_i is Levy stable process (2.3), we get

$$\Delta G_t(t) / N_t(t)^{1/\alpha} \quad (3:1)$$

and

$$\Delta G_t(t) / N_t(t) \sim N_c \quad (3:2)$$

From (2.6) and (3.2),

$$\begin{aligned} \Delta G_t(t) &= \text{zero} & (N_t(t) < n_0); \\ \Delta G_t(t) &\sim N_t(t)^{1/\alpha} & (n_0 < N_t(t) < n_1); \\ \Delta G_t(t) &= \text{constant}(\neq 0) & (n_1 < N_t(t)); \end{aligned} \quad (3:3)$$

Except for range $n_0 < N_t(t) < n_1$, $\Delta G_t(t)$ behaves as a bi-level stepwise function of $N_t(t)$. If we assume causal relation $\Delta G_t(t) \rightarrow N_t(t+1)$, probability distribution of $N_t(t+1)$ is

$$\begin{aligned} P_{t+1}(N_t) &= p_0(N_t) & (N_t(t) < n_0); \\ &= p_1(N_t) & (n_1 < N_t(t)); \end{aligned} \quad (3:4)$$

4. ISING MODEL

For convenience, introduce a new variable

$$x = \ln(N_t) - \ln\left(\frac{n_0 + n_1}{2}\right)$$

and assume symmetry between p_0 and p_1 of (3.4) as

$$p_0(x) = p_1(x) = p(x): \quad (4.1)$$

Represent N_t by bi-level of states $j = 1$ for $x < 0$ and $j = 0$ for $x > 0$. Transition amplitude among states is

$$\begin{aligned} P_{++} &= P_{11} e^+ = \frac{Z_1}{Z_0} \int_0^\infty dx p(x) \\ P_{+-} &= P_{10} e^- = \int_{-\infty}^0 dx p(x) \end{aligned} \quad (4.2)$$

where β represents strength of correlation in time series. If $\beta = 0$, there is no correlation among $N_t(t)$. This leads to Ising like model with spin variable $s_i = j = 1$ on 1 dimensional lattice (time with a tick t) and Hamiltonian is given by

$$H = - \sum_{(i,j)} s_i s_j: \quad (4.3)$$

When $\beta < \beta_c$ the system is in the disordered phase and trades happen randomly. If $\beta > \beta_c$ it is in the ordered phase and spins are aligned on a direction. In this case mean value of N_t is transiting and vacuum of the theory is unstable. If $\beta = \beta_c$ it is at a critical point and infinite length of correlation (power-law correlation) of spins (number of trade N_t) appears. Correlation function of magnetization in Ising model is given by

$$\langle s_{t+\tau} s_t \rangle = \tau^{-(d-2+\epsilon)} \quad (4.4)$$

where $\epsilon = \frac{1}{4}$ [7]. Regarding observed correlation of N_t , dimension of market Ising model is $d = 2.05$. One dimension is time and other may emerge from the network structure of interaction among ensemble of stocks. Equation (2.6) suggests the analogy of the mean field

$$hM_i / \frac{e^{+M} - e^{-M}}{e^{+M} + e^{-M}}: \quad (4.5)$$

It seems interaction in the space dimension (stock ensemble) is not restricted to its nearest neighborhood, rather is mediated by the mean field.

5. PRICE FLUCTUATION

From correlation (2.1), variation of sum of N_t asymptotically behaves as

$$\langle N_t^2 \rangle_T = \langle \sum_{i=1}^T N_t(i)^2 \rangle / T + T^2 \quad (5.1)$$

where α_0 and α_1 are constants of order $O(1)$. Around $T \gg 1$, this is approximated by

$$\langle N_t^2 \rangle_T \sim T^{1.35} \quad (5.2)$$

From (3.3), $\langle G \rangle$ is proportional to $\langle N_t^2 \rangle^{1/2}$ for small $\langle G \rangle$. For large $\langle G \rangle$, size of individual fluctuation does not depend on $\langle N_t^2 \rangle^{1/2}$ and it behaves as the Gaussian due to the Central Limit Theorem. This results

$$\begin{aligned} \langle G^2 \rangle^{1/2} &\sim \langle N_t^2 \rangle^{1/2} \sim T^{\frac{1.35}{2}} & (\langle G \rangle < 1) \\ \langle G^2 \rangle^{1/2} &\sim T^{\frac{1.35}{4}} & (1 < \langle G \rangle): \end{aligned} \quad (5.3)$$

This diffusion behavior of $\langle G^2 \rangle^{1/2}$ is consequence of distribution of G and we get observed power-law tail of the distribution function for G as

$$\begin{aligned} P(G) &\sim G^{-(\frac{2}{1.35} + 1)} \sim G^{-(1.48 + 1)} & (\langle G \rangle < 1) \\ &\sim G^{-(\frac{4}{1.35} + 1)} \sim G^{-(2.96 + 1)} & (1 < \langle G \rangle): \end{aligned} \quad (5.4)$$

6. CONCLUSION

In this paper, we see possible application of the Ising model to modeling of causal relation of activity in financial market. The structure of interaction is still obscure. Small world networks [8] that interpolates between the Ising model and the mean field theory is one of candidates.

References

- [1] B.B. Mandelbrot, J. Business 36, 394-419 (1963).
- [2] P. Gopikrishnan, V. Plerou, L.A.N. Amaral, M. Meyer, and H.E. Stanley, Phys. Rev. E 60, 5305-5316 (1999).
- [3] P. Gopikrishnan, V. Plerou, L.A.N. Amaral, Xavier Gabaix, and H.E. Stanley, Phys. Rev. E 62, 3023-3026 (2000).
- [4] P. Gopikrishnan, V. Plerou, Xavier Gabaix, and H.E. Stanley, Phys. Rev. E 62, 4493-4496 (2000).
- [5] P. Gopikrishnan, V. Plerou, Xavier Gabaix, and H.E. Stanley, Phys. Rev. E 66, 027104 (2002).
- [6] P. Gopikrishnan, V. Plerou, and H.E. Stanley, Nature 421, 130 (2003).
- [7] C. Itzykson and J-M. Drouot, Statistical field theory, Cambridge Univ. Press (1989).
- [8] D.J. Watts and S.H. Strogatz, Nature 393, 440 (1998).