

Calculation of Correlation Functions in Terms of Fluctuation Relation

Takaaki Monnai, Shuichi Tasaki

*Department of Applied Physics, Waseda University, 3-4-1 Okubo,
Shinjuku-ku, Tokyo 169-8555, Japan*

Abstract

We derive a relation similar to the fluctuation theorem for work done on a system obeying Langevin dynamics with thermal and colored noises. Then, we propose a method of calculating the correlation function of the colored noise by using this fluctuation relation.

Key words: fluctuation theorem, correlation function of colored noise, Langevin dynamics

PACS: 05.40.-a

1 Introduction

Fluctuation theorems (FT) [1] have been studied as one of exact relations valid in far from equilibrium regime. They state the symmetry of fluctuation of entropy production ΔS :

$$\left(\lim_{t \rightarrow \infty} \right) \frac{\pi_t(\Delta S = A)}{\pi_t(\Delta S = -A)} = A, \quad (1)$$

where $\pi_t(\Delta S = A)$ denotes the probability density that a system's entropy production during time interval t equals to A , and the limit of $t \rightarrow +\infty$ is taken or not depending on systems. Although interpretations and systems differ, this expression universally holds [7,8,9,10,11,12].

Jarzynski equality [2] was shown to be derived from FT [3] and was applied, e.g. to evaluate the free energy of biological systems [4]. It is even more inter-

Email address: monnai@suou.waseda.jp (Takaaki Monnai, Shuichi Tasaki).

esting to investigate direct applications of FT. In this letter, we derive a FT like relation (FR) for work done on a system obeying Langevin dynamics. In addition to the thermal noise, a colored noise is assumed to be applied. Then, this FR is shown to be used to evaluate a correlation function of the colored noise from the fluctuations of the work done. Such correlation functions would provide information about driving force in nonequilibrium regime and we expect that the FR-aided analysis is useful to investigate the behaviors of single molecular motors [5] or micro/nano machines.

2 Fluctuation Relation and Correlation Function

We consider a system described by a Langevin equation:

$$\int_{-\infty}^t \Gamma(t-t')v(t')dt' = f + \xi(t) + \Xi(t) \quad (2)$$

$$\langle \xi(t)\xi(t') \rangle = \frac{1}{\beta}\Gamma(t-t') \quad (3)$$

$$\langle \Xi(t)\Xi(t') \rangle = M(t-t') \quad (4)$$

$$\langle \xi(t) \rangle = \langle \Xi(t) \rangle = 0, \quad (5)$$

where $\xi(t)$ and $\Xi(t)$ are the thermal and colored noises, respectively, f stands for a constant external force, and β is the inverse temperature. The noises $\xi(t)$ and $\Xi(t)$ are assumed to be Gaussian and independent from each other. Eq.(3) is nothing but the fluctuation dissipation relation. We assume that the memory kernel $\Gamma(t)$ exponentially decays as $\Gamma(t) = \Gamma_0 e^{-\gamma|t|}$. Such a Langevin equation would provide a phenomenological description of the motion of molecular motors as pointed out by Harada [6], where the energy balance in the molecular motor was investigated.

Now, we introduce the work Σ_t done on the system by the external force f during the time interval t :

$$\Sigma_t \equiv \beta \int_0^t ds f v(s) \quad (6)$$

We denote the probability density of Σ_t being A as $\pi_t(\Sigma_t = A)$. As we shall show later, one then obtains a fluctuation relation (FR):

$$F(t) \equiv t \left(\left\{ \frac{1}{A} \log \frac{\pi_t(\Sigma_t = A)}{\pi_t(\Sigma_t = -A)} \right\}^{-1} - 1 \right)$$

$$= \frac{\beta}{2\gamma\Gamma_0} \int_0^t ds \int_0^t ds' \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega (\gamma^2 + \Gamma_0^2) \hat{M}(\omega) e^{i\omega(s-s')} , \quad (7)$$

where $\hat{M}(\omega)$ is the Fourier transformation of the correlation function $M(t)$ of the colored noise. Note that this FR is different from usual FT's, because it says nothing about entropy production, but only refers to the work done by an external force.

The FR (7) leads to a differential equation of $M(t)$:

$$-\frac{d^2}{dt^2}M(t) + \gamma^2 M(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega (\gamma^2 + \Gamma_0^2) \hat{M}(\omega) e^{i\omega t} = \frac{\gamma\Gamma_0}{\beta} \frac{d^2}{dt^2}F(t) . \quad (8)$$

The quantity $F(t)$ defined in (7) can be measured experimentally by observing individual trajectories and, thus, one can determine $M(t)$ provided two parameters γ and Γ_0 are known. We note that $F(t)$ is independent of A .

So far we have assumed $\langle \Xi(t) \rangle = 0$ for simplicity. However, the case of $\langle \Xi(t) \rangle \equiv \Xi_0 \neq 0$ can be treated in the same way and one can show that the function

$$\tilde{F}(t) \equiv t \left(\left\{ \frac{1}{A} \log \frac{\pi_t(\Sigma_t = A)}{\pi_t(\Sigma_t = -A)} \right\}^{-1} - \frac{f}{f + \langle \Xi_0 \rangle} \right)$$

instead of $F(t)$ satisfies

$$-\frac{d^2}{dt^2}M(t) + \gamma^2 M(t) = \frac{\gamma(f + \Xi_0)\Gamma_0}{\beta f} \frac{d^2}{dt^2}\tilde{F}(t) . \quad (9)$$

In this case again, the correlation function $M(t)$ of the colored noise can be evaluated experimentally.

3 Derivation of Fluctuation Relation

In this section, we shall derive (7). Because Σ_t is a Gaussian process, in order to determine $\pi_t(\Sigma_t = A)$, it is sufficient to calculate the mean value $\langle \Sigma_t \rangle$, and the variance $\langle \Sigma_t^2 \rangle - \langle \Sigma_t \rangle^2$.

With the aid of the Fourier transformation, one obtains following expression :

$$v(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{\hat{f}(\omega) + \hat{\xi}(\omega) + \hat{\Xi}(\omega)}{\hat{\Gamma}(\omega)} e^{i\omega t} , \quad (10)$$

where $\hat{\Gamma}(\omega) \equiv \frac{\Gamma_0}{\gamma + i\omega}$ is the Fourier-Laplace transformation of $\Gamma(t)$, and $\hat{f}(\omega) \equiv 2\pi f\delta(\omega)$, $\hat{\xi}(\omega)$ and $\hat{\Xi}(\omega)$ are the Fourier transformations of f , $\xi(t)$, and $\Xi(t)$, respectively. Therefore, (5) leads to

$$\langle \Sigma_t \rangle = \beta f^2 \frac{\gamma}{\Gamma_0} t \quad (11)$$

From $\langle \hat{\Xi}(\omega) \hat{\Xi}(\omega') \rangle = 2\pi \hat{M}(\omega) \delta(\omega + \omega')$ and the fluctuation dissipation theorem: $\langle \hat{\xi}(\omega) \hat{\xi}(\omega') \rangle = \frac{4\pi \Re \hat{\Gamma}(\omega) \delta(\omega + \omega')}{\beta}$, we obtain the variance of Σ_t :

$$\begin{aligned} \langle \Sigma_t^2 \rangle - \langle \Sigma_t \rangle^2 &= \beta \frac{2\gamma f^2}{\Gamma_0} t \\ &+ \beta^2 f^2 \int_0^t ds \int_0^t ds' \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} d\omega \int_{-\infty}^{\infty} d\omega' \int_{-\infty}^{\infty} d\omega'' \frac{\langle \hat{\Xi}(\omega) \hat{\Xi}(\omega') \rangle}{\hat{\Gamma}(\omega) \hat{\Gamma}(\omega')} e^{i\omega s} e^{i\omega' s'} \end{aligned} \quad (12)$$

$$= \beta \frac{2\gamma f^2}{\Gamma_0} t + \beta^2 f^2 \int_0^t ds \int_0^t ds' \frac{1}{2\pi} \int d\omega \frac{\gamma^2 + \omega^2}{\Gamma_0} \hat{M}(\omega) e^{i\omega(s-s')} . \quad (13)$$

Substituting the mean value $\langle \Sigma_t \rangle$ and the variance $\langle \Sigma_t^2 \rangle - \langle \Sigma_t \rangle^2$ into

$$\log \frac{\pi_t(\Sigma_t = A)}{\pi_t(\Sigma_t = -A)} = \frac{2\langle \Sigma_t \rangle A}{\langle \Sigma_t^2 \rangle - \langle \Sigma_t \rangle^2} , \quad (14)$$

one obtains the fluctuation relation for work done by an external force: (7).

To summarize, we derived a fluctuation relation for the work done on the system: eqs.(7) and (8). Eq.(7) has a form of the fluctuation theorem with modification due to $\langle \Xi(t) \Xi(t') \rangle$. An important point is that the right hand side of (8) is measurable experimentally, and thus, we can evaluate the correlation function $\langle \Xi(t) \Xi(t') \rangle$ of the nonthermal colored noise.

4 Acknowledgement

The authors are grateful to Professor P.Gaspard, Professor T.Yanagida, and Dr. M. Nishiyama for fruitful discussions. This work is supported by a Grant-in-Aid for Scientific Research (C) from JSPS, by a Grant-in-Aid for Scientific Research of Priority Areas ‘‘Control of Molecules in Intense Laser Fields’’ and the 21st Century COE Program (Holistic research and Education Center for Physics of Self-Organization Systems) both from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] D.J. Evans, E.G.D. Cohen and G.P. Morriss, (1993) Phys. Rev. Lett. **71**, 2401.
- [2] C. Jarzynski, (1996) Phys. Rev. Lett. **78** 2690; (1999) J. Stat. Phys. **95** 367.
- [3] G.E. Crooks (1999) Phys. Rev. E **60** 2721.
- [4] J. Liphardt, S. Dumont, S.B. Smith, I. Tinoco, and C. Bustamante, (2002) Science **296**(5574) 1832.
- [5] M. Nishiyama, H. Higuchi and T. Yanagida, (2002) Nature Cell Biol. **4** 790.
- [6] T. Harada, e-print cond-mat/0310547.
- [7] D.J. Evans and D. Searles (1994) Phys. Rev. E **50** 1645.
- [8] D.J. Evans and D. Searles, (2002) Adv. Phys. **51** 1529; E. Mittag and D.J. Evans, (2003) Phys. Rev. E **67** 026113.
- [9] J. Kurchan (1998) J. Phys. A **31** 3719.
- [10] J.L. Lebowitz and H. Spohn (1999) J. Stat. Phys. **95** 333.
- [11] G. Gallavotti and E.G.D. Cohen, (1995) Phys. Rev. Lett. **74** 2694; G. Gallavotti and E.G.D. Cohen, (1995) J. Stat. Phys. **80** 931; E.G.D. Cohen and G. Gallavotti, (1999) J. Stat. Phys. **96** 1343.
- [12] C. Jarzynski (2000) J. Stat. Phys. **98** 77.
- [13] G.M. Wang, E.M. Sevick, E. Mittag, D.J. Evans and D. Searles (2002) Phys. Rev. Lett. **89** 050601.
- [14] D.M. Carberry, J.C. Reid, G.M. Wang, E.M. Sevick, D.J. Searles, and D.J. Evans, to appear Phys. Rev. Lett. (2004).
- [15] R. van Zon and E.G.D. Cohen, (2003) Phys. Rev. E **67** 046102.