

Oscillations of magnetoresistance of 2D E G in a weak magnetic field under microwave irradiation

A. E. Patrakov and I. I. Lyapilin

Institute of Metal Physics, Ural Sibirsk Branch of RAS, Yekaterinburg, Russia

Under microwave irradiation, in 2D electron system with high filling factors oscillations of longitudinal magnetoresistance appear in the range of magnetic fields where ordinary SdH oscillations are suppressed. An unusual beat-like behaviour of these oscillations in weak magnetic fields ($B \approx 0.02$ T) was attributed to the zero spin splitting previously. We propose an alternate explanation of this beat-like structure on the basis of the Boltzmann kinetic equation, without any reference to the spin-orbit interaction.

The interest to theoretical investigations of non-linear transport phenomena in two-dimensional electron systems (2DES) has grown substantially due to new experimental results obtained in very clean 2DES samples. It has been discovered independently by two experimental groups [1, 2] that the resistance of two-dimensional high-mobility electron gas in GaAs-AlGaAs heterostructures reveals a series of new features in its dependence on magnetic field, temperature, radiation power, etc. Under microwave irradiation, in 2DES with high filling factors oscillations of longitudinal magnetoresistance appear in the range of magnetic fields where ordinary SdH oscillations are suppressed.

Among the new effects, an unusual behavior of longitudinal magnetoresistance in weak magnetic fields $B \approx 0.02$ T is worth mentioning. E.g., in experiments by Zudov [1] performed in GaAs-AlGaAs heterostructures (with electron mobility $\mu = 25 \cdot 10^6$ cm²/Vs and electron density $n_e = 3.5 \cdot 10^{11}$ cm⁻²) vanishing of amplitude of magnetoresistance oscillations has been observed at $B = 110$ T. Upon further lowering of the magnetic field, the amplitude of oscillations grows again (see the inset in Fig. 1). This feature of magnetoresistance has been interpreted as a beat. The analogous effect has been also observed by Mani [3] in GaAs-AlGaAs heterostructures (with electron mobility $\mu = 15 \cdot 10^6$ cm²/Vs and density $n_e = 3 \cdot 10^{11}$ cm⁻²). Vanishing of the amplitude of oscillations has been observed in the vicinity of $B = 240$ G (see Fig. 2). According to [3], the magnetic field at which the amplitude of magnetoresistance oscillations vanishes does not depend on microwave radiation frequency for the same sample, although the effect itself is visible only in the presence of radiation. The observed "beats" were associated by Mani with "zero-spin splitting" resulting from the spin-orbit interaction [4, 5].

In this paper we will show that the nature of the observed "beat" can be understood in a simple framework without the need for any spin-orbit interaction. First, let's mention important conditions of the experiments discussed above: the effect is seen at

$$\omega_c \approx \omega \approx \omega_c \approx \omega_c$$




FIG. 1: R_{xx} ($T = 1$ K) in the presence of microwave vs ω_c/ω (for clarity, plots have been offset vertically). The inset shows with magnification the part of the plot, corresponding to the frequency of 57 GHz and weak magnetic fields. At $B \approx 0.1$ kG the "beat" can be seen. (This figure was taken from [1])




FIG. 2: The resistance R_{xx} of 2DEG in the GaAs-AlGaAs heterostructure with and without microwave irradiation. The "beat" visible in resistance oscillations is expanded in the inset. (This figure was taken from [3])

It follows that the effect has a quasiclassical nature. Here τ is the momentum relaxation time, ω_c is the radiation and cyclotron frequencies respectively, ϵ_F denotes the Fermi energy, T is the temperature expressed in units of energy.

Since the "beat" expresses itself in the classical range of magnetic fields, we will start from the Boltzmann kinetic equation. Taking into account that the electrons




FIG. 3: Oscillations of 2DES photoconductivity in the magnetic field, $\mu = 10$

It follows from (14) that photon-assisted conductivity as a function of inverse magnetic field gives rise to slow oscillations: $\propto \cos(2\mu/\omega_c)$, whereas usual SdH oscillations denoted as $\sigma_{xx}^{ph; SdH}$ contain factors like $\cos(2\mu/\omega_c)$, are fast and their amplitude decreases rapidly when increasing temperature.

The dependency of photoconductivity upon μ/ω_c , calculated according to Eq. (14), is plotted on Fig. 3.

It also follows from Eq. (14) that at $\mu/\omega_c = 1$ the amplitude of radiation-induced magnetoresistance oscillations vanishes. This is the very fact responsible for the "beat" observed in experiments [1, 2]. Note that the observation of the node of the "beat" in stronger magnetic fields in experiments [2] as compared to [1] is (as one can easily verify) a direct consequence of the lower electron mobility in experiments [2].

-
- [1] M. A. Zudov, R. R. Du, L. N. Pfeiffer, K. W. West, Phys. Rev. Lett. **90**, 046807 (2003).
 [2] R. G. Mani, J. H. Simet, K. von Klitzing, V. Narayana-murti, Nature. **420**, 646 (2002).
 [3] R. G. Mani et. al, arXiv:cond-mat/0303034
 [4] Y. A. Bychkov and E. I. Rashba, J. Phys. C. **17**, 6039

- (1984).
 [5] G. Lommer, F. Malker, and U. Rossler, Phys. Rev. Lett. **60**, 728 (1988)
 [6] T. Ando, A. B. Fowler and F. Stem. Rev. Mod. Phys. **54**, 437 (1982).