

Kinetic theory of fluctuations in conducting systems

D. B. Gutman¹, A. D. Mirlin^{2,3} and Yuval Gefen⁴¹The James Franck Institute 5640 S. Ellis Avenue Chicago, IL 60637²Institut für Nanotechnologie, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany³Inst. für Theorie der Kondensierten Materie, Universität Karlsruhe, 76128 Karlsruhe, Germany⁴Department of Condensed Matter Physics, Weizmann Institute of Science 76100 Rehovot, Israel
(Dated: March 22, 2024)

We propose an effective field theory describing the time dependent fluctuations of electrons in conducting systems, generalizing the well known kinetic theory of fluctuations. We apply then the theory to analyze the effects of strong electron-electron and electron-phonon scattering on the statistics of current fluctuations. We find that if the electron-electron scattering length is much shorter than the transport mean free path the higher cumulants of current are parametrically enhanced.

Fluctuations of electric current in time have been the subject of fundamental and applied research since the time of Schottky. They have been a centerpiece in the study of systems ranging from vacuum tubes to quantum Hallbars (for a review see [1]). More recently noise has been recognized as an important diagnostic probe for the nature of disorder and interactions. Traditionally the number of electrons taking part in transport in macroscopic systems is large; for that reason current fluctuations obey nearly Gaussian statistics. Under such circumstances it is the power spectrum which is the object of primary interest when addressing the fluctuations. However, as the sample size shrinks down from a vacuum tube to a nano junction, and the currents involved become correspondingly smaller, the central limit theorem does not apply anymore, and information concerning higher current cumulants may be needed. Various techniques have been applied to address this issue [1, 2, 3, 4, 5, 6].

There is an interesting and relevant class of systems out of equilibrium in which the effect of quantum interference on transport quantities is negligible. Transport through such systems is often described by classical kinetic equations (e.g. the Boltzmann equation). The intensity of the noise spectrum can then be found employing a related approach, known as the kinetic theory of fluctuations (KTF) [7, 8, 9]. The appeal of this approach is its intuitive transparency, yet traditionally it was constructed to describe pair correlation functions, hence was limited to finding the second current cumulant. In a recent insightful work Nagaev [10] has proposed a generalization of the KTF, expressing higher order cumulants in terms of pair correlators. Using this idea (which was termed the "cascade approach"), Nagaev calculated the third and the fourth cumulants of the current. While this approach was shown to successfully address a number of problems, its (quantum-mechanical) microscopic foundations, as well as its formulation in terms of an effective field theory (which would allow for further generalization, such as including interactions or calculating the full counting statistics (FCS)) were not fully understood. An important step in the latter direction was an introduction of a stochastic path integral by Pilgram, Jordan, Sukhorukov, and Buttiker [11].

In the present paper we propose a phenomenological effective field theory describing the time dependent fluctuations in phase space of electrons in conducting systems.

This is a field-theoretical implementation of the Nagaev's regression scheme. This theory allows for various generalizations, including the effects of electron-phonon and electron-electron interactions. In a number of instances, this theory is equivalent (up to small quantum corrections) [12, 13] to the non-linear model formulation of the problem [6]. In particular, for non-interacting electrons it reproduces the FCS [14] of current fluctuations obtained earlier within quantum approaches. We apply the theory to systems with a strong electron-electron scattering ("hot electron regime"). We find that (in the absence of umklapp processes), the current statistics changes dramatically depending on the relation between the electron-electron scattering length l_{ee} and the elastic mean free path. Specially, in the regime $l_{ee} \ll l_{tr}$, when the electrons form a weakly disordered liquid, the current cumulants are parametrically enhanced due to the electron-electron scattering. We find the scaling law governing the dependence of the FCS of hot electrons on the ratio $l_{ee} = l_{tr}$. Finally, we show that under strong electron-phonon interaction the statistics is Gaussian.

The effective action. The KTF can be cast as a Boltzmann equation with a Langevin-type noise [9, 15]

$$\hat{L} + \hat{\Gamma} f(p; x; t) = J(p; x; t); \quad (1)$$

Here we have introduced the Liouville operator

$$\hat{L} = \frac{\partial}{\partial t} + \frac{p}{m} \frac{\partial}{\partial x} + eE \frac{\partial}{\partial p}; \quad (2)$$

while $f(p; x; t) dx dp$ is the number of electrons in a unit cell centered at $p(p; x)$ in a phase space at time t . With $\langle x; t \rangle = e \int p f(p; x; t)$ being the local density of electrons, the self-consistent electric field satisfies $\text{div} E = 4\pi \langle x; t \rangle$. The operator $\hat{\Gamma} = \hat{\Gamma}_{dis} + \hat{\Gamma}_{ph} + \hat{\Gamma}_{ee}$ is the collision integral [8] with the respective contributions coming from disorder, electron-phonon and electron-electron scattering. The Langevin random source $J(p; x; t)$ in (1) represents the fluctuations of the net flux of particle in phase space at $(p; x)$. These fluctuations are the consequence of the scattering processes

described by the respective collision integrals, and should be treated consistently along with the collision integrals to account for the conserved quantities. For the physics considered here (and disregarding quantum corrections) the interplay among the various scattering channels can be neglected. Within the same accuracy the Langevin flux is the sum of three uncorrelated random processes $J = J_{dis} + J_{ee} + J_{eph}$. Its correlation function can be found employing semiclassical considerations [9]. The random flux correlator can be written schematically as $\langle J(p; x; t) J(p^0; x^0; t) \rangle = \langle x - x^0 \rangle(t - t^0) H[p; p^0; x]$, where $H[p; p^0; x] = H_{dis}[p; p^0; x] + H_{ee}[p; p^0; x] + H_{eph}[p; p^0; x]$. Explicit expressions for the collision integrals and the pair correlation functions of the random fluxes are available [8, 15].

We consider a two-terminal system subject to an external voltage bias V ; the ambient temperature at the reservoirs is T . Transport through this system is a stochastic process. The charge transmitted over a time interval t is $Q(t) = \int_0^t dt I(t')$ where $I(t)$ is the electric current. The number of particles $n = Q/e$ is a fluctuating quantity described by probability distribution function $P(n)$. The Fourier transform of the latter $\langle e^{i n} \rangle = \int P(n) e^{i n} dn$ is the characteristic function which provides a complete statistical description of the problem. Constructing a

field theory it is natural to look for a characteristic function in a functional integral representation [6]

$$\tilde{Z} = \int \mathcal{D}f(p; x; t) \mathcal{D}\tilde{f}(p; x; t) \exp(iS[f; \tilde{f}]): \quad (3)$$

Here $f; \tilde{f}$ are independent bosonic fields and the measure of integration is \mathcal{D} . The field $f(p; x; t)$ is the phase space density operator within the KTF; $\tilde{f}(p; x; t)$ is an auxiliary 'counting field'. The field $\tilde{f}(p; x; t)$ accounts for fluctuations of this 'counting field'. The effective action consists of two parts

$$S[\tilde{Z}] = S_{D_{yn}} + S_{Noise}[\tilde{Z}]: \quad (4)$$

The propagation of fluctuations in phase space is described by the dynamical part of the effective action:

$$iS_{D_{yn}} = i \int_1^Z dx \int_{P^0}^Z dt \frac{1}{2} \left(\hat{L} + \hat{I} \right) f(p; x; t) \tilde{f}(p; x; t): \quad (5)$$

The second part of the action accounts for the generation of fluctuations:

$$iS_{Noise}[\tilde{Z}] = \int_1^Z dx \int_{P^0}^Z dt \frac{1}{2} \left(\tilde{f}(p; x; t) + f(p; x; t) H[p; p^0; x] \tilde{f}(p^0; x; t) + f(p^0; x; t) \right): \quad (6)$$

The effective action, Eqs. (5) and (6), can be obtained by exponentiating the equation of motion (1) with a help of the Lagrange multiplier $\tilde{f}(p; x; t)$ and averaging the resulting e^{iS} over the Langevin fluxes. The counting field, as a function of p, x , and t should be chosen accordingly to the problem one is interested in. It is illuminating to note that once the fluctuations of counting field are neglected in S_{Noise} , the effective action, (4) is equivalent to the standard KTF (i.e., it yields correctly the current-current fluctuations but not higher cumulants).

So far our treatment has been general. We now assume that the electron gas is degenerate and consider a quasi one-dimensional disordered conductor, where the momentum relaxation length is much shorter than the system length ($l_{tr} \ll L$). The problem then simplifies considerably. Additional important parameters are the diffusion coefficient, D , the single particle density of states at the Fermi level and the cross section S . Together they define the Ohmic conductance $G = e^2 D S / L$. Within the disordered approximation, (6) takes the form

$$iS_{D_{yn}} = 2 \int_0^L dx \int_1^Z dt f(x; t) \left(\frac{\partial}{\partial t} - D \frac{\partial^2}{\partial x^2} + eE \frac{\partial}{\partial x} \right) \tilde{f}(x; t) + \hat{I}_{eph} + \hat{I}_{ee} f(x; t); \quad (7)$$

where $\epsilon = p^2/2m$ is the kinetic energy. Eq.(7) describes propagation of fluctuations in disordered system in the presence of a long range Coulomb interaction. In the absence of the Coulomb interaction, one should set $E = 0$.

The noise-generating part of the action will also have three contributions, corresponding to elastic (impurity), electron-phonon and electron-electron scattering. The noise generated through elastic scattering is given by

$$iS_{Noise}^{dis} = \int dx dt D \left(\tilde{f}(x; t) + eE \frac{\partial f(x; t)}{\partial x} + \frac{\partial \tilde{f}(x; t)}{\partial x} \right)^2 f(x; t) (1 - f(x; t)): \quad (8)$$

One can check that the self-consistent field E does not affect the low frequency current statistics. Indeed,

a static electric field $E = -\partial_x \phi(x)$ can be eliminated from Eqs. (7), (8) by redefining the energy, $\epsilon \rightarrow \epsilon + e\phi(x)$. This implies that the counting statistics is the same in the interacting and non-interacting problems for frequencies much smaller than the inverse Thouless time, $\omega \ll D/L^2$. We will consider this regime below and thus set $E = 0$.

We are interested in the statistics of the number of particles traversing the system during a certain time window, irrespective of how this flow is distributed over that interval. We thus select the counting field to be

$$\chi(t) = \begin{cases} i\epsilon & ; 0 < t < \tau; \\ 0 & ; \text{otherwise}; \end{cases} \quad (9)$$

Here ϵ is the argument of the characteristic function $\chi(\epsilon)$.

To analyze the counting statistics we use a saddle point approach, which is justified by a large value of the dimensionless conductance $g = G/(\epsilon^2/h) \gg 1$. Note that this is the same condition which also allows us to neglect the quantum corrections to the current statistics due to the weak localization and the interaction (Altshuler-Aronov) effects. The boundary conditions for the distribution function are $f(\epsilon; 0) = f^L(\epsilon)$; $f(\epsilon; L) = f^R(\epsilon)$ where $f^{L(R)}$ are the equilibrium distribution functions of the electrons on the left (right) reservoir. The auxiliary field ϕ vanishes at the boundaries.

Current statistics and inelastic processes. When the inelastic processes can be neglected (which requires that the shortest inelastic length is longer than the sample size, $l_{in} \gg L$), the saddle-point equations following from (7), (8) permit an analytical solution [12, 19, 20]. In particular, in the shot-noise limit ($eV \gg T$) the resulting action reads (including a source term for the average current, $iS^1 = iGV \tau$)

$$iS^{dis} = iGV \tau \text{arcsinh}^2 \left(\frac{p}{e^{iL}} - 1 \right); \quad (10)$$

The found characteristic function, $\chi(\epsilon) = \exp[iS^{dis}g]$ eq.(10), agrees with the known result for the counting statistics obtained earlier [5, 6, 14] within the quantum approaches. This demonstrates the equivalence [12], up to $1/g$ corrections, between our effective theory and the model formalism [6]. Indeed, in this case the above effective action can also be obtained from a particular parametrization of non-linear model [13].

In the opposite limit, $l_{in} \ll L$, the collision integral plays a major role. There is only a small number of hydrodynamic modes (determined by the conservation laws) that propagate for a distances longer than l_{in} . For electron-electron collisions these are the fluctuations of the electron density and the energy (or, in other words, of the chemical potential and the temperature). By the same token, one has to keep only these two modes in the noise part of the action, S_{noise} , Eq. (8). In the presence of a strong electron-phonon scattering, $l_{eph} \ll L$, the collision integral \hat{I}_{eph} conserves only the number of electrons. The role of the electron-phonon scattering

thus amounts to projecting the action onto the mode of the density (chemical potential) fluctuations.

We consider now the case of a strong electron-electron scattering, $l_{ee} \ll L$, in detail. After projecting the fluctuating fields $\phi(p; x; t)$ and $\epsilon(p; x; t)$ in (5), (6) onto the relevant subspace, we are left with the fields ϕ and T that parametrize the fluctuations of a chemical potential and the temperature, respectively, and the conjugate fields ϕ and T . In terms of these fields the action (5), (6) takes the form

$$iS_{dyn} = i \int_0^{\tau} dx dt \left[\frac{D}{6} (\partial_x T)^2 + D \partial_x (\phi + \partial_x T) \right]; \quad (11)$$

$$iS_{noise} = \int_0^{\tau} dx \int_{-1}^1 dt T \left[D (\partial_x \phi + \partial_x T)^2 + D (\partial_x T)^2 \right]; \quad (12)$$

Here we introduced a thermal diffusion coefficient $D(T) = v_F^2 \left(\frac{1}{\tau_r} + l_{ee}^{-1}(T) \right)^{-1} \approx 3$, where

$$\frac{1}{l_{ee}(T)} = \frac{3}{2} \frac{1}{T} \int \frac{d\mathbf{p}}{4\pi} \frac{d\mathbf{p}'}{4\pi} n_0 n_0' H_{ee}[\mathbf{p}; \mathbf{p}'] \frac{1}{T}; \quad (13)$$

with $n; n'$ being the direction of the momenta $\mathbf{p}; \mathbf{p}'$ on the Fermi surface. Physically, l_{ee}^{-1} is the relaxation rate of the energy current due to the electron-electron collision processes. As we are going to show, there are two strikingly different cases. For $l_{tr} \ll l_{ee}$, which is the commonly considered hot-electron regime, the thermal diffusion and particle diffusion constants coincide, $D = D'$. Our action (11), (12) then agrees with that obtained in [17, 18] in the framework of a stochastic path integral formalism. In this regime, the electron-electron scattering does not affect the FCS qualitatively but only modifies the numerical coefficients in the expressions for current cumulants S_n . For example, for $eV \gg T$ one finds $S_2 = (3/4)e^2GV$, $S_3 = (8/15 - 9/16)e^3GV$, etc., instead of $S_2 = \frac{1}{3}e^2GV$, $S_3 = \frac{1}{15}e^3GV$;::: for non-interacting electrons.

The situation is qualitatively different in the regime of very strong electron-electron scattering (or, alternatively, of a very clean system), $l_{ee} \ll l_{tr}$, when the electrons form a liquid flowing through a weak disorder potential. In the absence of umklapp processes, the electron-electron scattering conserves the total momentum, so that the density diffusion is much faster than the thermal diffusion, $D' \gg D = D(T) \approx 1$. The saddle-point equations following from (11), (12) and determining the FCS of hot electrons do not permit an analytic solution. To obtain a qualitative understanding of the problem we first consider a simplified model, approximating $D(T)$ by a constant D . In this case, the action (11), (12) allows us to determine the scaling dependence of the FCS on τ . By performing a rescaling of fields $\phi \rightarrow \sqrt{\tau} \phi$, $T \rightarrow \sqrt{\tau} T$

in (11), (12) we find that () has the following scaling form ,

$$() = \exp f \quad ^2 G T t F (; eV=T) g; \quad (14)$$

where the function $F(x; y)$ governs the FCS at $\beta = 1$, $() = \exp f \quad G T t F (; eV=T) g$. This implies that in the equilibrium limit, $T \rightarrow eV$, the cumulants S_n with $n \geq 3$ in the electronic liquid are strongly enhanced compared to their values in the conventional hot-electron regime ($\beta = 1$). Specifically, the enhancement factor is 2^m for the cumulants S_{2m+1} and S_{2m+2} . In the opposite limit ($T \rightarrow 0$ at finite V), the enhancement starts already from S_2 , with S_n enhanced by the factor 2^{n-1} . Physically, the enhancement of higher moments of current fluctuations is due to the coupling between the mode of electric conductivity and the slow mode of the thermal conductivity.

So far we have used a model of a temperature independent thermal diffusion coefficient. On the other hand, in the Fermi-liquid theory, $D(T) = D_F v_F^2 = T^2$. We argue now that in a realistic situation the above strong enhancement of the cumulants in the $\beta_{ee} \rightarrow \beta_{tr}$ regime remains parametrically valid, with 2 replaced by $D = D(T_e)$, where T_e is the characteristic temperature in the system. In particular, in the equilibrium situation ($V = 0$), T_e is simply given by the temperature T of the reservoirs, $^2 \rightarrow \beta_{tr} = \beta_{ee}(T)$. In the non-equilibrium limit ($T = 0, V$ finite), the effective temperature is found to increase exponentially with V [16], yielding asymptotically $^2 \rightarrow \exp f(3=4^2) \beta_{tr} = \beta_{ee}(V) g$.

Finally, we consider the case of a strong electron-phonon scattering. The only soft modes allowed in this case ($\beta_{ph} \rightarrow \beta_L$) are fluctuations of chemical potential. The saddle point equations $\partial^2 = 0$; $\partial^2 = 0$ along with the boundary conditions lead to the trivial solution

$$(x) = (0) + (x=L) eV; \quad (x) = 0; \quad (15)$$

The resulting action

$$iS^{e-ph} = -tG [T^{-2} + iV] \quad (16)$$

represents Gaussian fluctuations, so that higher cumulants are suppressed by the electron-phonon scattering. This is consistent with earlier predictions based on qualitative arguments [1] and calculations of the third cumulant [6]. However, to the best of our knowledge, it is the first analytic derivation of this general result.

To summarize, we have proposed an effective theory describing the creation and evolution of fluctuations in phase space for degenerate electron gases and liquids. We have employed this theory to analyze statistical properties of noise in disordered conductors. We have reproduced known results for free fermion and confirmed the prediction of Gaussian statistics for the case of strong electron-phonon scattering. We have analyzed the effect of electron-electron scattering on the statistics of current fluctuations and established a scaling relation governing the dependence of the FCS on the ratio $\beta_{tr} = \beta_{ee}$, showing that the statistics differ parametrically between the $\beta_{tr} \rightarrow \beta_{ee}$ and $\beta_{tr} \rightarrow \beta_{ph}$ cases. Specifically, in the latter, when the electrons form a liquid flowing through weak disorder, the current cumulants are parametrically enhanced by electron-electron scattering. This is related to the coupling of the density mode to the slow mode of temperature fluctuations. The regime $\beta_{tr} \rightarrow \beta_{ee}$ is thus most favorable for experimental investigation of higher cumulants of current.

We acknowledge useful discussions and correspondence with D. Bagrets, M. Buttiker, I. Gruzberg, Y. Nazarov, S. Pilgram, and P. Wiegmann. This work was partially supported by the NSF MRSEC Program under grant DMR-0213745, by the Minerva Foundation, the US-Israel BSF, the ISF of the Israel Academy of Science and the Alexander von Humboldt foundation (through the Max-Planck Award), EC RTN on Spintronics. DBG and ADM acknowledge support by the Minerva Einstein Center during their visit to the WIS.

Note added. As this work was nearing completion we have learned of several related works [17, 18, 19, 20] that appeared in the cond-mat preprint archive.

[*] Also at Petersburg Nuclear Physics Institute, 188350 St. Petersburg, Russia.

[1] For a review see Ya. M. Blanter and M. Buttiker, Phys. Rep. 336, 1 (2000).

[2] G. P. Lesovik, JETP Lett. 49, 592 (1989).

[3] L. S. Levitov and G. B. Lesovik, JETP Lett., 58, 230 (1993).

[4] C. W. J. Beenakker and M. Buttiker, Phys. Rev. B 46, 1889 (1992).

[5] Yu. Nazarov, Ann. Phys. (Leipzig) 8, 193 (1999).

[6] D. B. Gutman, Y. Gefen, and A. D. Mirlin, Quantum Noise in Mesoscopic Physics (Delft, 2002), ed. Yu. Nazarov (Kluwer 2003), p. 497; cond-mat/0210076.

[7] L. D. Landau and E. M. Lifshitz, Statistical Physics 1,

(Pergamon Press, Oxford 1980), Ch. 118.

[8] E. M. Lifshitz and L. P. Pitaevsky Physical Kinetics, (Pergamon Press, Oxford 1981), Ch 10.

[9] A. Ya. Shul'man and Sh. M. Kogan, Sov. Phys. JETP 29, 3 (1969).

[10] K. E. Nagaev Phys. Rev. B 66, 075334 (2002); see also K. E. Nagaev, P. Samuelsson, S. Pilgram, Phys. Rev. B 66, 195318 (2002).

[11] S. Pilgram, A. N. Jordan, E. V. Sukhonukov, and M. Buttiker, Phys. Rev. Lett. 90, 206801 (2003).

[12] D. B. Gutman, A. D. Mirlin, and Yuval Gefen, cond-mat/0403436, v1.

[13] D. A. Bagrets, private communication.

[14] H. Lee, L. S. Levitov, and A. Yu. Yakovets, Phys. Rev. B

- 51, 4079, (1995).
- [15] Sh. Kogan, *Electronic Noise and Fluctuations in Solids* (Cambridge University Press, Cambridge 1996), Ch 3.
- [16] E. G. M. Ishchenko, *Phys. Rev. Lett.* 85, 4144 (2000).
- [17] S. Pilgram, *Phys. Rev. B* 69, 115315 (2004) (cond-mat/0401577).
- [18] S. Pilgram, K. E. Nagaev, and M. Buttiker, cond-mat/0401632.
- [19] A. N. Jordan, E. V. Sukhorukov, and S. Pilgram, cond-mat/0401650.
- [20] T. Bodineau and B. Derrida, cond-mat/0402305.