

Weak dynamical localization in periodically kicked cold atomic gases

C. Tian¹, A. Kamenev¹, and A. Larkin^{1,2,3}

¹ Department of Physics, University of Minnesota, Minneapolis, MN 55455, USA

² William I. Fine Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455, USA

³ L. D. Landau Institute for Theoretical Physics, Moscow, 117940, Russia

(Dated: April 14, 2024)

Quantum kicked rotor was recently realized in experiments with cold atomic gases and standing optical waves. As predicted, it exhibits dynamical localization in the momentum space. Here we consider the weak localization regime concentrating on the Ehrenfest time scale. The latter accounts for the spread-time of a minimal wavepacket and is proportional to the logarithm of the Planck constant. We show that the onset of the dynamical localization is essentially delayed by four Ehrenfest times and give quantitative predictions suitable for an experimental verification.

PACS numbers: 05.45.-a, 42.50.Vk, 72.15.Rn

Unprecedented degree of control reached in experiments with ultra-cold atomic gases [1] allows to investigate various fundamental quantum phenomena. A realization of quantum kicked rotor (QKR) is one such possibility that recently attracted a lot of attention [2, 3, 4, 5]. To this end cold atoms are placed in a spatially periodic potential $V_0 \cos(2k_L x)$ created by two counter-propagated optical beams. The potential is switched on periodically for a short time $\tau_p \ll T$, giving a kick to the atoms; here T is a period of such kicks. The evolution of the atomic momentum distribution may be monitored after a certain number of kicks. If the gas is sufficiently dilute [6], one may model it with the single-particle Hamiltonian, that upon the proper rescaling takes the form [7] of the QKR:

$$\hat{H} = \frac{1}{2} \hat{I}^2 + K \cos \frac{X}{n} \quad (t \rightarrow n): \quad (1)$$

Here $2k_L x$ and time is measured in units of the kick period, T . The momentum operator is defined as $\hat{I} = \hbar k \hat{\theta}$, where the dimensionless Planck constant is given by $k = 8\hbar T k_L^2 / (2m)$. Finally, the classical stochastic parameter is $K = kV_0 \tau_p / \hbar$.

The classical kicked rotor is known to have the rich and complicated behavior [8]. In particular, for sufficiently large K ($\gtrsim 5$), it exhibits the chaotic diffusion in the space of angular momentum [8]. The latter is associated with the diffusive expansion of an initially sharp momentum distribution: $\langle I^2(t) \rangle = \langle I(t) - I(0) \rangle^2 = 2D_{cl}t$ (dashed line on Fig. 1). For sufficiently large K , the classical diffusion constant may be approximated by $K^2/4$ [8]. The higher order correction is an oscillatory function of the stochastic parameter, i.e., $D_{cl}(K) = \frac{1}{4}K^2 (1 - 3J_2(K) + 2J_2^2(K))$ [9, 10]. It was realized a while ago [11, 12] that quantum interference destroys the diffusion in the long time limit and leads to localization: $\langle I^2(t) \rangle \sim K^2$ at $t > t_L$, $D_{cl} = K^2/2 = D_{cl}$, where the localization length is given by $\ell = D_{cl}/K$. For a large localization length $\ell \gg k$, there is a long crossover regime, $1 < t < t_L$, between the classical diffusion and quantum




FIG. 1: The momentum dispersion for $K = 6.1$ and $k = 0.6$ { full line; the classical limit ($K \rightarrow 0$) { dashed line, standard weak localization ($t_E = 0$) { dashed-dotted line; the limit $K \rightarrow 0$, Eq. (3) { long-dashed line.

localization.

It was suggested in Ref. [13], that the QKR may be mapped onto the one-dimensional Anderson localization with the long range disorder. The universal long-time behavior of the latter is described by the non-linear sigma model [14], resulting in the standard weak localization correction [15, 16]: $\langle I^2(t) \rangle = 2D_{cl}t (1 - 0.75 t/t_L)$ (dashed-dotted line on Fig. 1). Notice, that the correction is linear in k and non-analytic in time. In an apparent contradiction with this fact, explicit studies [17, 18] of the first few kicks show only renormalization of the diffusion constant starting from terms k^2 . The aim of this paper is to develop a quantitative description of the classical to quantum crossover for the QKR that, in particular, accounts for these conflicting observations.

It is known in various contexts [19] that such crossover involves an additional time scale, t_E , called an Ehrenfest (or breaking) time. For a generic quantum mapping, it was first shown by Berman and Zaslavski [20, 21], that quantum corrections become comparable to the classical limit at the time t_E . This is the time needed for a minimal quantum wavepacket, $\ell_0 \sim 1/k$, to spread uniformly over the angular direction. Due to the chaotic mo-

tion, trajectories diverge as $\langle r \rangle(t) = r_0 e^{\lambda t}$, where λ is the classical Lyapunov exponent. For $K \gg 1$, $\lambda = \ln(K/2)$ [8]. Estimating $\lambda_0 \approx K/2$ (a typical momentum dispersion after one kick, one finds for the Ehrenfest time:

$$t_E = \frac{1}{\lambda} \ln \frac{r}{r_0} = \frac{1}{\ln(K/2)} \ln \frac{r}{r_0} \quad (2)$$

It is widely believed [12, 20, 21, 22] that this intermediate, $1 < t_E < t_L$, time scale is indeed relevant for the quantum evolution of classically chaotic systems. This observation was put on the quantitative basis in Ref. [23] in the context of localization caused by classical scatterers. In this paper we adopt methods of Ref. [23] to the essentially different problem of the QKR.

In the leading order in k we found for the momentum dispersion:

$$\hbar^2 \langle t \rangle = 2D_{cl} t - \frac{8k^2 D_{cl}}{3} (t - 4t_E) (t - 4t_E)^{3/2}; \quad (3)$$

where $\langle t \rangle$ is the step function (long-dashed line on Fig. 1). At relatively large time, $t > t_L$ our result approaches the standard weak localization, mentioned above. However, corrections of the order of k are absent for $t < 4t_E$. The delay is caused by the interference nature of the localization. Indeed, the first correction originates from the interference of two closed-loop counterpropagating trajectories, see Fig. 2. It takes time about t_E for classical trajectories passing through (almost) the same value of the momentum to diverge and take counterpropagating roots. As a result, the interference effects are practically absent at smaller times and show up only after $4t_E$.

One may show that the time interval $0 < t < 4t_E$ is protected from higher order weak localization corrections as well. For example, in the second order weak localization correction there are two diagrams [24] proportional to $k^2 (t - mt_E) (t - mt_E)^2$ with $m = 6$ and $m = 8$ correspondingly. This fact agrees with the perturbative studies of the QKR dynamics after a few kicks [2, 11, 12], where no localization effects were seen. (Though the classical diffusion coefficient, D_{cl} , is renormalized as an analytic function of k^2 .) It is important to mention, however, that in reality there is no nonanalyticity at the point $t = 4t_E$ as may seem from Eq. (3). The small localization corrections, nonanalytic in k , do exist for $t < 4t_E$. They are associated with the fluctuations of the Ehrenfest time. Since in the quantum mechanics the minimal separation between the trajectories is about $r_0 \approx \hbar/k$, it takes a finite time (t_E) for them to diverge. This time may fluctuate depending on initial conditions. The fluctuations are characterized by the time scale $t_E = t_E^2 = t_E^2$, where (cf. Eq. (2)) $t_E^2 = \hbar^2 / \langle \dot{r}^2 \rangle \approx 0.82$ for sufficiently large K and the angular brackets denote averaging over the initial angle. The interference between rare trajectories, diverging




FIG. 2: The first quantum correction to the density-density correlator: (a) one-loop weak localization diagram; (b) its image in the momentum space; (c) semiclassical Hilbert box.

faster than the typical ones, leads to quantum corrections at $t < 4t_E$ of the form:

$$\hbar^2 \langle t \rangle = 2D_{cl} t - \frac{5}{3} \frac{k^2 D_{cl}}{256} (t_E)^{3/2} f \frac{4t_E}{t_E} t; \quad (4)$$

where $f(0) = 1$ and $f(x) = \frac{5}{4} x^{5/2} e^{-x^2} = 1.6$ for $x = 1$. Localization correction including effect of the Ehrenfest time fluctuations is depicted on Fig. 1 by a full line. It is rather close to the prediction of Eq. (3) (long-dashed line), however the singularity at $t = 4t_E$ is rounded.

Having outlined our main results, we turn now to some details of the calculations. One starts from introducing the exact one period evolution operator as: $\hat{U} = \exp i(K=k) \cos \hat{p} \exp i\hat{p}^2/(2k)$. All physical quantities may be expressed in terms of the matrix elements of \hat{U}^n , where n stays for the number of kicks (time). We shall be particularly interested in the four-point density-density correlator, defined as:

$$D(l_+; l_-; l_+^0; l_-^0; l_+; l_-) = \sum_{n, m=0}^{\infty} \langle l_+ | \hat{U}^{n+m} | l_- \rangle \langle l_+^0 | \hat{U}^n | l_-^0 \rangle \langle l_+ | \hat{U}^m | l_- \rangle \langle l_+^0 | \hat{U}^m | l_-^0 \rangle; \quad (5)$$

where $|l_{\pm}\rangle$ denote momentum eigenstates. First we note that averaging over $(l_+ + l_-)/2$ leads to $n = n^0$ [14]. Then performing the standard Wigner transform, one passes to the variables $l = (l_+ + l_-)/2$ and $\{l, l'\}$ (the Fourier transform of $l_+ - l_-$ (and similarly for the prime variables)). Since the most unstable direction of the underlying classical dynamics is along the direction [8], any initially smooth distribution quickly relaxes in this direction. Thus, averaging over l and l' may be performed.

$D(\mathbf{l}) = D_{cl} + D(\mathbf{l})$ with

$$D(\mathbf{l}) = \frac{kD_{cl}}{W^2} \sum_{\mathbf{l}'} \frac{d'}{l' + D_{cl}'} : \quad (13)$$

Equations (12) and (13) constitute the main analytical results of this work. They describe quantitatively dynamical weak localization of the QKR with the account for the Ehrenfest time phenomena. One may naturally express the time evolution of the momentum dispersion in terms of the frequency-dependent diffusion coefficient. The exact relation reads as:

$$\hbar l^2(t) = \sum_{\mathbf{l}'} \frac{d!}{l'^2} \frac{e^{i\mathbf{l}'t}}{l'^2} D(\mathbf{l}') : \quad (14)$$

Neglecting fluctuations of the Ehrenfest time ($\tau_2 \rightarrow 0$ in Eq. (12)) and performing frequency and angle integrations in Eqs. (14), (13), one obtains Eq. (3) for the momenta dispersion. Notice that in this approximation the evolution is purely classical at $t \gg t_E$. To account for the quantum corrections at $t < 4t_E$ one needs to keep the τ_2 term in Eq. (12). The straightforward integration leads to Eq. (4).

Our results, Eqs. (3), (4), are expected to be quantitatively accurate if separations between the relevant time scales: $1 < t_E < t_L$ are large enough. This is the case when the two dimensionless constants satisfy inequalities: $k < 1 < K$. (In the experiments, we are aware of [2, 3], $k > 2$, and thus $t_E < 1$.) Another restriction has to do with the dephasing time, τ_2 [28]. The latter may originate from nonperfect periodicity of the kicks (noise) [29], spontaneous emission [3], as well as from many-body collisions between the atoms [6]. Whatever the nature of the dephasing time, one needs to ensure $t_E < \tau_2$ to observe the weak localization.

We would like to thank A. Altland, J. Liu and C. Zhang for helpful discussions. Useful comments from S. Fishman are gratefully acknowledged. C.T. and A.L. are supported by NSF Grant No. 0120702; A.K. is A.P. Sloan fellow.

s-wave scattering length. The corresponding dimensionless scattering time is $\tau_s = 1/(T v)$, where v is a typical atomic velocity, that may be estimated as $v \sim \frac{\hbar k_F}{m}$. This leads to the self-consistent estimate of the dimensionless dephasing time: $(1/k_L = K)^{2=3}$.

- [7] The difference with the conventional QKR, is in the boundary condition on the wave-function: $(\mathbf{l} + 2) = (\mathbf{l})e^{i\mathbf{l} \cdot \mathbf{b}}$, where $\mathbf{l} \in [0; 1]$ is a fractional part of an atom's momentum in units $2\hbar k_L$. It may be eliminated by the gauge transformation that introduces an Aharonov-Bohm flux into the kinetic energy $(\hat{\mathbf{l}} - \mathbf{l}_0)^2 = 2$. Such flux does not affect dynamical localization, though averaging over \mathbf{l}_0 helps to eliminate resonances [22].
- [8] B.V. Chirikov, Phys. Rep. 52, 263 (1979); A.L. Lichtenberg and M.A. Leiberman, Regular and Chaotic Dynamics (Springer-Verlag, Berlin 1991).
- [9] A.B. Rechester and R.B. White, Phys. Rev. Lett. 44, 1586 (1980); A.B. Rechester, M.N. Rosenbluth, and R.B. White, Phys. Rev. A 23, 2664 (1981).
- [10] M. Khodas and S. Fishman, Phys. Rev. Lett. 84, 2837 (2000); Erratum, ibid. 84, 5918 (2000); M. Khodas, S. Fishman, and O. Agran, Phys. Rev. E 62, 4769 (2000).
- [11] G. Casati, B.V. Chirikov, J. Ford, and F.M. Izrailev, in Stochastic Behavior of Classical and Quantum Hamiltonian Systems, Lecture Notes in Physics 93, edited by G. Casati and J. Ford, 334 (Springer, New York 1979).
- [12] B.V. Chirikov, F.M. Izrailev, and D.L. Shepelyansky, Sov. Sci. Rev., Sect. C 2, 209 (1981).
- [13] S. Fishman, D.R. Grempel, and R.E. Prange, Phys. Rev. Lett. 49, 509 (1982).
- [14] A.A. Itland and M.R. Zimbauer, Phys. Rev. Lett. 77, 4536 (1996); ibid. 80, 641 (1998); G. Casati, F.M. Izrailev, and V.V. Sokolov, ibid. 80, 640 (1998).
- [15] A.A. Itland, Phys. Rev. Lett. 71, 69 (1993).
- [16] D.M. Basko, M.A. Skvortsov, and V.E. Kravtsov, Phys. Rev. Lett. 90, 096801 (2003); V.E. Kravtsov, cond-mat/0312316.
- [17] D.L. Shepelyansky, Physica D 28, 103 (1987).
- [18] A.J. Daley and A.S. Parkins, Phys. Rev. E 66, 056210 (2002); G. Du et. al., cond-mat/0401346.
- [19] A.I. Larkin and Yu.N. Ovchinnikov, Zh. Eksp. Teor. Fiz. 55, 2262 (1968) [Sov. Phys. JETP 28, 1200 (1969)].
- [20] G.P. Bernan and G.M. Zaslavsky, Physica A 91, 450 (1978).
- [21] G.M. Zaslavsky, Phys. Rep. 80, 157 (1981).
- [22] F.M. Izrailev, Phys. Rep. 196, 299 (1990).
- [23] I.L. Aleiner and A.I. Larkin, Phys. Rev. B 54, 14423 (1996).
- [24] C. Tian and A.I. Larkin, cond-mat/0310429.
- [25] M.R. Zimbauer, in Supersymmetry and Trace Formulas, Chaos and Disorder, edited by I.V. Lerner, J.P. Keating and D.E. Khmelnitskii (Kluwer Academic/Plenum, New York 1999).
- [26] S. Hikami, Phys. Rev. B 24, 2671 (1981).
- [27] In contrast to Ref. [23], the Hikami box for the QKR may be found explicitly without introducing a regularization.
- [28] D. Cohen, Phys. Rev. A 44, 2292 (1991).
- [29] D.A. Steck et. al., Phys. Rev. E 62, 3461 (2000); V. Miron et. al., Phys. Rev. E 61, 7223 (2000).

-
- [1] For a review see e.g. S. Chu, Science 253, 861 (1991).
 - [2] F.L. Moore et. al., Phys. Rev. Lett. 75, 4598 (1995); C.F. Bharucha et. al., Phys. Rev. E 60, 3881 (1999).
 - [3] H. Ammann et. al., Phys. Rev. Lett. 80, 4111 (1998).
 - [4] M.G. Raizen, Adv. At., Mol., Opt. Phys. 41, 43 (1999).
 - [5] C. Zhang, J. Liu, M. Raizen, and Q. Niu, Phys. Rev. Lett. 92, 054101 (2004).
 - [6] The two-particle scattering mean free path is $l = 1/(na^2)$, where n is the atomic concentration and a the