arXiv:cond-mat/0403541v1 [cond-mat.str-el] 21 Mar 2004

Instability Criteria for Vibrations of a Vortex Subject to a Gyro-force

B. A. Ivanov[1] and A. Yu. Galkin[2]

[1] *Institute of Magnetism, National Academy of Sciences and Ministry of Education of Ukraine, 36(b) Vernadskii avenue, 03142 Kiev, Ukraine*

[2] *Institute of Metal Physics, National Academy of Sciences of Ukraine, 36 Vernadskii avenue, 03142 Kiev, Ukraine*

PACS. 67.40.Vs – Vortices and turbulence.
PACS. 74.20.-z – Superconductivity, theory.
PACS. 75.60.-d – Domain effects, magnetization curves, and hysteresis.

Abstract. – Here we study the instability of the vortex motion caused by vortex velocity non-monotonous dependence of the friction force in the presence of a gyro-force. We demonstrate that even a weak gyro-force renders the condition for onset of instability more strict than for a vortex not subject to gyro-forces: either the friction force must exceed a threshold, or its non-monotonic velocity dependence must be a rather steep function.

The linear topological defects, vortices are one of the most widely studied objects in modern physics. The vortex instances can be found in conventional [1] and high-temperature [2] superconductors, superfluid liquids [3], dilute Bose-Einstein condensates [4–6], Bloch lines in the domain walls of ferromagnets [7,8], magnetic vortices in ferromagnets and antiferromagnets [9, 10], and even singular lines in non-linear optical beams (optical vortices) [11]. For the vortex dynamics description the classical concept of the elastic string is widely employed. The vortex possess elastic properties of the string and many peculiarities of the string dynamics are manifested in the vortex dynamics. It is well-known that if the friction force acting on the string depends non-monotonically on the string velocity, instability of the string motion results [12, 13]. When the string moves in a medium where the friction force acting on it, $\vec{F} = -(\vec{V}/V)F(V)$, depends non-monotonically on velocity within some range, and moves with a velocity V_0 such that $dF(V)/dV|_{V_0} < 0$, then small amplitude vibrations of the string, superimposed on this background motion, can be described as linear vibrations under negative friction. The elastic string provides one of the best-known examples of auto-oscillation. The auto-oscillations of a violin string, when the violin string is under action of a bow moving with a velocity V_0, are an example.

For vortices in condensed media a non-monotonous dependence $F(V)$ is often realized. The physical mechanisms can be very various: e.g. quasi-particle spectra rearrangement in the core of a moving Abrikosov vortex in superconductors [14], vortex interactions with a system of microdefects [15] or Cherenkov radiation of sound [8, 9]. For these cases a non-linear friction

© EDP Sciences

force, additional to the conventional viscous force, with the vortex velocity dependence $1/V^n$, where n varies from 0.5 [15] to 2 [8,9,14] appears.

An important feature common to dynamics of many types of vortices is the presence of a gyro-force, see references [1] - [11]. The gyro-force is formally equivalent to the Lorentz force that acts on a charged elastic string placed in the magnetic field parallel to the string line.

A gyro-force $|\vec{F}_g| = G \cdot V$ (G is the gyro-constant) directed perpendicular to the vortex velocity $\vec{V}$ is an inherent feature of vortex dynamics. Experimentally it has been observed in ferromagnets [7,8], in high-Tc superconductors in super-clean limit [16], and in optics [11]. The presence of a gyro-force for these media is a direct consequence of the topological character of the vortex [7–9]. For models having Lorentz-invariant dynamics a gyro-force is not an inherent feature of the vortex. An example of such a model is a nonlinear sigma–model, used for description of antiferromagnets [8, 9] and quantum nematic phases in Bose-Einstein condensates [6] and spin–1 non-Heisenberg magnets [17]. Even for these cases, a gyro-force could appear in the presence of an external magnetic field parallel to the vortex line [18, 19].

In this Letter we demonstrate that the question of the general stability of the motion of string-like objects (vortices) in viscous media needs to be revised in the presence of a gyro-force. It is shown that in this case the presence of non-monotonic dependence of the friction force, as well as the negative differential mobility does not serve as a stability condition and more complicated stability conditions arise. We will use the simple model of the elastic string, which has a gyro-force, which is a rather general one and its applicability falls outside the framework of the theory of vortices in ordered media.

Consider the vortex directed along the z-axis. Its dynamics may be described on the basis of the effective equation for a two-dimensional vector, $\vec{U} = \vec{U}(z,t)$ lying in the (x,y) plane and determining the displacement of the vortex center:

$$m\ddot{\vec{U}} - \sigma(\partial^2\vec{U}/\partial z^2) + G(\widehat{\vec{z}} \times \dot{\vec{U}}) = \vec{F}. \tag{1}$$

Here $\dot{\vec{U}} \equiv \partial\vec{U}/\partial t$, m, G and σ are the vortex mass, the gyro-force constant and the energy per unit length of the vortex, respectively. The friction force $\vec{F} \equiv \vec{F}(\dot{\vec{U}}) = -(\dot{\vec{U}}/|\dot{\vec{U}}|)F(|\dot{\vec{U}}|)$ is considered to be collinear to the instantaneous velocity of the vortex with positive value of $F(V)$.

In order to investigate the stability of the motion of the vortex when moving with a velocity close to V_0 we assume $\vec{U} = \vec{V}_0 t + \vec{u}(z,t)$, where $\vec{u}(z,t)$ is a small perturbation of the steady motion at $\vec{V}_0$, and linearize (1) (in effect, the friction force $\vec{F}$) over $\vec{u}$. In this linear approximation the friction force can be represented as a sum of components perpendicular and parallel to $\vec{V}_0$:

$$\vec{F} = -\vec{V}_0(\dot{\vec{u}}\vec{V}_0)(F'/V_0^2) - (\vec{V}_0 \times (\dot{\vec{u}} \times \vec{V}_0))(F/V_0^3), \tag{2}$$

where $F = F(V_0), F' = [dF(V)/dV]_{V=V_0}$. Then we can find the solution in Fourier representation, $\vec{u}(z,t) \propto \vec{u}_q \exp(\Lambda t + iqz)$, where Λ defines the character of the time evolution of small deviations from the steady motion. If the real part of Λ is negative then the value $-\mathrm{Re}\Lambda > 0$ serves as a damping coefficient for small vibrations of the vortex. Otherwise, if the real part of Λ is positive for some values of the wave number q, weak vibrations cause instability with the incremental factor $\mathrm{Re}\Lambda > 0$.

If the gyro-force is absent the situation is very simple. The equations for the components of $\vec{u}$ parallel and perpendicular to the $\vec{V}_0$ are then independent. They have different effective friction coefficients F' and F/V. The value of F/V is necessarily positive while F' can be either positive or negative within any given small velocity range. It is therefore evident that

for $G = 0$ vibrations with $\vec{u} \perp \vec{V}_0$ are damped, while the possible instability of vibrations with $\vec{u} \parallel \vec{V}_0$ is related to the condition $F' < 0$. Thus for $G = 0$ the sole criterion for instability of steady motion of the vortex at velocity $\vec{V}_0$ is $[dF(V)/dV]_{V=V_0} < 0$. If $G \neq 0$, then oscillations with $\vec{u} \perp \vec{V}_0$ and $\vec{u} \parallel \vec{V}_0$ become coupled and the characteristic equation for Λ is then a 4-th power equation. It can be presented in the simple form:

$$[m\Lambda^2 + (\Lambda/2)(F' + F/V) + \sigma q^2]^2 = (\Lambda/2)^2[(F' - F/V)^2 - 4G^2], \tag{3}$$

and solved exactly. The analysis demonstrates that the real parts of all four roots of this equation are negative, i.e. the motion is stable when two criteria are *simultaneously* satisfied: $F'(F/V) + G^2 > 0$ and $F' + (F/V) > 0$. (It is evident that these criteria do not depend on the mass and elasticity.) Thus, two *independent* criteria for instability exist, which correspond to violation of one or other of these two stability criteria. The first condition for the vortex motion to be unstable can be written as

$$-(F')(F/V) > G^2. \tag{4}$$

In the limit $G \to 0$ this condition coincides with the inequality $F' < 0$. The second instability condition

$$F' + (F/V) < 0 \tag{5}$$

can be satisfied however weak the friction and however large the gyro-constant G. Therefore this second criterion can also be significant for the case of interest when G is non-small.

Suppose that the friction force $F(V)$ is the sum of a regular viscous friction force with a constant velocity coefficient β and a non-linear friction force $f(V)$, which can be a decreasing function of V, so that $F(V) = \beta V + f(V)$. The condition $F' < 0$ requires the value of $f'(V)$ to be negative and that the maximal value of its modulus, reached at $V \cong V_m$, exceeds β, $|f'|^{max} > \beta$. It is evident that the condition $F' < 0$ can be met however weak the friction coefficient $\beta \sim |f'|^{max} \approx (f/V) \ll G$; it is only important that the values β and $|f'|^{max} \sim f/V$ should be comparable. However, at fairly large G, instability condition (4) is certainly more restrictive as it involves G. Estimating $|f'|^{max} \sim f(V_m)/V_m$, one can express the first instability condition (4) as $|f'|^{max} > \sqrt{G^2 + \beta^2}$, which cannot be fulfilled for large enough G and however weak the friction force f.

The instability condition (4) can acquire a clear physical meaning if one introduces the non-linear differential mobility $\mu(V)$ in respect to the external force F_e, $\mu(V) = dV(F_e)/dF_e$. In fact, the motion under action of a constant external force takes place at the Hall angle α_H regarding to $\vec{F}_e$, and the velocity is determined by the condition $G^2V^2 + F^2(V) = F_e^2$. The $\mu(V)$ can be presented as

$$\mu(V) = \frac{\sqrt{F^2(V) + G^2V^2}}{VG^2 + F(V)F'(V)} \tag{6}$$

and the instability condition (4) can be considered as the negative differential mobility $\mu(V) < 0$. The condition of the negative differential mobility for small friction and finite G could be met with difficulty, even if $F' < 0$.

Thus, for this case - large G, the condition (5) provides the better opportunity for instability: it does not involve G and can be met at low friction. However, there is another strong restriction that can prevent instability. If one extrapolates the function $f(V)$ by a power function, $f(V) = \eta/V^n$, where η is some coefficient, then F' can be negative at any ratio between η and β and at any $n > 0$. However, the condition (5) yields $(1-n)\eta/V^{n+1} + 2\beta < 0$ and is not fulfilled with $n \leq 1$. So here at any small G instability requires a rather steep dependence of $f(V)$.

As an example, for the dynamics of a single vortex in a media with weak inhomogeneities, $n = 1/2$ [15], the value of $f' + f/V = f/2V$ is always positive and the stability of the motion is not destroyed by the condition (5). Nevertheless, when a dense vortex lattice moves through defects $n = 2$ and instability due to the criterion (5) could appear [20].

In summary, the onset of instability of motion of a vortex subject to a gyro-force cannot be attributed merely neither to negative incremental friction $F' < 0$ nor of the negative differential mobility. It requires additional restrictions on the velocity dependence of the friction force $F(V)$ or on the value of the gyro-force G. In fact, two independent instability criteria arise. One of them is associated with appearance of the negative differential mobility, dV/dF_e and can be discussed as stability with respect to the *longitudinal* perturbations. The other condition is nothing to do with differential mobility and corresponds to the negative effective dumping coefficient averaged over the longitudinal and *transverse* (regarding to the velocity $\vec{V}$) oscillations of the vortex. For $G = 0$ the condition $dV/dF_e < 0$ coincides with $F' < 0$, however at $F', F(V)/V \ll G$ these two conditions are essentially different.

It is worth to note that most of above considered examples are related to vortex-like linear defects, especially vortices in superconductors and magnets. Nevertheless, the model of the string, which has a gyro-force is a rather general one and falls outside the framework of the theory of vortices in ordered media.

* * *

We thank J. E. L. Bishop and C.E.Zaspel for their useful comments on the manuscript. One of us (B.I.) thanks the grant I/75895 from the Volkswagen-Stiftung for partial support.

REFERENCES

[1] DeGennes P. G., *Superconductivity of Metals and Alloys* (Benjamin, New York) 1966.
[2] Blatter G., Feigelman M. V. , Geshkenbein V. B. , Larkin A. I., and Vinokur V. M., *Rev. Mod. Phys* , **66** (1994) 1125.
[3] Donnely R. J., *Quantized Vortices in Helium II*, edited by Goldman A. M., McClintoc P. V. E., and Springford M., Vol. **3** (Cambridge University Press, Cambridge, England) 1991.
[4] Mathews M. R., Anderson B. P., Halljan P. C. et al, *Phys. Rev. Lett.* , **83** (1999) 2498.
[5] Madison K. W., Chevy F., Wohlleben W., and Dalibard J., *Phys. Rev. Lett.* , **84** (2000) 806.
[6] Zhou Fei,, *Quantum Spin Nematic States in Bose-Einstein Condensates, ArXiv:cond-mat/0108473* 2002.
[7] Malozemoff A. P. and Slonczewski J. C., *Magnetic domain walls in bubble materials* (Applied Solid State Science, Supplement I, Academic, New York) 1979.
[8] Baryakhtar V. G, Chetkin M.V., Ivanov B. A. and Gadetskii S. N., *Dynamics of topological magnetic solitons. Experiment and theory* (Springer Tract in Modern Physics, Springer-Verlag, Berlin), Vol. **139**, 1994.
[9] Baryakhtar V. G. and Ivanov B. A., *Soliton thermodynamics of low-dimensional magnets*, edited by Khalatnikov I.M. *Sov.Sci.Rev., Section A.-Physics*, **16** (1992) 1 - 192.
[10] Kosevich A. M., Ivanov B. A. and Kovalev A. S., *Phys.Reports*, **194** (1990) 117.
[11] Cristou J., Tikhonenko V., Kivshar Yu.S., and Luther-Davids B., *Opt. Lett.*, **21** (1996) 1649; Kivshar Yu.S. and Luther-Davids B., *Phys. Reports*, **298** (1998) 81.
[12] Lord Rayleigh, John William Strett, *The theory of sound, 2-nd edition, in two volumes* (MacMillan, London) 1929.
[13] Andronov A. A., Vitt A. A., and Khaikin S. E., *Theory of oscillations* (Addison-Wesley, Reading, Mass.) 1966.

[14] Larkin A. I. and Ovchinnikov Yu. N., *Sov. Phys. JETP.*, **41** (1976) 960.
[15] Galkin A. Yu. and Ivanov B. A., *Phys. Rev. Lett.*, **83** (1999) 3053.
[16] Harris J. M. et al, *Phys. Rev. Lett.*, **73** (1994) 1711.
[17] Ivanov B.A. and Kolezhuk A. K., *Phys. Rev.*, **B68** (2003) 052401
[18] Ivanov B. A. and Sheka D. D., *Phys. Rev. Lett.*, **72** (1994) 404.
[19] Ivanov B. A. and Kolezhuk A. K., *Low. Temp. Phys.*, **21** (1995) 275.
[20] Galkin A. Yu. and Ivanov B. A., *Phys. Rev. B*, **66** (2002) 054507.