

Drift causes anomalous exponents in growth processes

Gunnar Pruessner

Department of Mathematics, Imperial College London, 180 Queen's Gate, London SW 7 2BZ, UK
(dated: 23 October, 2003)

The effect of a drift term in the presence of fixed boundaries is studied for the one-dimensional Edwards-Wilkinson equation, to reveal a general mechanism that causes a change of exponents for a very broad class of growth processes. This mechanism represents a relevant perturbation and therefore is important for the interpretation of experimental and numerical results. In effect, the mechanism leads to the roughness exponent assuming the same value as the growth exponent. In the case of the Edwards-Wilkinson equation this implies exponents deviating from those expected by dimensional analysis.

PACS numbers: 05.70.Np, 68.35.Ct, 68.35.Rh

The Edwards-Wilkinson (EW) equation [1], as it is discussed below, is probably the best-studied equation describing surface growth processes. Due to its linearity it is solvable by standard methods and has been studied analytically as well as numerically in great detail [2, 3, 4]. The equation is very well-behaved, so that the outcome of these investigations are usually quite predictable. There is no reason to suspect that well-accepted methods, such as dimensional analysis and coarse graining, produce wrong results, even if applied to the EW equation with an extra drift term (EWd), which still represents a linear problem.

However, it is shown below that, depending on the boundary conditions, such a drift term changes the exponents dramatically to anomalous values, which apparently have been missed in the literature. The drift in conjunction with the boundary condition poses a relevant perturbation to the original equation. While it is not possible to capture its effect by the simple methods mentioned above, it can be understood using physical arguments. The mechanism turns out to be very powerful and extends far beyond the EW problem.

The Edwards-Wilkinson equation [1] describes the temporal evolution of an interface characterized by its height $h(x;t)$ over a substrate of length L , $x \in [0;L]$, at time t under the influence of a thermal noise $\xi(x;t)$. In one dimension it reads

$$\partial_t h(x;t) = D \partial_x^2 h(x;t) + \xi(x;t); \quad (1)$$

with a diffusion constant or surface tension D . The initial conditions are usually [4, 5] chosen to be $h(x;t=0) = 0$ and periodic boundary conditions (PBC) are applied. The central property one is interested in is the roughness, defined as

$$w^2(t;L) = \frac{D}{2} \overline{E} - \frac{D}{2} \overline{E}^{-2}; \quad (2)$$

where \overline{A} denotes the spatial average, $\overline{A} = \frac{1}{L} \int_0^L dx A(x)$ and \overline{h} is the ensemble average, averaging over all realizations of the noise ξ . In order to determine w^2 , the

only property of ξ which enters is

$$\langle h(x;t) \xi(x';t') \rangle = \delta(x-x') \delta(t-t'); \quad (3)$$

where δ parameterizes the strength of noise. To fully specify the noise, the higher order correlations are usually chosen to be those of Gaussian white noise and the average is set to $\langle h(x;t) \rangle = 0$. Assuming a Family-Vicsek scaling behavior [6], three exponents, β , ν , and z , are defined for the asymptotic behavior of w^2

$$w^2(t;L) = a L^{2-\beta} G\left(\frac{t}{L^z}\right); \quad (4)$$

with appropriate, system dependent parameters a and b which make the universal scaling function $G(x)$ a dimensionless function of a dimensionless argument. This function is supposed to behave like $G(x) \sim x^{-2}$ for small arguments x and to converge to a non-zero value for large arguments. In the initial growth phase, w^2 is supposedly independent of L , so that $\beta = z$.

Because of the small number of independent parameters, the exponents characterizing (1) can be determined immediately, simply by using dimensional analysis. The roughness based on the ensemble of solutions of (1) can be written as

$$w^2(t;L;D;\xi) = \frac{2L}{D} G\left(\frac{tD}{L^z}\right); \quad (5)$$

with all independent parameters being listed on the left hand side. Eq. (5) is the unique way of writing w^2 in the form prescribed by (4) with all t -dependence being absorbed into a dimensionless function of a dimensionless argument. By assuming the existence of the appropriate limits or by simply comparing (5) to (4), the exponents are determined to be:

$$\beta = 1=2 \quad \nu = 1=4 \quad z = 2 \quad (\text{standard EW}) \quad (6)$$

Of course, this can easily be confirmed by exact solutions [4, 5].

The exponents (6) remain unchanged, if fixed boundary conditions (FBC), $h(x=0;t) = h(x=L;t) = 0$,

are introduced; the exact solution changes, but as the new boundary condition does not contain any new non-zero parameter and therefore cannot introduce any new scale, the scaling form Eq. (5) must necessarily remain unchanged; there are just not enough (independent) parameters for the form Eq. (5) to change.

On the face of it, the EW equation becomes much more complicated, if a drift or convection term is introduced into (1),

$$\partial_t \langle x; t \rangle = D \partial_x^2 \langle x; t \rangle + v \partial_x \langle x; t \rangle + \langle x; t \rangle; \quad (7)$$

where v denotes the drifting velocity. Such a term is probably present in most experimental setups, for example due to the presence of a gravitational field and a small tilt of the substrate but it is not obvious whether the resulting drift is strong enough to have a significant effect, (see also the crossover length calculated below). Here, it is worth mentioning the general role of a drift term as discussed in the context of anomalous aging by Luck and Mehta [7].

In case of PBC, the solution of Eq. (7) is related to the solution of Eq. (1) by a simple Galilean transformation, $\langle x; t \rangle = \langle x + vt; t \rangle$, which leaves the noise correlator (3) unchanged. Because of translational invariance, the correlator $\langle x; t \rangle \langle x^0; t^0 \rangle$ (spatially) depends only on $x - x^0 + v(t - t^0)$ and becomes independent of v for equal times $t^0 = t$. Because w^2 in (2) is a functional of the equal-time correlator, it is also independent of v . Thus, in case of PBC, the exponents remain those listed in Eq. (6),

$$\alpha = 1=2 \quad \beta = 1=4 \quad z = 2 \quad (\text{EW with PBC and drift}) \quad (8)$$

Dimensional analysis, however, leads to a scaling form

$$w^2(t; L; D; v) = \frac{L^2}{D} G\left(\frac{tD}{L^2}; \frac{vL}{D}\right); \quad (9)$$

the behavior of which is unknown; as is shown below, $G(x; y)$ might behave like $y^{-1=2}$ for large x .

It is very tempting and in fact leads to the exponents previously reported in the literature [8], to argue that the drift term in (7) is relevant compared to the diffusion term. In fact, this is what an ansatz $\langle x; b^z t \rangle = b^{-\alpha} \langle x; t \rangle$ suggests. The idea behind this much-used method [9, 10], is to determine the terms dominating the large scale behavior of Eq. (7). It is often applied to determine relative relevancy of higher order terms in Langevin equations. In fact, the exact relation

$$\langle x; t; bL; D; v \rangle = b^{\frac{z-1}{2}} \langle x; t; b^z; L; D; b^z; v \rangle \quad (10)$$

seems to indicate that w^2 behaves on large scales like on small scales with a diffusion constant D reduced by

a factor b^{-1} compared to the reduction of v . Thus, the diffusion constant is expected to drop out for large system sizes. Repeating the dimensional analysis without the diffusion term, i.e. setting $D = 0$, then yields

$$\alpha = 0 \quad \beta = 0 \quad z = 1 \quad (\text{EW d from coarse graining}); \quad (11)$$

This is consistent with a scaling law which can be derived from (10). It states that any derivable from dimensional analysis must obey [4]

$$\alpha = \frac{z-1}{2}; \quad (12)$$

Below it is shown that this is not the case. There is no reason to assume that w^2 becomes asymptotically self-averaging.

As shown above, this result, suggesting a smooth or flat interface [8], is wrong at least in the presence of periodic boundary conditions, because the velocity v simply disappears due to translational invariance. The key question is then: Are the exponents (11) recovered if the drift term cannot be gauged away? Does the coarse graining and dimensional analysis arguments presented above fail only because of translational invariance, which effectively removes v as free parameter from the problem? The answer to both of these questions turns out to be negative. The reason for that is an extremely efficient mechanism, effectively "wiping out" the stationary roughness.

In order to answer the questions raised above, fixed boundaries are applied again, so that the drift cannot be gauged away. It is worth stressing that fixed boundaries correspond to a finite substrate or a finite region of exposure to the noise, i.e. these boundaries are much more natural than periodic boundary conditions.

Eq. (7) with FBC can be solved using a saddle point approximation. In the following only a few technical details of this calculation are presented. After expressing (7) in a dimensionless form, the solution can be written as

$$\langle y; q \rangle = \int_0^1 dy^0 \int_0^1 dy^1 \int_0^1 dy^2 \dots \int_0^1 dy^z \quad d^0 \dots d^z \quad \langle y^0; y^1; \dots; y^z \rangle \quad (13)$$

where $y = x/L$, $p = t/(L^2=D)$, $q = vL/D$ and $\langle y; q \rangle = \int_0^1 \int_0^1 \dots \int_0^1 \langle x; t; L; D; v \rangle$ as well as $\langle y^i \rangle = L^{z-2} = (L^2/D) \langle x; t \rangle$ are dimensionless quantities. The propagator $\langle y^i \rangle$ of this problem essentially consists of two Gaussians "wrapped around a circle of radius 2 " from a mirror charge trick, which are closely related to Jacob's $\#_3$ -function [11, 12]. This sum is multiplied by an exponential $e^{-\frac{1}{2}(y^i - y^j)q} e^{-\frac{1}{4}q^2}$, which accounts for the drift:

$$\phi_0(y; i; \phi; q) = \frac{1}{4} \sum_{n=1}^{\infty} e^{-\frac{(y-y_0+2n)^2}{4}} e^{-\frac{(y+y_0+2n)^2}{4}} e^{-\frac{1}{2}(y-y_0)q} e^{-\frac{1}{4}q^2} \quad (14a)$$

$$= \frac{1}{2} \sum_{n=1}^{\infty} \sin(ny) \sin(ny_0) e^{-(n)^2} e^{-\frac{1}{2}(y-y_0)q} e^{-\frac{1}{4}q^2} \quad (14b)$$

where by convention $\sum_{n=1}^{\infty}$ and y_0 is the initial position. The saddle-point approximation is required when the spatial averages (2) are taken. One finds

$$w^2(t; L) = \frac{2}{3} \frac{t}{2D} \frac{t}{L} \frac{yL^{3/2}}{3L^2 \frac{t}{2D}} + \frac{t^2 yj}{2L^2} \quad \text{for } t \ll \frac{L}{vj} \quad (15a)$$

$$\frac{2}{3} \frac{t}{2D} \frac{t}{L} \frac{1}{2vj} \quad \text{for } t \gg \frac{L}{vj} \quad (15b)$$

which becomes exact for divergent $vL=D$, i.e. especially in the thermodynamic limit. Inspecting the leading terms immediately gives

$$z = 1/4 \quad z = 1/4 \quad z = 1 \quad (\text{EW with FBC and drift}) \quad (16)$$

This result is surprising, because these exponents are anomalous, as they do not correspond to what seems to be suggested by dimensional analysis: Not only do they contradict (6), (8) and (11), they do not even obey (12). In fact, neither the drift term, nor the diffusion term become irrelevant [19]; they are both crucial for (16). Moreover, the set of exponents switches from (8) to (16) if the boundary condition is changed from periodic to fixed in the presence of a drift term. The Edwards-Wilkinson equation with drift term and fixed boundary conditions is a linear problem displaying anomalous exponents.

The fact that the above result cannot be reconciled with dimensional analysis might suggest that the mechanism leading to these exponents is very subtle. However, it turns out that it can be understood quite easily.

The drift term makes the entire interface configuration move from one boundary to the other. Without noise, a peak starting somewhere in the bulk gets slowly moved by the drift to one of the boundaries, while diffusively broadening. It eventually disappears at the boundary. The time it spends between the boundaries depends on the starting position and the direction of the movement. The maximum time is $L=v$, which is also the maximum time, any noise-generated structure has to develop. However, as known from the model without drift, it takes time $L^2=D$ in order to fully develop the roughness. Thus, for $L^2=D < L=v$, i.e. $L < D=v = L_x$, the interface will remain in its initial growth phase; the characteristic length scale L_x represents an effective cutoff for correlations. At the same time it enables the system to display anomalous exponents. Depending on the direction of the drift, the interface "comes out" of the left boundary initialized




FIG. 1: A qualitative picture of an interface snapshot with its "local roughness", obtained in a numerical simulation; scales are irrelevant. Upper panel: An example of an interface profile with fixed boundaries and drift term. Lower panel: The ensemble averaged local width squared (numerical data, circles) is proportional to the square root of the position where measured, $x^{1/2}$ (fitted, dashed line).

to $x=0$ and moves to the right boundary, as shown in Fig. 1. The average "age" is proportional to $t_x = L/v$, so that according to (4) $w^2 / t_x^{1/2} / L^{1/2}$, therefore $z = 1/4$. Regarding ϕ , the interface cannot "see" the drift initially, so that $z = 1/4$ just like for the case without drift, (6). Indeed, even the amplitude of the leading term in (15a) corresponds to the amplitude obtained for the problem with periodic boundaries. The identity $z = 1/4$ already indicates $z = 1$, which can also be derived from the fact that saturation should be reached as soon as the interface has swiped through the system once, i.e. after $t_x = L=v$.

To test the validity of the results, Eq. (15) has been

compared to numerical simulations, based on a straightforward integration of (7) using Euler's method. For system sizes not too small ($L \geq 128$), even the higher order corrections were reproduced. The mechanism is illustrated in Fig. 1: The upper panel shows a snapshot of an interface configuration. The "local age" of the interface can be read off the local roughness (in an appropriate ad hoc definition) as shown in the panel below, because spending more time between the boundaries increases the local roughness according to $w^2 / t^{1-z} = (x-v)^{1-z}$, with x being the position where the roughness is measured.

The physical explanation presented above goes beyond the EW equation; provided that the crossover time of the original model without drift scales faster in L than t_x , i.e. $z > 1$, the argument should apply, so that at sufficiently large system sizes obtains the value of z and therefore $z = 1$. It is a very efficient mechanism, which works under very general circumstances even in the most simple, linear case. It therefore speaks a clear warning as to the interpretation of numerical and experimental studies: the true value of z might have been "washed away" by a very small drift.

The KPZ [2, 3, 4, 13] equation ($z = 3/2$) is particularly interesting: Its non-linearity is only important during the initial growth phase ($z = 3/2$) and becomes insignificant in the stationary regime ($z = 1/2$). However, with an additional drift term the equation should remain in the initial growth phase, with the non-linearity present in the stationary state. Indeed, using a Cole-Hopf transformation [3], the problem can be reduced to an equation similar to Eq. (7). However, preliminary numerical tests did not fully confirm this correspondence.

It is not yet completely clear how to generalize this argument to higher dimensions. For two dimensions it is tempting to speculate whether exponents observed in experimental molecular beam epitaxy are related to such a drift term, for example when $z = 1/4$ or when z is close to $z_{KPZ} = 0.24$ [10, 15]. Interestingly, only one boundary needs to be fixed in order to observe the phenomenon, namely the boundary the velocity points away from, in Fig. 1 the left boundary.

The mechanism stresses once more the relevance of boundary conditions as prominently pointed out by Landau and Binder [16]. However, it is worth emphasizing that in the present case, the change of boundary conditions leads to a change of the bulk critical exponents.

Even though the physical explanation provides a very clear picture of the mechanism, it is not obvious how a drift term exactly affects other models, with, for example, quenched or conserved noise, with an additional term or with additional scaling laws.

In conclusion we have presented a remarkably simple mechanism which reduces the roughness exponent to the value of the growth exponent for any small amount of drift in the Langevin equation in the presence of fixed boundary conditions, provided that in the original model,

the dynamical exponent z is larger than unity. On sufficiently large scale, this mechanism should be visible in many experimental and numerical systems. Most unexpectedly, it can even be found in the Edwards-Wilkinson equation, which consequently shows anomalous exponents, depending on the boundary conditions imposed.

The author would like to thank E. Brunet, E. Frey, H. J. Jensen, J. M. Lopez, J. M. Luck, A. Parry, B. Schmittmann, U. C. Tauber and R. K. P. Zia for very helpful discussions and for their hospitality. Also, he thanks H. Bruhn, C. Porter, I. Ruf, the Thoenen family, and especially the Max-Planck-Institut für Physik komplexer Systeme in Dresden for hospitality during NES-PHY 03. The author gratefully acknowledges the support of EPSRC.

Note added in proof: The author wants to thank J. Krug for pointing out that the effect of a drift term is discussed and referenced in Sec 4.6.4 of [4] in relation to KPZ, see especially the work by Derrida et al. [17], also with respect to the EW equation [18].

-
- Electronic address: gunnar.pruessner@physics.org;
 URL: <http://www.ma.imperial.ac.uk/~pruess/>
- [1] S. F. Edwards and D. R. Wilkinson, Proc. R. Soc. Lond. A 381, 17 (1982).
 [2] J. Krug and H. Spohn, in Solids far from Equilibrium, edited by C. Godreche (Cambridge University Press, Cambridge, UK, 1991), pp. 479-582.
 [3] T. Halpin-Healy and Y.-C. Zhang, Phys. Rep. 254, 215 (1995).
 [4] J. Krug, Adv. Phys. 46, 139 (1997).
 [5] T. Nattermann and L.-H. Tang, Phys. Rev. A 45, 7156 (1992).
 [6] F. Family and T. Vicsek, J. Phys. A: Math. Gen. 18, L75 (1985).
 [7] J. M. Luck and A. Mehta, Europhys. Lett. 54, 573 (2001).
 [8] P. Biswas, A. Majumdar, A. Mehta, and J. K. Bhattacharjee, Physica A 248, 379 (1998).
 [9] A.-L. Barabasi and H. E. Stanley, Fractal concepts in surface growth (Cambridge University Press, Cambridge, UK, 1995).
 [10] M. Kardar, in Annual Reviews of Computational Physics, edited by D. Stauffer (World Scientific, Singapore, 2001), Vol. V III, pp. 1-47.
 [11] Z. Farkas and T. Fulop, J. Phys. A: Math. Gen. 34, 3191 (2001).
 [12] W. Magnus, F. Oberhettinger, and R. P. Soni, Formulas and Theorems for the Special Functions of Mathematical Physics (Springer-Verlag, Berlin Heidelberg New York, 1966).
 [13] M. Kardar, G. Parisi, and Y.-C. Zhang, Phys. Rev. Lett. 56, 889 (1986).
 [14] H. You, R. P. Chiarello, H. K. Kim, and K. G. Vandervoort, Phys. Rev. Lett. 70, 2900 (1993).
 [15] E. A. Eklund, R. Bruinsma, J. Rudnick, and R. S. Williams, Phys. Rev. Lett. 67, 1759 (1991).
 [16] D. P. Landau and K. Binder, J. Appl. Phys. 63, 3077 (1988).

- [17] B. Derrida, J. L. Lebowitz, E. R. Speer, and H. Spohn, Phys. Rev. Lett. 67, 165 (1991).
- [18] B. Derrida, J. L. Lebowitz, E. R. Speer, and H. Spohn, J. Phys. A :M ath. Gen. 24, 4805 (1991).
- [19] Field-theoretically, the diffusion term might be regarded as dangerously irrelevant.