## Dust Control in Finite A ir Volum es at Zero G ravity – M ean-Field Like A nalysis

T.R.Krishna Mohan and Surajit Sen

Department of Physics, State University of New York, Bu ab, New York 14260-1500<sup>y</sup>

(Dated: April 14, 2024)

We study a simple 1D model of dust rods, with mean size , passing through a parallel 1D alignment of pores as a problem of clogging of a liter by dust grains; is kept less than the pore size, s. We assume that the liter is \sticky", characterized by some parameter 0 1, which means that dust grains slightly smaller in size than s can get trapped in the pores. Our analyses suggest that the number of clogged pores, N<sub>cl</sub>, grows in time as  $t_N^{clog} / N_{cl}$ , where = (;) is a non-universal exponent that depends upon the dust size distribution and liter properties.

PACS num bers: 47.55 K f,47.55 M h,89.20 Bb

Common household dust typically comprises of particulate matter such as carbon particles, skin akes, dust m ites, ber, hair and other irregular shaped ob jects with length-scales between 10  $^2\,$  m and 10  $^2\,$  m (see, for example, [1]). Typically, the larger sized hairs/ bers act as nucleation centers for agglom eration of ne scale dust and the eventual developm ent of the fam iliar dust bunnies. Presence of dust bunnies reduces the e ectiveness of dust lters. W e study the problem of control of dust at all length-scales to ensure that dust bunny grow th becom es di cult. In particular, we consider dust control at zero gravity in an air volum e that is nite and cannot be replenished. Ourmean-eld like analysis is inspired by the problem of air quality maintenance in the International Space Station. The present study has the potential to facilitate the design of lters which can retain air-quality below speci ed thresholds, for long periods.

Signi cant work has been done to describe the pressure drop in uids with ne particulates that is owing through Iters with given speci cations (see, for exam – ple, [2]). The analyses have been performed typically via rate equations using state variables, akin in spirit to therm odynam ical analysis. The resultant blocking equations reveal the behavior of pressure drop with time as Itrate volumes are modi ed and Iter criteria are tuned [3]. Limited information appears to be available on speci cs such as time-dependent degradation of Iters as functions of dust size distributions and Iter pore distributions.

The present work focuses on liter clogging as a function of time. We refrain from addressing the dust agglom eration process in this Letter. Fine scale dust is continuously generated in most inhabited environments and naturally agglom erates. Dust mites ( $10^2$  m), plentiful in most environments, tend to feed on smaller length scale dust. Presence of dust mites therefore leads to a natural integration of smaller length scale dust.

In our analyses, we will be considering dust size distributions such that the m ean dust size is comparable to the

Iterpore size. This turns out to be the regime of interest in which one can understand how Iter and dust properties lead to the time evolution of the clogging process of the Iter. Our studies show that dust size distributions can a lect the details of the clogging process.

The M odel System : The present study is a 1D, m eaneld-like, analysis. Dust grains are irregular in shape and com e in all sizes. In our 1D m odel, we only consider the size of the dust grains, i.e. e ectively consider rod shapes. Consideration of all sizes in 1D is accomplished by assuming a G aussian distribution of dust sizes. Our studies suggest that an exponential tail to the distribution or a Lorentzian distribution will a ect the rate at which liters clog. However, the major qualitative features remain the same for the dimention of and the size considered and, therefore, G aussian distribution will form the basis for our study.

R andom ly sized rods from the distribution, assumed to be aligned vertically for easier geometric conceptualization, are allowed to impinge at random positions on a vertical 1D liter with vertical slits or \pores". The walls between the pores are considered to be of negligiblewidth. Rods of all sizes travelat xed speeds towards the liter. We assume that our time steps are su ciently

ne and/or the air is su ciently dilute in dust content such that a single rod reaching the lter is a reasonable approximation. This justi es measurement of time, t, in term s of particle release events at 'unit' tim e intervals of  $t_i n^{th}$  particle is released at t = nt. Not all particles contribute to clogging. W e will only count clogging events, and m easure clogging rates, with respect to com plete clogging of a pore. The number of clogged pores at any time is denoted as N<sub>cl</sub>. One would expect N<sub>cl</sub> to initially grow rapidly in time. But as time progresses, it will become progressively di cult for the rods to nd the next available pore to clog. Thus, the tem poral grow th of N<sub>cl</sub> will eventually saturate. To best describe the manner in which N<sub>cl</sub> grows, we de ne a new parameter to characterize the instant at which the N<sup>th</sup> clogging event happens and call it  $t_{\rm N}^{\rm clog}$  . This also leads us to the param – eter  $(t_N^{clog}, t_{N-1}^{clog})$ , which is a waiting time between

On leave from CSIR Centre for Mathematical Modelling and Computer Simulation (C-MMACS), Bangalore 560017, India.

<sup>&</sup>lt;sup>y</sup>E lectronic address: km ohan@ bu alo.edu, sen@ dynam ics.physics.bu alo.edu




FIG.1: In Fig.1(a), is kept xed while varying and, in Fig.1(b), vice versa. It is clear from these graphs that  $t_N^{clog}$  / N<sub>cl</sub><sup>(;)</sup> and this is explicitly shown in Fig.1(c), where all the graphs have been collapsed onto a single curve through =  $^{0:62}$  (1)  $^{0:2}$ .

the  $(N = 1)^{th}$  and  $N^{th}$  clogging. A stime progresses and the liter gets progressively clogged, we expect the average value of to grow in magnitude.

The mean of rod size, r, is referred to as and is kept smaller than the lter pore size, s. The standard deviation of rod sizes is taken to be unity. It is to be understood, how ever, that the variance will a ect lter life,

Itering e ciency etc. and will be discussed later. We study cases of Iters with xed pore sizes, as well as Iters with a Gaussian ditsribution of pore sizes; lter pore size is initially kept xed in the results reported below, and is to be considered so unless otherwise mentioned. Ideally, when r < s, rods go through open pores, and, when r > s, they block open pores while, also, partially blocking adjacent pores by the am ount by which r > s. However, we allow a rod with r slightly < s to stick to a pore; this m odels the various sticking tendencies induced by diverse physical mechanisms, and also generates the well docum ented bridge form ation ' [2], where several particles link together to arch over a pore. For this, we de ne a sticking threshold, 0 1, which prescribes the rod length that can block an open pore as a fraction of s. If = 1, only rods with r s can block a pore, and hence this is the hon-stick' lim it. If < 1, particles with r < s will pass through the pore, without sticking. Typically, 0:5

0:9 in our calculations, is set arbitrarily equal to 5 units, and s to 9 units.

We also allow for the form ation of \cakes", another commonly observed phenomenon (see, for example, [2]), with the additional deposition of rods on top of previously deposited rods, which may not contribute to additional blocking unless they extend over adjacent open pores. Rods are not assumed to hit the barrier between pores with any signi cant probability and get trapped by the same; this eliminates the need to consider the partial blocking of adjacent pores by a rod that drapes over the barrier. Further, adjacent pores are considered only to one side (in the direction of increasing array index) for convenience; these e ects do not comprom ise the validity of the nal results. In the following, we present our results on the clogging process of the 1D lter. It may be noted that a 2D liter can be thought of as an array of ad jacent 1D liters. In the lim it of rods being incident at the same rate on each 1D unit' liter, the results of our present study are expected to remain applicable in the 2D case.

Results: W e consider a lter with  $N_{\rm F}$  pores at time  $t=0;\,N_{\rm F}$ , in the following, is kept at  $10^5$  for adequate statistical robustness. In Fig. 1, we plot the time taken for the N  $^{\rm th}$  clogging event,  $t_{\rm N}^{\rm clog}$ , versus the number of clogged pores, N  $_{\rm cl}$ , for various choices of the two control parameters, and , in log-log mode. In Fig. 1(a), is held xed while varying and, in Fig. 1(b), is held

xed while is varied. All the cases are characterized by two di erent regimes of power law growth where, rstly, there is a smooth increase in  $t_N^{clog}$  with N<sub>cl</sub>, followed by an abrupt changeover to a region of in nite slope; the latter marks the divergent times, on the average, required to clog the nalfew pores, which are scattered random ly on the 1D liter. Clearly, the interesting physics is in the

rst region, and we see that  $\frac{c^{log}}{N} / N_{cl}$ , where = (, ). The scaling pattern is made explicit in Fig. 1(c), where we have collapsed all the various graphs for the di erent 's and 's onto a single graph, and it emerges that

$$(;) = (1:0)^{2}:$$
 (1)

where  $_1 = 0.62$  and  $_2 = 0.2$  for this case of G aussian distribution of rod sizes in pinging on a liter with

xed pore size. W e found that these exponents are som ewhat \fragile" and can vary depending on the nature of the rod and liter pore size distributions. For example, with an exponential distribution of rod sizes and a lter of xed pore size, the exponents are  $_1 = 0.2$  and  $_2 = 0.1$ . Nevertheless, the scaling behavior is robust.

Figs.1(a)-(b) reveal that the time required to clog all the pores tend to grow dram atically when N<sub>cl</sub> becomes su ciently large. We recall that the time period between successive cloggings is . In Fig.2, we present data of the distribution of , P (), versus , for two typical cases; the plots are in log-log m ode and only 's up to  $10^4$  are shown. In one case, P () decays smoothly while, in the other




FIG. 2: A plot of the distribution of waiting times, P (), against , in a log-log plot, shows a plateau region preceding the decay region when is reduced; the graphs are with xed pore size, s = 9:0. A similar plateau region results if is reduced, keeping xed. This indicates the enhanced role for sm aller particles in the distribution, which are characterized by sm aller 's, through the incremental clogging mechanism.

case, the decay is preceded by a plateau region where the

's have the same density of occurrence. In general, it was observed that the plateau region increases with relative decrease of with respect to s. A lso, the same e ect is generated if is decreased. In both these cases, we note that the sm aller particles have an enhanced role to play in the clogging process and the plateau region is increased due to the phenom enon that we call increm ental clogging (IC).

0 ne can see that the role of the big (r s) rods is necessary for the initiation of the clogging process; note that these come from the tail of the distribution, have sm aller probabilities and, hence, larger 's associated with them . Once the process has been initiated, the smaller rods can contribute to additional clogging through the IC process, i.e. in the next stage, only a rod of size bigger than the fraction of the remaining open portion of the pore is required to further block the pore and so on. These sm aller rods have larger probabilities and sm aller 's associated with them . A lso, if, at any stage, the rod that sticks on is big enough to protrude into the adjacent pore, it can reduce the size of that pore, which initiates IC in this ad jacent pore. W hile the big rods contribute to clogging when IC is active, their role is especially signi cant in the initial stages of the clogging process. IC increases the proportion of sm all 's and, as a corollary, when the role of sm aller particles is enhanced either by reduction in

or by reduction in , IC becomes a more active mechanism. It is to be emphasized here that the random nature of the clogging events mandates non-zero probability of obtaining small 's even in the nalstages of the clogging process; indeed, this has been clearly seen in plots of 's versus N  $_{\rm cl}$  [4].

At the m acroscopic level, it is of interest to probe the average rate of clogging at the time of clogging of the N  $^{\rm th}$  pore. We de ne this average rate of clogging via  $K_{\rm cl}$ 

 $N_{cl}=t_N^{clog}$ . This quantity is shown in Fig. 3, plotted against  $N_{cl}$ , for various choices of control parameters, and . In Fig. 3(a), is held xed and is varied while, in Fig. 3(b), is held xed and is varied. The data suggests two di erent power-law growth regimes of  $K_{cl}$  as a function of  $N_{cl}$ . There is a smooth increase in  $K_{cl}$  in the early stages followed by a steep decrease in the rate when there are only a few pores left to be clogged; the form er regime corresponds to the functional period in the life of the lter. It turns out that the plots in Fig. 3(a) and Fig. 3(b) can be united, as shown in Fig. 3(c). The scaling relation used on the data in Fig. 3(a) and Fig. 3(b) turns out to be

$$K_{cl} = N_{cl}^{l} :$$
 (2)

where =  $\exp(f(;))$ , and f(;) = 163 + 1.04 + 4.66. The relationship with respect to N<sub>cl</sub> is to be expected given the de nition of K<sub>cl</sub> and Eq. (1). However, further complexity of the scaling relationship via the prefactor results because we are considering a rate quantity in a problem with discontinuous kinetics.

It is easy to calculate (see, for example, [5]), based on the probabilities for particles of size r > k, where

is the standard deviation of rod size distribution and k any positive value, the probability of nding a rod of size r s as against nding a rod of size r s; it turns out that the latter is greater by three orders of m agnitude if = 0.8, with = 5.0, = 1.0 and xed pore size s = 9.0 [4]. Di erences in probability translate to comparable di erences in  $\frac{c^{log}}{M}$ . C learly, variance of the distribution also controls the rate of clogging. Similar di erences in probability can be achieved by varying with respect to s.

If we introduce a distribution of pore sizes around a m ean value, instead of having a xed pore size, the decreases; in this comparison, the same value of the xed pore size was assigned for the mean. If we introduce a second mode at a lower value in a corresponding unim odal G aussian distribution,  $t_{\rm N}^{\rm clog}$  increases because of the enhanced proportion of smaller sized rods; in this com parison, a value of 3.0 was used for the second m ode with the rst mode at 5.0. An exponential distribution, which has a higher proportion of bigger rods than a G aussian, leads to faster clogging of the pores. These results are summarized in Fig. 4 in a log-log plot of  $t_{M}^{clog}$  versus N<sub>cl</sub>. A corresponding ordering, with respect to the distributions, can be observed in plots of K  $_{\rm cl}$  versus N  $_{\rm cl}$ plots; note that larger tim es translate to low er K cl values in this case.

Concluding Remarks: Our studies indicate that adjusting the lter pore size (or mean of the same in case of a pore size distribution) to be slightly higher than the mean of the dust particle distribution is an important step in obtaining e cient ltering as well as longer lter




FIG.3: Plots of K<sub>c1</sub> ( $N_{c1} = t_N^{clog}$ ; cf. Fig. 1) versus  $N_{c1}$ , in log-log mode; in Fig. 3(a), is kept xed while varying and, in Fig. 3(b), vice versa. A scaling relationship akin to Fig. 1(c) m ay be suspected; this is indeed so, as shown by Fig. 3(c), where all the graphs have been collapsed onto a single curve through proper scaling, with  $= \exp(f(;)), f(;) = 16:3+1:04+4:66$ .




FIG. 4: E ect, on clogging rates, of di erent rod size distributions, with and without lter pore size distribution are com pared. The notation in the legend is: BM { B in odalG aussian (two m odes being at 5.0 and 3.0), G { G aussian (m ean at 5.0), E { Exponential (m ean at 5.0); FP { xed pore size (s = 9.0), PD { distribution of pore sizes with m ean at 9.0. It is seen that faster clogging takes place with a pore size distribution in general, and bim odal distribution is the slowest and exponential distribution fastest in clogging the lter.

lifetimes. The variance as well as the type of distribution of the lter and dust particles are also important determ inants. The scaling law that we propose in the relationship of  $t_{\rm N}^{\rm clog}$  and N  $_{\rm cl},$  with respect to the control parameters and , can play an important role in suitably choosing the lter parameters. Even though the scaling exponent is som ew hat \fragile" with respect to di erent distributions of dust particles, the fact that its m agnitude is only altered by a factor of the order of 2, rather than by an order of magnitude, in going from a Gaussian distribution to an exponential distribution, is very encouraging. A key parameter for further analysis is the resistance to the ow caused by lter clogging which requires modelling of uid ow; the relationship of cakes to the model parameters would also be interesting to study. The present study is expected to be applicable to any problem where small particles can lead to clogging of any uid ow. The model may have broad applicability with respect to problem swhere the concept of increm ental clogging can be applied.

A cknow ledgem ents: The research has been supported by NSF CM S0070055. We thank A dam Sokolow and M asam iN akagawa for their interest in this study.

- J.H. van Bronswijk, House Dust Biology, for Allergists, Acarologists and Mycologists (N IB Publishers, Zeist, The Netherlands, 1981).
- [2] R.J.W akem an and E.S.Tarleton, Filtration | Equipment Selection M odelling and Process Simulation (Elsevier A dvanced Technology, Elsevier Science Ltd., 1999).
- P.H.Herm ans and H.L.Bredee, Rec.Trav.Chin.54, 680 (1935); H.P.Grace, AIChE J.2, 307 (1956); M.Shirato, T.Aragaki, and E.Iritani, Chem.Engng.J. (Japan)

12,162 (1979); J.Hermia, Trans.Inst.Chem.E 60,183 (1982); J.G ranger, D.Leclerc, and J.A.Dodds, Filtr. Separ. 22, 58 (1985); M.Hlavaœk and F.Bouchet, J. Membr.Sci.82,285 (1993).

- [4] T.R.Krishna Mohan and Surajit Sen, (Unpublished).
- [5] S. Lipschutz, Schaum 's outline of theory and problem s of probability (M cG raw -H ill, N ew York, 1968).