

Multi-Scaling Comparative Analysis of Time Series and a Discussion on "Earthquake Conversations" in California

Nicola Scafetta¹ and Bruce J. West^{1,2}

¹ Physics Department, Duke University, Durham, NC 27708 and

² Mathematics Division, Army Research Office, Research Triangle Park, NC 27709.

(Dated: March 22, 2024)

Time series are characterized by complex memory and/or distribution patterns. In this letter we show that models obeying to different statistics may equally reproduce some pattern of a time series. In particular we discuss the difference between Levy-walk and fractal Gaussian intermittent signals and show that the adoption of complementary scaling analysis techniques may be useful to distinguish the two cases. Finally, we apply this methodology to the earthquake occurrences in California and suggest the possibility that earthquake occurrences are described by a colored (= "long-range correlated") Generalized Poisson model.

PACS numbers: 91.30.Dk, 05.40.Fb, 05.45.Tp, 89.75.Fb

Herein we introduce a method of multi-scaling comparative analysis (MSCA) for the study of intermittent signals. We show that to distinguish between fractal Gaussian intermittent noise and Levy-walk intermittent noise the scaling results obtained using division entropy analysis (DEA) should be compared with those obtained from both finite variance scaling methods (FVSM) and probability distribution functions (pdf) [1, 2]. Finally, we apply MSCA to the seismic data of California and suggest that, instead of being described by a statistics according to which the waiting times between Omori's earthquake clusters are uncorrelated from one another, as the traditional Generalized Poisson model [3, 4] or a recent Levy-walk like model assume [4], the data may also be characterized by intercluster $1/f$ long-range correlations that may disclose the earthquake conversations recently suggested by Stein [3].

Hurst, in his pioneering work [5], introduced the notion of rescaled range analysis of a time series that takes the scaling form of $(R=S)(t)/t^H$ (H is now called the Hurst exponent). This stimulated Mandelbrot to introduce the concept of fractional Brownian motion (fBm) [6]. In a random walk context the value $H = 0.5$ indicates uncorrelated noise, $0 < H < 0.5$ indicates anti-persistent noise and $0.5 < H < 1$ indicates persistent/long-range correlated noise [6]. Alternative scaling methods applied to a time series f_{ig} , where $i = 1, 2, \dots$, focus on the autocorrelation function $C(t)/h_{i+i+1}/t^{2H}$; on the power spectrum representation $P_S(f)/f^{1-2H}$ [7] and on the evaluation of the variance of the division generated by f_{ig} [8] ($V(t)/t^{2H}$). All such scaling methods are related to the original Hurst's analysis and yield his H -exponent. These techniques, referred by us as FVSM, assume a finite variance and according to the central limit theorem (CLT) [9] the underlying statistics are Gaussian.

Recently, Scafetta et al. [1] introduced a complementary scaling analysis, the DEA, that focuses on the scaling exponent evaluated through the Shannon entropy $S(t)$ of the division generated by the fluctuations f_{ig}

of a time series [1, 2]. Here, the pdf of the division process, $p(x;t)$, is evaluated by means of the subtrajectories $x_n(t) = \sum_{i=0}^t x_{i+n}$ with $n = 0, 1, \dots$. The pdf scaling property for a fractal anomalous division takes the form $p(x;t) = t^{-F} x^{-t}$; and its entropy increases in time as $S(t) = \int_0^1 p(x;t) \ln[p(x;t)] dx = A + \ln(t)$; where A is a constant. One can also examine the scaling properties of the second moment for the same process using the FVSM. One version of FVSM is the standard deviation analysis (SDA) [2] which is based on the evaluation of the standard deviation $D(t)$ of the same variable x and pdf $p(x;t)$, and yields $D(t) = \sqrt{\int_0^1 x^2 p(x;t) dx} / t^H$ [2, 6].

Note that the entropy $S(t)$ does not require the variance of the pdf $p(x;t)$ to be finite [2]. The existence of scaling for a process with a diverging second moment implies that DEA is complementary to and not simply an alternative to FVSM. So, the scaling exponent is conceptually different from the Hurst exponent H measured by means of the FVSM. This suggests that the scaling exponents and H may fulfill multiple relations according to the process under study and, therefore, the combined use of DEA, SDA and pdf analysis may increase our understanding of complex phenomena through a MSCA.

Herein we focus on the statistics of intermittent noises. The simplest way to represent intermittent noise f_{ig} is through a dichotomous representation in which the value $= 1$ indicates the occurrence of an event and the value $= 0$ represents no-event [10, 11, 12, 13]. An intermittent noise is characterized by the correlation properties of the waiting time sequence f_{ig} between consecutive events and by its waiting time distribution (τ) . There are two basic distinct forms of intermittent noises:

1) Fractal Gaussian intermittent noise is characterized by a long-range correlated waiting time sequence, h_{i+i+1}/t^{2H} , and by a finite variance waiting time distribution (τ) whose form may be, for example, that of a Gaussian, exponential or Poisson distribution. The division generated by a fractal Gaussian intermittent noise is a particular type of fBm and satisfies the asym-




FIG. 1: DEA and SDA of [A] a fractal Gaussian intermittent noise with $\langle t \rangle / \exp(\langle t \rangle) = 25$ and $H = 0.75$; the fractal Gaussian relation (1) of equal exponents is fulfilled; [B] a Levy-walk intermittent noise with $\langle t \rangle / \exp(\langle t \rangle) = 2.5$; note the bifurcation between $H = 0.75$ and $\delta = 0.67$ caused by the Levy-walk relation (2).

otic scaling relation between indices

$$\delta = H : \quad (1)$$

We refer to (1) as the fractal Gaussian diffusion relation. If the long-range correlations of f_{ij} are destroyed via shuffling, the new intermittent sequence is characterized by the value $H = 0.5$ of random time series. Fig. 1A shows the scaling properties of a computer generated fractal Gaussian intermittent noise with an exponential waiting time distribution and $H = 0.75$.

2) Levy-walk intermittent noise is characterized by an uncorrelated waiting time sequence, h_{ij} / t_{ij} , and a Levy or an inverse power law waiting time distribution $\langle t \rangle / (T + 1)$; with $2 < \alpha < 3$ that ensures that although the first moment of t is finite, the second moment diverges. The presence of a Levy-walk process in a given time series can be detected [2, 10] by mean of the following asymptotic relation among the three exponents

$$0.5 < \delta = (3 - 2H)^{-1} = (\alpha - 1)^{-1} < 1 : \quad (2)$$

We refer to (2) as the Levy-walk diffusion relation. Interesting applications of this type of noise have been found in several Levy phenomena including the distribution of solar flares [10, 11, 12, 13, 14]. In the particular case in which the sequence f_{ij} is correlated the scaling exponents δ and H are larger than the values predicted by Eq. (2). Fig. 1B shows the scaling properties of a computer generated random Levy-walk intermittent noise with $\langle t \rangle = 2.5$ that has $H = 0.75$ and $\delta = 0.67$.

We stress that the Levy-walk relation (2) is fulfilled if the waiting times f_{ij} are uncorrelated, in which case any shuffling of f_{ij} would not alter the scaling exponents H and δ . In fact, the super-diffusion scaling exponents $0.5 < \delta < H < 1$ of a Levy-walk intermittent noise are related to the fatness of the waiting time inverse power law tail, as measured by the exponent α . Contrary to a fractal Gaussian intermittent noise, this Levy scaling does not imply a temporal correlation, or a historical

memory, among events because the occurrence of future events is independent of the frequency of past events.

We also observe that there exist particular intermittent sequences obtained by mixing Levy and Gaussian noises [12], with a Levy memory beyond memory [15] or by substituting an event with a cluster of events [4]. In these cases the asymptotic properties of the scaling exponent H and δ are expected to depend on the component, Gaussian or Levy, with the strongest persistence.

The relations (1) and (2), and the correlation/shuffling effects indicate that the DEA should be jointly used with the FVSM and/or pdfs. The adoption of a single technique can lead to a misinterpretation of the characteristics of a phenomenon, because Levy-walk intermittent noise can be confused with fractal Gaussian intermittent noise, and uncorrelated noise of one kind of statistics can be mistaken for correlated noise with another kind of statistics. Figs. 1A and 1B clearly show that the determination of only one of the two exponents H and δ is not sufficient to conclude whether a phenomenon is characterized by a Levy-walk intermittent statistics or by a fractal Gaussian intermittent statistics. So, we suggest a MSCA by combining complementary techniques.

Recently, DEA has been applied by Mega et al. [4] to study the time distribution of earthquakes in Southern California (20° - 45° N latitude and 100° - 125° W longitude) from 1976 to 2002. The catalog [16] is complete for local events with magnitude $M \geq 3$ since 1932, for $M \geq 1.8$ since 1981 and for $M \geq 0$ since 1984. The time intervals between large earthquakes was studied in Ref. [4] by setting a temporal variable $t_i = 1$ at the occurrence of an earthquake with a magnitude larger than a given threshold M_t , and by setting $t_i = 0$ when no earthquake of the specified magnitude occurs. We refer to f_{ij} as the waiting time sequence between consecutive earthquakes with $M \geq M_t$. So, the intermittent sequence t_i was analyzed by means of the DEA and the measured scaling exponent was $\delta = 0.94 \pm 0.01$. The authors of Ref. [4] concluded that the time intervals, t_i , between two consecutive Omori's earthquake clusters [17] is modeled by an inverse power law $\langle t_i \rangle / \langle t_i \rangle$ with an exponent $\alpha = 2.06$ calculated via Eq. (2). This calculation was based on the traditional assumption [4] that the waiting times between such clusters are uncorrelated ($\langle t_i \rangle \langle t_j \rangle = t_{ij}$), implying that the observed superdiffusion is induced by a Levy-walk between the Omori's clusters. Finally, Mega et al. [4] showed that a synthetic sequence produced with Omori's uncorrelated clusters, $\langle t_i \rangle \langle t_j \rangle = t_{ij}$, temporally distributed according to an inverse power law $\langle t_i \rangle / \langle t_i \rangle$ with $\alpha = 2.06$, generates a superdiffusive process with $\delta = 0.94$.

However, the authors of Ref. [4] did not make the important distinction between Levy-walk and fractal Gaussian intermittent noises such as we did above. We showed that a scaling exponent in the range $0.5 < \delta < 1$ can be




FIG. 2: pdf of the waiting times τ of earthquakes with a magnitude $M_t = 1, 2, 3$ and 4. The initial $P(\tau)/1 =$ is the Omori's law [17].

associated with either a correlated fractal Gaussian intermittent noise or with an uncorrelated Levy-walk noise. Consequently we apply a M SCA to determine which of the two statistics better describes the data.

Fig. 2 shows the waiting time pdfs between earthquakes using four magnitude thresholds $M_t = 1, 2, 3$ and 4. The pdfs show an initial Omori's law [17] ($P(\tau)/1 =$), but the pdf tails present a large inverse power law exponent > 4 and may even approach an exponential (or Poisson) distribution asymptotically. The Omori's law is determined by the short-range correlated aftershocks [17] and lasts for a time that increases with the magnitude threshold. If the waiting time distribution between Omori's clusters were an inverse power law with $\alpha = 2.06$ it might be expected that by increasing the magnitude threshold, most of the aftershocks could be cut off and the tail of the distribution could converge to an inverse power law with $\alpha = 2.06$. This does not seem to happen. Therefore, such an $\alpha = 2.06$ inverse power law, if it is real, cannot be observed in this way.

Fig. 3 compares the DEA and SDA applied to the intermittent sequence (t) of earthquakes with $M_t = 1$. Different magnitude thresholds give similar results. If these data corresponded to a random intermittent Levy-walk and if the curves shown in the figure corresponded to the asymptotic regime, the condition (2) interrelating the exponents should hold. A rigorous DEA fit in the range $2^7 : 2^{15}$ gives $\delta = 0.944 \pm 0.008$ implying a Levy-walk $H = 0.97 \pm 0.005$. Instead, the same-range SDA fit gives $H = 0.943 \pm 0.004$. The error analysis seems to confirm better the Gaussian relation of equal scaling exponents (1) because δ and H overlap within the statistical error as in Fig. 1A, while the difference between the measured H and the Levy-walk H is statistically significant ($p < 0.01$). By shuffling the earthquake waiting time intervals f_{ig} we get $H = 0.5$. Finally, by directly applying DEA




FIG. 3: DEA and SDA of the intermittent time signal (t) for magnitude $M_t = 1$. The data are fitted with scaling exponents $\delta = 0.944 \pm 0.008$ and $H = 0.943 \pm 0.004$. The uppermost solid line with $H = 0.97$ corresponds to the expectation of H if the Levy-walk condition (2) holds true.

and SDA to the waiting time series f_{ig} , we again get $\delta = H = 0.94$. These findings suggest that the data do not fulfill the Levy-walk relation (2) and that it might be more likely that the Californian earthquakes are long-range temporal correlated according to the persistence of a fractal Gaussian intermittent noise with $H = 1$ known as 1=f or pink noise [7].

The curve with circles in Fig. 4 shows the DEA applied to a synthetic earthquake catalog obtained by coloring a kind of Generalized Poisson model for earthquakes. First we generated several Omori's clusters exactly as done in Ref. [4], that is, by assuming that the number of earthquakes in a cluster follows an inverse power law distribution with exponent equal to 2.5 and that the events within the same cluster are temporally distributed according to the Omori's law, that is, an inverse power law with exponent equal to $p = 1$. We generated a total number of events equal to the total number of earthquakes in the catalog with a magnitude threshold $M_t = 1$. However, contrary to what was done in Ref. [4], we do not randomly $(h_i^{[m]}, j_i^{[m]})$ position these clusters according to an inverse power law intercluster waiting time distribution $(h_i^{[m]})/1 =$ with $\alpha = 2.06$. Instead, we distribute the clusters according to a 1=f fractal Gaussian intermittent noise f_{ig} that is, with $h_i^{[m]} h_{i+t}^{[m]} / t^{2H-2}$ and $H = 1$. The intercluster waiting time distribution $(h_i^{[m]}) / \exp(h_i^{[m]})$ shown in the insert of Fig. 4 could be substituted with any other distribution with finite variance. Fig. 4 shows that the model is able to reproduce the same superdiffusion pattern shown by the data. Finally, the curve with triangles shows the reduction of long-range persistency of a synthetic catalog obtained with the same clusters of above but temporally distributed after having shuffled, to randomize, the same




FIG. 4: DEA applied to a 1=f long-range correlated Generalized Poisson model (circle) for earthquakes for $M < 1$. The intercluster waiting time pdf ($V(t)$) is exponential and $h_i^{[m]} h_j^{[m]} / t^{2H-2}$ with $H < 1$. The curve with triangles refers to the case in which the intercluster waiting time sequence $f_j^{[m]}$ is randomized such that $h_i^{[m]} h_j^{[m]} / t_{ij}$.

intercluster waiting time sequence $f_j^{[m]}$.

However, Eqs. (1) and (2) are fulfilled only asymptotically where the Central Limit Theorem for Gaussian processes or its generalization for Levy processes apply [9]. There might be the possibility that Fig. 3 as well as figure 2 in Ref. [4] do not show the asymptotic limit but a transition regime that is strongly superdiffusive ($H < 1$) because of the Omori's intracluster short-range correlations. A long transition regime is also evident in the curve with triangles shown in Fig. 4 that refers to an uncorrelated intercluster waiting time sequence and should asymptotically converge to $H = 0.5$. In fact, also a random Generalized Poisson model or an "ETAS" model [18] with appropriate parameters may generate a synthetic catalog showing superdiffusive properties similar to the real data within a time range [19]. However, we observe that such a result may depend too strongly on the particular parameters used in the models.

In conclusion, we have discussed some of the difficulties that can be encountered in interpreting intermittent sequences and shown that models with alternative statistics can reproduce some pattern of a time series equally well. This fact suggests the need of an analysis involving complementary tests. In particular we showed how to distinguish fractal Gaussian intermittent noise from Levy-walk intermittent noise using M-SCA. This methodology has important application in the analysis of phenomena having intermittent signals because different statistics imply different dynamics. Our analysis supports the idea that earthquakes generate strain diffusion, whose propagation over hundreds of kilometers induces remote seismic activity [3, 20]. This propagation according to our statistical analysis produces correlations in the time intervals between earthquake clusters. In fact, the thesis that earth-

quakes are assembled into uncorrelated Omori's clusters, $h_i^{[m]} h_j^{[m]} / t_{ij} = 1/t_{ij}$, as both the standard Generalized Poisson model [4] and the Levy-walk model [4] require, seems unrealistic. We suggest that it is more plausible that earthquake clusters are 1=f long-range correlated and, perhaps, they are subclusters of a larger Omori cluster. In fact, a 1=f noise can be generated by the superposition of relaxation processes within a wide range of energies [7] that may well describe the coexistent stress alterations caused by old and recent, as well as large and small shocks. Thus, the 1=f long-range correlations may imply that earthquake occurrences may strongly depend on the geological history of a vast region.

Acknowledgment: N.S. thanks the ARO for support under grant DAAG 5598D 0002.

-
- [1] N. Scafetta, P. Hamilton, and P. Grigolini, *Fractals* 9, 193 (2001).
 - [2] N. Scafetta, P. Grigolini, *Phys. Rev. E* 66, 036130 (2002).
 - [3] R. S. Stein, *Scientific American*, 288 Jan 72 (2003).
 - [4] M. S. Mega, P. Allegrini, P. Grigolini, V. Latora, L. Palatella, A. Rapisarda and S. Vinciguerra, *Phys. Rev. Lett.* 90 188501 (2003).
 - [5] H. E. Hurst, R. P. Black, and Y. M. Simaika, *Long Term Storage: An Experimental Study* (Constable, London, 1965).
 - [6] B. B. Mandelbrot, *The Fractal Geometry of Nature*, (Freeman, New York, 1983).
 - [7] R. Badii and A. Politi, *Complexity, Hierarchical structures and scaling in physics*, (Cambridge University Press, UK 1997).
 - [8] C.-K. Peng, S. V. Buldyrev, S. Havlin, M. Simons, H. E. Stanley, and A. L. Goldberger, *Phys. Rev. E* 49, 1685 (1994).
 - [9] B. V. Gnedenko and A. N. Kolmogorov, *Limit Distributions for Sums of Random Variables*, Addison-Wesley, Reading, MA, (1954).
 - [10] P. Grigolini, L. Palatella, and G. Raelli, *Fractals* 9, 439 (2001).
 - [11] P. Grigolini, D. Leddon, N. Scafetta, *Phys. Rev. E* 65, 046203 (2002).
 - [12] N. Scafetta, V. Latora, P. Grigolini, *Phys. Rev. E* 66, 031906 (2002).
 - [13] N. Scafetta and B. J. West, *Phys. Rev. Lett.* 90, 248701 (2003).
 - [14] M. F. Shlesinger, B. J. West and J. Klafter, *Phys. Rev. Lett.* 58, 1100-1103 (1987).
 - [15] P. Allegrini, P. Grigolini, P. Hamilton, L. Palatella and G. Raelli, *Phys. Rev. E* 65 41926 (2002).
 - [16] Southern California Earthquake Data Center. <http://www.scecdc.scec.org/ftp/catalogs/SCNS/>
 - [17] F. Omori, *J. College Sci. Imper. Univ. Tokyo* 7, 111 (1895). P. Bak, K. Christensen, L. Danon and T. Scanlon *Phys. Rev. Lett.* 88 178501 (2002).
 - [18] Y. Y. Kagan and L. Knopov, *Science* 236, 1563 (1987).
 - [19] A. Helmstetter and D. Sornette, *physics/0307134 v1*.
 - [20] A. Helmstetter, *Phys. Rev. Lett.* 91, 058501 (2003).