

Neel-Dimer Transition in Antiferromagnetic Heisenberg Model and Deconfinement of Spinons at the Critical Point

Daisuke Yoshioka, Gaku Arakawa and Ikuo Ichinose
Department of Applied Physics, Graduate School of Engineering,
Nagoya Institute of Technology, Nagoya, 466-8555 Japan

Tetsuo Matsui
Department of Physics, Kinki University, Higashi-Osaka, 577-8502 Japan
(April 14, 2024)

Quantum phase transition from the Neel to the dimer states in an antiferromagnetic (AF) Heisenberg model on square lattice is studied. We introduce a control parameter for the exchange coupling which connects the Neel ($\gamma = 0$) and the dimer ($\gamma = 1$) states. We employ the CP^1 representation of the $s = \frac{1}{2}$ spin operator and integrate out the half of the CP^1 variables at odd sites to obtain a CP^1 nonlinear model. The effective coupling constant is a function of γ and at $\gamma = 0$ the CP^1 model is in the ordered phase which corresponds to the Neel state of the AF Heisenberg model. A phase transition to the dimer state occurs at a certain critical value of γ as γ increases. In the Neel state, the dynamical composite $U(1)$ gauge field in the CP^1 model is in a Higgs phase and low-energy excitations are gapless spin wave. In the dimer phase, a confinement phase of the gauge theory with $s = 1$ excitations is realized. For the critical point, we argue that a deconfinement phase, which is similar to the Coulomb phase in 3 spatial dimensions, is realized and $s = \frac{1}{2}$ spinons appear as low-energy excitations.

Quantum phase transition (QPT) is one of the most interesting problem in these days.¹ It is often argued that the simple Ginzburg-Landau theory does not apply to certain class of the QPT's. In this paper we shall study $s = \frac{1}{2}$ antiferromagnetic (AF) Heisenberg model on 2-dimensional square lattice with nonuniform exchange couplings,

$$H_{AF} = \sum_{x,j} J_{xj} S_x \cdot S_{x+j} \quad (1)$$

where x denotes site of the spatial lattice, j is the direction index ($j = 1; 2$) and S_x is the spin operator at site x . We rename the even lattice sites $x = (o; i)$ where o denotes odd site and the index $i = 1; 2; 3$ and 4 specifies its four nearest-neighbor (NN) even sites (see Fig.1).




FIG. 1. Numbering and dimer picture: cross symbol is odd site, and spot symbol is even site. Solid line means that its correlation is stronger than dotted lines.

The exchange couplings $J_{xj} = J_{oi}$ are position dependent and we explicitly consider the following case which corresponds to the dimer configuration,

$$\begin{aligned} J_{oi} &= J + J_{oi}; \\ J_{oi} &= \begin{cases} J_{oi} = J; & i = 1 \\ J_{oi} = J; & i = 2; 3; 4 \end{cases} \end{aligned} \quad (2)$$

where $0 \leq \gamma \leq 1$ is a control parameter which connects the uniform Heisenberg model to the dimer model.

It is not so difficult to derive the CP^1 field-theory model^{2,3} from Eq.(1)⁴. The spin operator S_x can be expressed in terms of the CP^1 variable $z_x = (z_x^1; z_x^2)^t$ as

$$S_x = \frac{1}{2} z_x^y \sim z_x; \quad (3)$$

where \sim are the Pauli matrices and the CP^1 constraint $\sum_{a=1,2} z_x^a z_x^a = 1$ guarantees the magnitude of the localized spin as $\frac{1}{2}$.

From our assignment of J_{oi} (2), it is obvious that J_{o1} is larger than the others. We use the path-integral formalism and parameterize the CP^1 variable z_o by referring to $z_{o1} = z_e$,

$$z_o = p_o z_e + \frac{1}{1 - p_o} e^{i\phi} z_e; \quad (4)$$

where p_o is a parameter, $e^{i\phi}$ is a phase factor and $z_e = i \frac{1}{2} z_e$. At vanishing temperature (T), spins tend to point antiparallel their NN spins, and then the parameter p_o can be treated as a small parameter. We expand $\frac{1}{1 - p_o} \approx 1 + \frac{1}{2} p_o + \dots$ and retain only terms up to quadratic of p_o . Then we perform the Gaussian integration of p_o 's to obtain an effective model of z_e 's for which smooth configurations dominate at $T = 0$.

Calculation is rather long but straightforward⁴ and we obtain an effective field theory of the AF Heisenberg model under study,

$$L_{CP} = \sum_x \sum_h \left(\frac{1}{2} z_x^a z_x^a + \frac{1}{2} z_x^a z_{x+h}^a \right) + \frac{1}{f_e} \sum_x \frac{1}{f_e} + L_B; \quad (5)$$

where $D_z = (\partial + iA)_z$, $A = iz\partial_z$ is the $U(1)$ gauge field, λ_x is the Lagrange multiplier for the CP^1 constraint and L_B is the Berry phase term. We have rescaled the imaginary time and x as taking the continuum limit (the spatial coordinate x in Eq.(5) denotes the even site of the original lattice). The effective coupling constant f_e and the "speed of light" c , which is often set unity, are explicitly given as

$$\frac{1}{f_e} = \frac{1}{2} \frac{1}{\bar{a}} \frac{1}{2} \frac{r}{2(2+r)} \frac{1}{1} \frac{1}{f_{AFH} b(r)}; \quad (6)$$

$$c = \frac{2aJ}{h} \frac{(2+r)(1+r)}{2} c_{AFH}(r); \quad (7)$$

where a is the lattice spacing of the original lattice and $f_{AFH} = 2 \frac{1}{\bar{a}}$ and $c_{AFH} = \frac{2aJ}{h}$.

Here we should comment on the Berry phase terms, L_B , which appear in the CP^1 nonlinear model representation of the AF Heisenberg model. First we consider AF spin chains in one dimension in order to obtain important insight for the effect of the Berry phase terms. For the uniform AF chains, the Berry phase reduces to the θ -term ($i \int d\mathbf{x} F_x$) with $\theta = \pi$. The gauge dynamics crucially depends on the value of θ and only the case of $\theta = (\text{mod } 2\pi)$, the deconfinement phase is realized. Topologically nontrivial configurations are suppressed by the θ -term when $\theta = \pi$.

On the other hand for the bond-alternating (BA) AF Heisenberg chain with $J_x = J + (-1)^x J$, the value of the parameter is $\theta = \frac{J_+ - J_-}{J_+ + J_-}$, and therefore the above suppression by the θ -term does not occur and gauge dynamics is essentially the same with the $\theta = 0$ case. Then the confinement of the spinon z_x^a occurs, and low-energy excitations are gapful $s = 1$ excitations $\pi_x = \frac{1}{2} z_x^y \sim z_x$ even for a very small value of J_- . This is nothing but the spin-Peierls transition.

For the 2-dimensional uniform AF Heisenberg model, the Berry phase term similarly appears as $L_B = \frac{i}{2} \sum_x A_x$, where $x = 1(1)$ for even (odd) site x (of the lattice with the lattice spacing \bar{a} for the CP^1 model (5)) and A_x is the area enclosed by the curve given by the time evolution of $\pi_x(t)$. It can be proved that it gives only vanishing contribution for an arbitrary smooth configuration. However for singular configurations like instantons (monopoles), the Berry phase is nonvanishing and it gives terms like $L_B = i \sum_{i=1}^4 \frac{1}{2} \pi_x^i m_x^i$; where π_x^i are $0; 1; 2; 3$ ($i = 1; 2; 3; 4$), respectively and m_x^i is the instanton number.^{5,6} (The index i refers to the four dual sublattices of the original lattice.) From the above L_B , it is obvious that the cancellation mechanism of the instanton contribution occurs by the destructive interference unless $m_x^i = 4 (\text{mod } 4)$.

For the nonuniform cases which are considered in this paper, the coefficient of the Berry phase term becomes irrational as in the case of the BA spin chains and therefore cancellation of instanton effect does not occur or at least weakens. Then we can expect that the (global) phase

structure of the effective gauge model for the nonuniform AF Heisenberg model is the same with that of the CP^1 model without the Berry phase terms. More comments will be given after showing results of nonperturbative studies on the gauge dynamics.

It is straightforward to obtain the effective potential of π_x and $z^2 = h z_x^2$ by integrating out z_x^1 ,

$$V_e = \int d^2x \int \frac{1}{f_e} + \frac{1}{6} \left(\frac{1}{2} \right)^p \frac{1}{2} \frac{1}{3} \frac{1}{p-1};$$

where the cutoff $\Lambda = \frac{p}{a}$. The effective potential V_e indicates that there exists a critical coupling $f_c = \frac{4}{3}$. The existence of the phase transition has been verified by the numerical calculation of the equivalent $O(3)$ nonlinear model in $(2+1)$ dimensions⁷. In the weak-coupling region $f_e < f_c$, the spontaneous symmetry breaking occurs and $h z_x^2 \neq 0$. As a result, the Higgs phase is realized in the gauge-theory terminology. Low-energy excitations are gapless spin waves which are described by z_x^1 . In the strong-coupling phase $f_e > f_c$, $h z_x^2 \neq 0$ whereas $h z_x^2 = 0$. Local Maxwell terms appear in the effective action of the gauge field A , and the confinement phase is realized. Low-energy excitations are $s = 1$ composite of the spinons which correspond to π_x with $(\text{mass})^2 / h x$.

The effective coupling f_e in (6) first decreases as θ increases but above certain value of θ it starts to increase and goes to infinity at the dimer limit $\theta = 1$. In the uniform case $\theta = 0$, $f_e(\theta = 0) < f_c$ and this means that the ordered Neel state is realized at the vanishing T in the AF Heisenberg model in two spatial dimensions as it is now widely believed. The behavior of the effective coupling $f_e(\theta)$ shows the existence of a critical value f_c at which the phase transition occurs. This result indicates that the strong-coupling phase of the CP^1 model (5) corresponds to the dimer phase in which the ground state is nothing but spin-singlet pairs formed by the alternative strong exchange couplings and excitations have $s = 1$. In fact this Neel-dimer transition was observed by the numerical calculations some years ago⁸.

Hereafter we are interested in the critical point at $f_e = f_c$ which separates the Neel and dimer phases. In order to investigate that "phase", study on the gauge dynamics is required. At f_c , $h z_x^2 = h z_x^a = 0$. Effective action of the gauge field and the field π_x is obtained by integrating out the gapless spinon field z_x^a ($a = 1; 2$). The resultant effective action becomes nonlocal and therefore it is possible for the gauge dynamics to belong to different universality class from that of the usual gauge theory in $2+1$ dimensions. By the continuum field-theory calculation, the effective action of the gauge field A is obtained as

$$L_A / \int d^3x \int d^3y \frac{1}{F(x)} \frac{1}{F(y)}; \quad (8)$$

where $F = \partial A - \partial A$. Similarly the effective action of the field π_x is obtained as

$$\frac{1}{L} \int d^3p \sim(p) \frac{1}{p_j} \sim(p); \quad (9)$$

where $\sim(p)$ is the Fourier transformed field of ϕ_x . Equation (9) shows that fluctuations of the field ϕ_x are strongly suppressed at large distances. In the CP¹ model on the 3-dimensional space-time cubic lattice, a similar expression of the action L_A and L is obtained by the hopping expansion of z_x^a ($a = 1; 2$). In the effective action S_A of the compact gauge field $U = e^{iA}$, the following nonlocal terms appear

$$S_A = \sum_{\mathbf{x}} \sum_{\mathbf{y}} \sum_{\mathbf{z}} U(\mathbf{x}; \mathbf{y}, \mathbf{z}); \quad (10)$$

where the summation over closed loops includes loops of an arbitrary large size, j is the length of and the parameter is estimated as $\frac{1}{2d}$ for massless z_x^a with the dimension of the lattice $d = 3$. We shall focus our interest on the gauge dynamics of the above nonlocal action which is one of the most important problems in the theoretical studies on the strongly-correlated electron systems in these days and is still controversial. At present it is known that there exists only the confinement phase in the (2 + 1)-dimensional compact U(1) gauge theory without matter couplings.⁹ However phase structure is not clear when the U(1) gauge field couples to matter fields, particularly gapless matter fields^{10,14}. In particular in Ref.^{10,11}, it is argued that a deconfinement phase is realized by the gapless fermion couplings. Simple mean-field type argument is not applicable for the nonlocal gauge systems and numerical studies are required.

We shall study lattice gauge model with a nonlocal action which is related with (10) but slightly more tractable. The summation over \mathbf{z} in S_A (10) becomes (logarithmically) divergent for the condensation $U = 1$ ¹⁵ since the massless relativistic bosons z_x^a ($a = 1; 2$), which appear at the critical point, give divergent hopping expansion for $U = 1$. From the above discussion we shall consider the following (2 + 1) dimensional lattice gauge model with the long-range interaction in the \hat{x} direction,

$$S_G = g_1 \sum_{\mathbf{x}} \sum_{\mathbf{y}} \sum_{\mathbf{z}} \frac{1}{n} U(\mathbf{x}) W_{\mathbf{x}+\mathbf{z}}(n) U^\dagger(\mathbf{x}+\mathbf{z}) W_{\mathbf{x}}^\dagger(n) + g_2 \sum_{\mathbf{x}} \sum_{\mathbf{y}} U(\mathbf{x}); \quad (11)$$

where $W_{\mathbf{x}}(n) = U_0(\mathbf{x}) U_0(\mathbf{x} + \hat{x}) \dots U_0(\mathbf{x} + n\hat{x})$, N is the system size in the \hat{x} direction and g_i ($i = 1; 2$) are coupling constants for the time and spatial directions, respectively. From action (11), it is obvious that the gauge model under study has nonlocal coupling in the \hat{x} direction whereas it has the usual local Maxwell-type correlation in the spatial directions.¹⁶ Reason why we take the action (11) is that the \hat{x} direction terms logarithmically divergent for $U = 1$ and also Monte-Carlo simulations are easier for the model than those with full-nonlocal interaction terms. We think that studies of the model (11)

give important insight for the full-nonlocal gauge system (10). More comments on this point will be given after showing the results of the Monte-Carlo simulations of the model (11).




FIG. 2. Fluctuation of energy as a function of g_1 is plotted for the fixed $g_2 = 1.0$ and 3.0 . Lattice size is $12^2 \times 16$ and $9^2 \times 12$. The results show the existence of a phase transition.




FIG. 3. The correlations of Polyakov lines as functions of their distance are plotted. The results for $g_1 = 0.12$ and 0.14 are shown in (a), and one for $g_1 = 0.16$ is shown in (b). Value of g_2 is fixed as $g_2 = 1.0$. The results show the existence of the confinement-deconfinement phase transition.

We studied phase structure of the model (11) by the Monte-Carlo simulations of the standard Metropolis algorithm. We calculated the fluctuation of the energy E , i.e., $\langle (E - \langle E \rangle)^2 \rangle^{1/2}$, as a function of the coupling g_1 for fixed g_2 . The results are shown in Fig. 2 for $g_2 = 1.0$ and 3.0 , and they indicate the existence of a phase transition at critical coupling $g_{1c} \approx 0.14$ and $g_{1c} \approx 0.13$, respectively. We varied the system size and verified that the peak gets sharper as the system size increases. This means that the transition is of second order. In order to investigate gauge dynamics in each phase we calculated the correlations of the Polyakov lines $G(l) = \langle P(\mathbf{x}) P^\dagger(\mathbf{x} + l\hat{x}) \rangle$ where $P(\mathbf{x}) = U_0(\mathbf{x}) U_0(\mathbf{x} + \hat{x}) \dots U_0(\mathbf{x} + N\hat{x})$.¹⁷ In Fig. 3, we show the results of $G(l)$ for various values of g_1 and it is obvious that the behavior of $G(l)$ changes at $g_1 = g_{1c}$ from $\ln G(l) \sim -l$ to $G(l) \sim \text{constant}$ for large l . Then the phase transition from the confinement to the deconfinement phases occurs as g_1 increases.

Numerical calculations show that the spatial coupling of the gauge field and larger system size enhance the deconfinement phase. From the above results we expect that the full-nonlocal gauge model (10), which results from the integration of the massless relativistic boson z_x^a at the criticality, is in the deconfinement phase. In fact

the coupling constant $g_1 = 1$ corresponds to (10) in which the damping factor γ_j balances the entropy factor of the paths.

In the deconfinement phase of (11), topologically non-trivial configurations are suppressed and the field-theory result (8) gives a qualitatively correct picture. Charges interacting through A have the potential $V(r) \propto 1/r$ where r is the spatial distance between the two charges.

Let us comment on the effects of the Berry phase. Since L_B is neglected in (11), one may doubt the deconfinement phase transition observed above. However, as the Berry phase generates extra phases for topologically nontrivial configurations in the path integral, the Berry phase enhances the deconfinement. In fact without these extra phases, all instanton configurations contribute additively to disorder the gauge system. Thus the existence of the deconfinement phase in the gauge system (11) guarantees its existence even in the presence of the Berry phase. Similar argument was used for the deconfinement transition at finite T ¹⁸, which is established at present.

We summarize the phase structure of the original spin model. In the region $\beta < \beta_c$ ($f_e < f_c$), the low-energy excitations are the massless spin waves whereas in the region $\beta > \beta_c$ ($f_e > f_c$), they are $s = 1$ excitations π_x . On the critical point $\beta = \beta_c$ ($f_e = f_c$), the gauge dynamics is in the "Coulomb" phase and the low-energy excitations are the $s = 1/2$ bosonic spinons z_x^a ($a = 1, 2$) which are interacting with each other by the potential $1/r$. The spin correlation function decays algebraically both in the Neel state and at the criticality but exponent is different. In the Neel state, the spin operator is given as $S_x = \frac{1}{2} z_x^y \sim z_x^y - i z_x^x \sim z_x^1$ whereas at the criticality π_x , the bilinear of z_x^a and $z_x^{a\dagger}$ ($a = 1, 2$). Phase structure of the nonuniform AF Heisenberg model is schematically shown in Fig. 4.




FIG. 4. Schematic phase diagram of the Heisenberg model. Phases A, B and C are the Higgs (Neel), Coulomb (critical) and confinement (dimer) phases, respectively.

Recently a similar phase transition from the Neel to the dimer states was discussed¹⁹. There they conclude that instanton effects are irrelevant at the critical point. Our numerical investigation is consistent with their result but our study shows the long-range interactions of the gauge field, which appear as a result of the coupling to the massless boson z_x^a , play an essentially important role. Results of more detailed studies on the long-range gauge

theories will be reported in near future.²⁰

- ¹ See for example, S. Sachdev, "Quantum Phase Transitions", (Cambridge University Press, Cambridge, England, 1999).
- ² A. D'Adda, P. Di Vecchia and M. Luscher, Nucl Phys B 146, 63 (1978); E. Witten, Nucl Phys B 149, 285 (1979).
- ³ I. Ya. Arefeva and S. I. Zakov, Nucl Phys B 162, 298 (1980).
- ⁴ For the $s = \frac{1}{2}$ AF Heisenberg models, standard derivations were given by I. Ichinose and T. Matsui, Phys Rev B 45, 9976 (1992); Phys Rev B 57, 13790 (1998); See also D. P. Arovas and A. Auerbach, Phys Rev B 38, 316 (1988).
- ⁵ F. D. M. Haldane, Phys Rev Lett. 61, 1029 (1988).
- ⁶ N. Read and S. Sachdev, Phys Rev Lett. 62, 1694 (1989); Phys Rev B 42, 4568 (1990).
- ⁷ See for example, M. Kamal and G. Murthy, Phys Rev Lett. 71, 1911 (1993) and references cited therein.
- ⁸ P. Tomczak and J. Richter, J Phys A 34, L461 (2001) and references cited therein; see also R. R. P. Singh, M. P. Gelfand, and D. A. Huse, Phys Rev Lett. 61, 2484 (1988).
- ⁹ A. M. Polyakov, Nucl Phys B 120, 429 (1977).
- ¹⁰ I. Ichinose and T. Matsui, Phys Rev Lett. 86, 942 (2001).
- ¹¹ I. Ichinose and T. Matsui, and M. Onda, Phys Rev B 64, 104516 (2001).
- ¹² H. Kleinert, F. S. Nogueira, and A. Sudb, Phys Rev Lett. 88, 232001 (2002); Nucl Phys B 666, 361 (2003).
- ¹³ I. F. Herbut and B. H. Seradjeh, Phys Rev Lett. 91, 171601 (2003).
- ¹⁴ I. F. Herbut, B. H. Seradjeh, S. Sachdev, and G. Murthy, Phys Rev B 68, 195110 (2003).
- ¹⁵ In the strong-coupling phase, the field z_x acquires nonvanishing mass by $h_x \neq 0$. In this case, the summation over z_x is convergent even for $U = 1$.
- ¹⁶ The model (11) can be extended for the strong-coupling phase in which $h_x \neq 0$. The nonlocal terms become as $\sum_{n=1}^{\infty} \frac{1}{n} e^{i \int d^3x U(x) W_x + (n) U^y (x + n \hat{e}) W_x^y (n)}$.
- ¹⁷ We calculate the correlation of the Polyakov lines instead of the Wilson loop since the system is anisotropic. The spatial correlation of the Polyakov lines can be an order parameter of the deconfinement transition in the gauge system (11).
- ¹⁸ A. M. Polyakov, Phys Lett B 72, 477 (1978); L. Susskind, Phys Rev D 20, 2610 (1979).
- ¹⁹ T. Senthil, L. Balents, S. Sachdev, V. V. Ishwanath, and M. P. A. Fisher, cond-mat/0312617, "Quantum criticality beyond the Landau-Ginzburg-Wilson paradigm".
- ²⁰ G. A. Rakawa, I. Ichinose, T. Matsui, and K. Sakakibara, work in progress.