

Projecting Fermion Pair Condensates into Molecular Condensates

Roberto B. Diener and Tin-Lun Ho

Department of Physics, The Ohio State University, Columbus, Ohio 43210

We offer strong evidence that the recent observations by M. Greiner, C. Regal, and D. Jin [1] and by MIT group [2] are signatures of a fermion superfluid in the strongly interacting regime made up of large fermion pairs. Our conclusions are based on calculations using crossover theory for different potentials including those with the characteristics of two-channel models. Our results demonstrate clearly universality near resonance. The T_c predicted by crossover theory is a perfect match with the observed boundary of vanishing condensate fraction with no adjustable parameters.

In an exciting recent paper, M. Greiner, C. Regal, and D. Jin reported evidence for condensation of fermion pairs of ^{40}K near Feshbach resonance [1]. Similar results in ^6Li were reported later by Ketterle's group at MIT but with a different interpretation [2]. Feshbach resonance is produced by Zeeman shifting the energy of a bound state in closed channel to zero energy. When this energy (linear in magnetic field B and usually referred to as "detuning") vanishes, a low energy fermion pair in the open channel strongly resonates with the bound state through hyperfine interaction, leading to considerable scattering. The regions $\Delta > 0$ and $\Delta < 0$ (which has negative and positive scattering length a_s respectively) are referred to the BCS and BEC side of the resonance because the ground state deep inside these regions are BCS superfluid and Bose-Einstein condensate (BEC) of molecules respectively. The intriguing region is near resonance ($\Delta \approx 0$) where the system becomes strongly interacting. The experiments in ref. [1] and [2] explore this region.

In ref. [1], an equilibrium state was prepared at an initial field B_i , which was quickly changed to B_f deeper in the BEC side. It is found that when B_i is sufficiently close to resonance, a molecular condensate will emerge at B_f such that the fraction of condensed molecules (N_0/N_m) is independent of the (fast) sweep rate [1]. The JILA group considers this as evidence for condensation of fermion pairs in the initial state. Their picture is the BCS-BEC crossover theory, which describes the BCS to BEC evolution as a smooth reduction in size of the condensed pair [3, 4, 5]. The immediate question is that near resonance the size of the pair should be of interparticle spacing, then the direct overlap between these pairs and the tightly bound molecules at B_f should be very small. In that case, the final molecular condensate must be generated by the fast sweeping field, and the relation between the momentum distributions in the final state and the initial state is not immediately clear.

In contrast, the MIT group thinks that their similar results are due to a BEC of tightly bound molecules in closed channel persisting onto the BCS side. They argue that even though these molecules are unstable in vacuum on that side, they can be stabilized by the Fermi sea. Somewhat earlier, Falco and Stoof (FS) [6] had

also suggested that the molecular fraction N_0/N_m observed at B_f in ref. [1] is given by the molecular component $Z(\Delta)$ in the condensed fermion pair in the initial state, $\text{pair}_i = \frac{Z(\Delta)}{Z(\Delta) + 1} \text{closed}_i + \frac{1}{Z(\Delta) + 1} \text{open}_i$, where closed_i and open_i are closed channel and open channel fermion pairs. Using a phenomenological resonance model, FS calculated the boundary of vanishing Z and found reasonable match with the boundary of vanishing N_0/N_m observed in ref. [1]. FS argue that even though when Z vanishes, the system still has pairing in open channel, and the observed boundary of vanishing N_0/N_m is a sharp crossover from molecular rich pair to fermion pair, (see fig. 1 in ref. [6]). This "persisting molecular BEC" pictures [2, 6] are very different from the usual crossover picture, which does not predict a sharp division between molecule rich and pure fermion pairs.

Adding to this confusion is the question of whether the many-body physics near a Feshbach resonance can only be described by "two channel" (or "resonance") models which incorporate open and closed channel physics specially [5], or can be described by simpler "single channel" models which consist of two types of fermions interacting with a tunable potential [3, 4]. Should these models have very different properties at resonance, and if the physics near resonance is universal as generally believed, it would imply that there are two distinct universality classes, and that only the two channel universality class gives the right physics of Feshbach resonance. The other possibility is that two channel models are not universal, and that the notion of universality inferred from current experiments [7, 8] is misconstrued.

In this paper, we point out that these confusions can be eliminated by deriving the precise expression for the observed molecular fraction, calculating the properties of the system explicitly using crossover theory, and examining the nature of the broken symmetry. We first perform $T = 0$ calculations using crossover theory [3, 4] for three different types of potential: V_I : a square well, V_{II} : a well with a high barrier, and V_{III} : a delta-function potential. Since V_{II} can accommodate a very long lived quasi-bound state, it has the major characteristics of a two channel (or "resonance") model. We shall show that all these potentials give rise to identical properties (such as chemical potential, energy gap, and coherence factors) near res-

onance, demonstrating universality in this regime. The fact that V_{II} carries all the characteristics of the two-channel model suggests both types of models may have identical properties near resonance. To our surprise, our calculations show that there is non-trivial overlap between the pairs at resonance (denoted as $x = 0$, where $x = (k_F a_s)^{-1}$ and k_F is the Fermi wavevector) and those at x as large as 12, which corresponds to the thermal field B_F in ref.[1]. This eliminates a serious concern formerly expressed by one of us (T.L.H.) [9] about the validity of projection argument in ref.[1]. To the extent that direct projection describes the experimental process, one can show that $N_0 = N_m$ is directly proportional to the superfluid order parameter, and the boundary for vanishing $N_0 = N_m$ in T - B plane observed in ref.[1, 2] is precisely the T_c between superfluid and normal gas, not a crossover between different types of pairs. Since V_I to V_{III} have identical properties near resonance, one can use the simplest potential (V_{III}) to study the finite temperature properties in this region. Such study had been performed in ref.[4] by studying Gaussian fluctuations about the mean field state. Remarkably, the phase boundary predicted in ref.[4] matches very well with the observed boundary for vanishing condensate fraction ($N_0 = N_m = 0$) [1, 2] with no adjustable parameters.

(A) The fraction of condensed molecules: Consider a fermion pair with total momentum q , $D_q^y(x) = \sum_{k; f_k; (x)} a_{k+q=2}^y a_{k+q=2}^y = 2$, where $a_{k; \uparrow}^y$ creates a fermion with momentum k and hyperfine spin \uparrow , $x = (k_F a_s)^{-1}$ describes the distance from resonance, and $f_k; (x)$ is the Fourier transform of the pair wavefunction $f(r; x) = \int_{-1/2}^{1/2} \frac{d^3r}{(2\pi)^3} e^{ik \cdot r} f_k; (x)$ which depends on x ; V is the volume of the system. A condensate of N zero momentum pairs is $|\Psi\rangle = N^{-1/2} \sum_{q=0}^N (x) |\Psi_q\rangle$. It reduces to a weak coupling BCS superfluid and a molecular condensate as $x \ll -1$ and $x \gg 1$ respectively [3, 4]. In grand canonical description, this pairing state becomes

$$|\Psi\rangle = N^{-1/2} \sum_{k; \uparrow} u_k(x) + v_k; (x) a_{k; \uparrow}^y a_{k; \downarrow}^y |\Psi\rangle \quad (1)$$

where N is the normalization constant, u and v are related to the Fourier transform of $f(r)$ as $f_k; (x) = v_k; (x) = u_k$, and V is a normalization constant.

Let x_0 and x be the distances of the initial and final state from resonance, and x be on the BEC side. When the system is jumped from x_0 to x , the number of condensed and un-condensed molecules emerge at x are $N_0 = \int d^3x D_0(x) i \int_{x_0}^x dx$ and $N_{ex} = \int_{x_0}^x d^3x D_q(x) D_q(x) i_{x_0}$ respectively. We then have from eq.(1),

$$N_0 = \int d^3x D_0(x) i \int_{x_0}^x dx f_k; (x) k; (x_0) = 2; \quad (2)$$

$N_{ex} = \int_{k,q} d^3k \int_{x_0}^x dx f_k; (x) \int_{n_q} d^3n_q n_q 2k; = 2$, where $k; (x_0) = \hbar a_{k; \uparrow}^y a_{k; \downarrow}^y i_{x_0}$, $n_k = \hbar a_{k; \uparrow}^y a_{k; \downarrow}^y i_{x_0}$, and we have ignored

in eq.(2) a term down by a factor of N^{-1} . Eq.(2) shows that N_0 is the overlap of the initial order parameter with the normal pair wavefunction.

Let us first discuss the single channel case because it is illuminating. Denoting the two spin states as \uparrow and \downarrow , the s -wave pairing function is $f(r) = f(r) i^y$, and

$$\frac{N_0}{N_m} = \frac{\int d^3x \int_{k,q} d^3k \int_{n_q} d^3n_q f_k(x) f_k(x) f_{k,q} f_{k,q}}{\int d^3x \int_{k,q} d^3k \int_{n_q} d^3n_q f_k(x) f_k(x) f_{k,q} f_{k,q}}; \quad (3)$$

where $N_m = N_0 + N_{ex}$, $(k) = \hbar a_{k; \uparrow}^y a_{k; \downarrow}^y i_{x_0}$, $n_k = \hbar a_{k; \uparrow}^y a_{k; \downarrow}^y i_{x_0} = \hbar a_{k; \uparrow}^y a_{k; \downarrow}^y i_{x_0}$. Since k vanishes at $T = T_c$, the curve $N_0 = N_m = 0$ in the T - B plane is therefore the boundary of superfluid to normal transition.

(B) The Crossover Picture: The basic assumption of crossover theory is that the ground state near resonance is still given by eq.(1). We have studied eq.(3) using the single channel crossover theory for: (I) a square well, $V_I(r) = \int_0^j$ (or 0) for $r < r_0$ (or $r > r_0$); (II) a well plus a barrier, $V_{II}(r) = \int_0^j$ for $r < r_0$. $V_{II}(r) = \int_1^j$ for $r_0 < r < r_1$, and $V_{II}(r) = 0$ for $r > r_1$; (III) a contact potential $V_{III}(r) = g(r)$, where $g = 2 \hbar^2 a_s M$, where M is the mass of the fermion. The Hamiltonian is $H = \sum_{k; \uparrow} \hbar^2 k^2 a_{k; \uparrow}^y a_{k; \downarrow}^y + \sum_{k,q} V(k) a_{q=2+k; \uparrow}^y a_{q=2+k; \downarrow}^y a_{q=2+k; \uparrow}^y a_{q=2+k; \downarrow}^y$, $k = \hbar^2 k^2 = 2M$. The range of the potential of V_I and V_{II} (r_0 and r_1) are taken to be of atomic size, much smaller than inter-particle spacing, i.e. $k_F r_0, k_F r_1 \ll 1$. By varying \int_0^j a bound state can be peeled off from the continuum, causing a_s to jump from -1 to $+1$. The main difference between V_I and V_{II} is that the latter can accommodate a quasi-bound state of atomic size, similar to the feature of the two channel or the resonance model.

For all cases, u_k and v are given by the well known expressions $\int_k \int = (E_k + \epsilon_k) = (2E_k)$, $\int_k \int = (E_k - \epsilon_k) = (2E_k)$, where $\epsilon_k = \frac{\hbar^2 k^2}{2M}$, $E_k = \frac{\hbar^2 k^2}{2M} + \frac{\hbar^2 k^2}{2M}$, and the energy gap ϵ_k at $T = 0$ is given by the gap equation $\epsilon_k = \int_{k_0}^{\infty} V(k) \epsilon_k^0 \epsilon_k^0 = (2E_{k_0})$. The corresponding expressions for ϵ_k and n_k are $\epsilon_k = u_k v_k = \epsilon_k = (2E_k)$, $n_k = v_k^2$. The chemical potential is determined by the number density n as $n = \sum_{k; \uparrow} n_k = n(T = 0;)$.

(C) Results: We have solved the gap equation numerically and have inverted the relation $n = n(\epsilon)$ so that quantities like $\epsilon_k; u_k; v_k; \epsilon_k; n_k$ are functions of the parameters of the potential (denoted collectively as fY_{ig}) and density n . Our results are displayed in figure 1 to 9. In these figures, we use the dimensionless parameter $x = (k_F a_s)^{-1}$ instead of potential parameters fY_{ig} . The results for potential V_I, V_{II} , and V_{III} will be colored in black, red, and green respectively. For both V_I and V_{II} , we take $k_F r_0 = 0.05$. For V_{II} , we choose $r_1 = r_0 = 1.2$ and $U_1 = 5 \int_0^{(m)} j$, where $\int_0^{(m)} j = \hbar^2 a_s^2 = (4M r_0^2)$ is the value for a square well to produce a bound state.

(C1) ϵ_k, ϵ_0 , and n_k : In figure 1a, we have plotted $\epsilon_k = 0$ versus $k r_0$ for $x = 2; 0; 2$, denoted as dashed,




FIG. 1: Fig.1a to 1c: Black, red, and green represent $V_I; V_{II}; V_{III}$, $x = (k_F a_s)^{-1}$, and $k_F r_0 = 0.05$. Dashed, solid, and dotted lines denote $x = 2; 0; -2$. Within the range $-2 < x < 2$, there are little changes in $\Delta(k)/\Delta(0)$, although $\Delta(0)$ and μ undergo significant changes (see Fig.1b), leading to very different $\Delta(k)$ in Fig.1c.

solid, and dotted lines respectively. The inset shows that $\Delta(k)/\Delta(0)$ for different potentials have similar values at low momenta (even up to a few k_F 's) but differ significantly at higher momenta, though it vanishes for realistic potentials V_I and V_{II} . Within the range $-2 < x < 2$, this ratio is essentially unchanged. The value of $\Delta(0)$ and μ , however, changes a lot as x varies from -2 to 2 , as shown in Fig.1b. These variations lead to the large differences in $\Delta(k)$ as one moves across the resonance. It is clear from Fig.1b and 1c that universal behavior sets in around resonance ($x = 0$) since different potentials give identical results. For all potentials, we find that n_k is accurately described by that of V_{III} . Its figure is given in ref.[4].

(C 2) Pair wavefunction f : To display f , we plot $f(k) = k_F r_0 f(r)$ versus $k_F r$ in Fig.2a to 2c. One sees that all potentials have essentially the same pair wavefunctions unless one goes deeper into the BEC side. The function f on the BEC side for all potentials have essentially the Feshbach form $f \propto e^{-r/a_s} = r/r_0$. Until a_s reduces to the atomic size r_0 , f has a non-trivial fraction beyond r_0 . It can therefore have significant overlap with the pair wavefunction near resonance. In fact, the overlap between f at $x = 0$ and that at $x = 12$ (which is the final state in ref.[1]) is as big as 18%. Thus, the direct projection picture suggested in ref.[1] (formerly criticized by one of us[9]) is in fact reasonable. In the case of V_{II} , we note that while a molecular component of range r_0 is visible on the BCS side near resonance, it essentially vanishes for $x < -2$. However, the $T = 0$ condensate fraction N_0/N_m is still very large for initial state $x_0 = 3$ which has no molecular component. This shows that the molecular component is unimportant in contributing to the projected condensate fraction N_0/N_m .

(C 3) The condensate fraction N_0/N_m , and the superfluid-normal phase boundary: In figure 3a, we show the $T = 0$ molecular fraction N_0/N_m at $x = 12$ as a function of the initial state x_0 . The data of ref.[1] and [2] are represented by circles and square respectively. The former is obtained at $T=T_F = 0.05$. The temperature of the latter is uncertain. Since the decrease of N_0/N_m on the BEC side appears to be related to three

body effects[1], we apply equilibrium theories only to the BCS side ($x < 0$). The large difference in scale between the results at $T = 0$ and at $T=T_F = 0.05$ is anticipated since T_c is strongly suppressed in the strongly interacting regime[3, 4], which will reflect in a strong suppression of N_0/N_m . The full effect of the crossover theory is best illustrated by showing T_c in the $T-x$ plane. In Fig.3b and 3c, we have superposed the T_c predicted in ref.[4] on the contour plots for N_0/N_m in the JILA [10] and the MIT [2] experiments. (Because of the universality near resonance demonstrated in Fig.1 and 2, we can use the finite temperature results for V_{III} [4] to compare with experiments.) The matches are remarkable, in view that there are no adjustable parameters.

In Fig.3a, we also show that as the final state moves deeper into the BEC side, the boundary of vanishing N_0/N_m moves to the opposite, i.e. BCS side. This is because while both N_0 and N_{ex} decrease as the final state becomes more tightly bound, the decrease of N_{ex} is faster than that of N_0 so that N_0/N_m actually increases. This fact can be demonstrated numerically, but can be established analytically if we take $f(k) = (k^2 + \mu^2)^{-1}$ and study the change of N_0/N_{ex} as a function of μ^2 . As $\mu^2 \rightarrow 1$, N_0/N_m reaches the asymptotic curve: $N_0/N_m = A/(A+B)$, where $A = \int d^3k f_k^2$, and $B = \int d^3k n_k n_{k-q}$, a prediction that can be verified experimentally. Since jumping to fields where molecular size are comparable r_0 (such as a deeply bound state) means sampling short distance behaviors of the order parameter (see Fig.1a), it will not be universal, as illustrated in the dashed and dotted curve in Fig.3a.

(D) Two channel models: For Feshbach resonances, f in Section (A) is $f(r) = \frac{1}{Z} f^{(c)}(r) f^{(o)}(r) + \frac{1}{Z} f^{(o)}(r) f^{(c)}(r)$, where f and $f^{(o)}$ are normalized orbital and spin functions, $f^{(c)}$ and $f^{(o)}$ denote close and open channel, and $f^{(c)}$ and $f^{(o)}$ are orthogonal. The zero momentum pair is $D_{q=0}^y = \frac{1}{Z} b_{q=0}^y + \frac{1}{Z} f_k^{(o)} f_k^{(c)} a_k^y a_k^y$, where $b_{q=0}^y = \frac{1}{Z} f_k^{(c)} f_k^{(c)} a_k^y a_k^y$ is the closed channel boson in the resonance model[5]. From eq.(2), we have




FIG. 2: Fig 2a to 2c: Pair wavefunctions for different values of x , $x = \frac{p}{4} = k_F r$. For V_{II} , f has a peak at $r < r_0$, yet most of the weight for all potentials is outside the atomic range r_0 . At resonance the size of f is given by k_F^{-1} .




FIG. 3: Fig 3a: The $T = 0$ value of N_0/N_m at $x = 12$ as a function of initial state x_0 is represented by the solid line. The data of ref.[1] and [2] are denoted as circles and squares. As the initial state x_m moves toward the BEC side, the curves move toward the BCS side, reaching a fixed asymptote, which depends on the microscopic nature of $\epsilon(k)$ at large momenta (see Fig. 1a). The asymptotes of V_I and V_{II} are represented by the dotted and dashed curve resp. In Fig 3b and 3c, we superpose the phase boundary calculated in ref.[4] (red triangle) on the data of ref.[10] and [2]. In Fig 3b, light color means large N_0/N_m . The numerical correspondence of the color code is given in ref.[10]. Dark blue corresponds to $N_0/N_m = 0$. In Fig 3c, we have used the approximate formula $x = (832 - B)/(B - 640)$ to convert magnetic field into $x = (\mu_B a_s)^{-1}$ [11].

$N_0 = \sum_k \frac{p}{Z} f_k^{(c)} f_k^{(c)} + \frac{p}{1-Z} f_k^{(o)} f_k^{(o)}$, where $f_k^{(c/o)} = \text{Tr} \rho^{(c/o)}_{k=2}$ are the components of the order parameter k ; in closed and open channel. If the initial state is a tightly bound molecule ($Z = 1$), then the vanishing of N_0 is given by the vanishing of $f_k^{(c)}$. However, to infer from a vanishing $f_k^{(c)}$ of a possible non-zero $f_k^{(o)}$, (like the vanishing Z in f means the open channel component $(1 - Z) \neq 1$), is incorrect. This is because k ; has no normalization constraint, and both $f_k^{(c)}$ and $f_k^{(o)}$ are dynamically connected through the coupling term $b_{q=0}^\dagger a_k a_k + \text{h.c.}$ in the Hamiltonian. Any non-vanishing pairing order in one channel will generate a non-vanishing order in the other.

We have also learned from D. Jin that for ^{40}K there is a shallow bound state (C) in the open channel. Thus, as the close channel bound state (B) is Zeeman shifted down, it will cross the zero energy state (A) and then the bound state (C) in the open channel. One can then go from A to B to C using a magnetic field sweep, so that a pair of fermions in the open channel can be swept into a bound state in the same channel. In that case, the system can be described by single channel models.

This work is supported by NASA GRANT NAG 8-1765 and NSF Grant DMR-0109255.

-
- [1] C. A. Regal, M. Greiner, and D. S. Jin, Phys. Rev. Lett. 92, 040403 (2004).
 - [2] M. Zwierlein, et al., Phys. Rev. Lett. 92, 120403 (2004)
 - [3] D. M. Eagles, Phys. Rev. 186, 456 (1969); A. J. Leggett, J. Phys. (Paris), Colloq. 41, 7 (1980); P. Nozières and S. Schmitt-Rink, J. Low Temp. Phys. 59, 195 (1985).
 - [4] C. S. de Melo, M. Randeria, and J. Engelbrecht, Phys. Rev. Lett. 71, 3202 (1993)
 - [5] M. Holland, et al. Phys. Rev. Lett. 87, 120406 (2001), E. Timmermans, Phys. Lett. A 285, 228 (2001), Y. Ohashi and A. Gri n, Phys. Rev. Lett. 89, 130402 (2002).
 - [6] G. Falco and H. Stoof, Phys. Rev. Lett. 92, 140402 (2004).
 - [7] K. M. O'Hara, et al., Science 298, 2179 (2002). M. E. Gehm, et al., Phys. Rev. A 68, 011401(R) (2003)
 - [8] T. Bourdel, Phys. Rev. Lett. 91, 020402 (2003)
 - [9] Search and Discovery, Physics Today, March 2004.
 - [10] See additional scientific information in <http://jila-www.colorado.edu/~jin/>.
 - [11] We use $a = a_{bg} (1 + W/(B - B_0))$, $B_0 = 823\text{G}$, $W = 180\text{G}$, $a_{bg} = 2000a_0$, $k_F^2 = 2000a_0^{-2}$ [2].