## Probing m icrocavity polariton super uidity through resonant Rayleigh scattering

I. Carusotto<sup>1,2</sup>, and C. Ciuti<sup>3</sup>

<sup>1</sup>Laboratoire Kastler Brossel, Ecole Normale Superieure,

24 rue Lhom ond, F-75231 Paris Cedex 05, France

<sup>2</sup>CRS BEC-INFM and Dipartimento di Fisica, Universita di Trento, I-38050 Povo, Italy

<sup>3</sup>Laboratoire Pierre Aigrain, Ecole Norm ale Superieure,

24 rue Lhom ond, F-75231 Paris Cedex 05, France

(Dated: April 14, 2024)

We investigate the two-dimensional motion of polaritons injected into a planar microcavity by a continuous wave optical pump in presence of a static perturbation, e.g. a point defect. By nding the stationary solutions of the nonlinear mean-eld equations (away from any parametric instability), we show how the spectrum of the polariton Bogoliubov-like excitations rejects onto the shape and intensity of the resonant R ayleigh scattering emission pattern in both momentum and real space. We ind a super uid regime in the sense of the Landau criterion, in which the R ayleigh scattering ring in momentum space collapses as well as its normalized intensity. More generally, we show how collective excitation spectra having no analog in equilibrium system s can be observed by tuning the excitation angle and frequency. P redictions with realistic sem iconductor microcavity parameters are given.

F

PACS num bers: 71.36.+ c, 42.65.-k, 03.75.K k

The concept of a quantum uid has played a central role in many elds of condensed matter and atom ic physics, ranging from superconductors to Helium uids [1] and, more recently, atom ic Bose-E instein condensates [2]. One of the most exciting manifestations of quantum behavior is super uidity, i.e. the possibility of the uid to ow without friction around an in purity [3].

In this Letter, we investigate the super uid properties of a two-dimensional gas of polaritons in a sem iconductor m icrocavity in the strong light-matter coupling regime [4]. In this system, the normal modes are superpositions of a cavity photon and a quantum well exciton. Thanks to their photonic component, polaritons can be coherently excited by an incident laser eld and detected through angularly or spatially resolved optical spectroscopy. Thanks to their excitonic component, polaritons have strong binary interactions, which have been dem onstrated to produce spectacular polariton am pli – cation e ects [5, 6] through matter-wave stimulated collisions [7], as well as spontaneous param etric instabilities [8, 9, 10].

Here, we study the propagation of a polariton uid in presence of static in perfections, which are known to produce resonant Rayleigh scattering (RRS) of the exciting laser eld [11, 12, 13, 14]. We show that super uidity of the polariton uid manifests itself as a quenching of the RRS intensity when the ow velocity in printed by the exciting laser is slower than the sound velocity in the polariton uid. Furtherm ore, a dram atic reshaping of the RRS pattern due to polariton-polariton interactions can be observed in both momentum and real space even at higher ow velocities. Interestingly, the polariton eld oscillation frequency is not xed by an equation of state relating the chem ical potential to the particle density, but it can be tuned by the frequency of the exciting laser. This opens the possibility of having a spectrum of collective excitation around the stationary state which has no analog in usual system s close to therm al equilibrium. We show in detail how these peculiar excitation spectra can be probed by resonant R ayleigh scattering.

A commonly used model for describing a planar microcavity containing a quantum well with an excitonic resonance strongly coupled to a cavity mode is based on the Ham iltonian [15]:

$$\begin{array}{rcl} & Z & X & h & i \\ H &= & dx & & \stackrel{Y}{_{i}}(x) & h_{ij}^{0} + V_{i}(x) & _{ij} & \stackrel{}{_{j}}(x) \\ & & & \stackrel{ij=fX,Cg}{Z} \\ & & + & \frac{-g}{2} & dx & \stackrel{Y}{_{X}}(x) & \stackrel{Y}{_{X}}(x) & \stackrel{}{_{X}}(x) & \stackrel{}{_{X}}(x) & + \\ & & Z \\ & & & + & dx & \sim F_{p} e^{i(k_{p}x & !_{p}t)} & \stackrel{Y}{_{C}}(x) + h.c.; \end{array}$$
(1)

where x is the in-plane spatial position and the eld operators  $x_{,c}(x)$  respectively describe excitons (X) and cavity photons (C). They satisfy Bose commutation rules,  $\begin{bmatrix} 1 & x \\ j & x \end{bmatrix} = 2 (x = x^{0})_{ij}$ . The linear Ham iltonian  $h^{0}$  is:

$$h^{0} = \sim \frac{!_{X}(ir)}{R};$$
 (2)

where  $!_{C}(k) = !_{C}^{0} \frac{p}{1+k^{2}=k_{z}^{2}}$  is the cavity dispersion as a function of the in-plane wavevector k and  $k_{z}$  is the quantized photon wavevector in the growth direction. R is the Rabi frequency of the exciton-cavity photon coupling. A at exciton dispersion  $!_{X}(k) = !_{X}$  will be assumed in the following. In this fram ework, polaritons simply arise as the eigenmodes of the linear H am iltonian (2);  $!_{LP(UP)}(k)$  denotes the dispersion of the lower (upper) polariton branch Fig.1 (a)].

The external force term proportional to  $F_p$  describes a coherent and monochrom atic laser eld of frequency  $!_{p}$  (called the pump), which drives the cavity and injects polaritons. Spatially, it is assumed to have a planewave pro le of wavevector  $k_p = \sin_p !_p = c_{r_p}$  being the pump incidence angle, so to generate a polariton uid with a non-zero ow velocity along the cavity plane. The nonlinear interaction term is due exciton-exciton collisional interactions and, as usual, is modelled by a repulsive (q > 0) contact potential. The anham onic excitonphoton coupling has a negligible e ect in the regim e considered in the present study [15].  $V_{X,C}$  (x) are externalpotential term s acting on the excitonic and photonic elds which can model the presence of disorder. Here, results for the specic case of a point defect will be presented. Note that point defects can be naturally present in state-of-the-art sam ples [16] or even be created deliberately by m eans of lithographic techniques.



FIG. 1: (a): linear dispersion of the lower (LP) and upper (UP) polariton branches. Typical position of the pump wavevector  $k_{\rm p}$  (arrow), well in the parabolic region of the LP dispersion; the dashed line is the parabolic approximation at small k. (b): internal vs. incident intensity curve showing bistable behavior. The dashed branch is unstable: the inversion points A and B are respectively due to a single-mode (Kerr) or a multimode (parametric) instability. Excitation parameters:  $k_{\rm p}$  = 0:314 m $^{-1}$ ,  $!_{\rm p}$ ,  $!_{\rm LP}$ ,  $(k_{\rm p})$  = 0:47 meV,  $!_{\rm X}$  =  $!_{\rm C}^{\rm C}$  = 1:4eV, 2  $_{\rm R}$  = 5 meV.

W ithin the mean-eld approximation, the timeevolution of the mean elds  $_{X,C}(x) = h^{2}_{X,C}(x)i$  under the Ham iltonian (1) is given by:

$$i\frac{d}{dt} = \begin{cases} x & (x) \\ c & (x) \end{cases} = \begin{cases} 0 \\ F_{p} e^{i(k_{p}x \cdot !_{p}t)} \\ + \end{cases} + h^{0} = \frac{i}{2} 1 + \\ V_{X} & (x) + gj_{X} & (x)^{2} \\ 0 \\ V_{C} & (x) \end{cases} = \begin{cases} 0 \\ c & (x) \end{cases}$$
(3)

In the quantum uid language, these are the Gross-Pitaevskii equations [2] for our cavity-polariton system. For simplicity, an equal rate is assumed for the dam ping of both the excitonic and the photonic elds. In the present work, we will be concerned with an excitation close to the bottom of the LP dispersion, i.e. the region m ost protected [10] from the exciton reservoir, which may be responsible for excitation-induced decoherence [17].

In the hom ogeneous case  $(V_{X,C} = 0)$ , we can look for spatially hom ogeneous stationary states of the system in which the eld has the same plane wave structure  $X_{j,C}(x;t) = \exp[i(k_p x !_p t)] \underset{X_{j,C}}{\overset{ss}{\underset{c}}}$  as the incident laser eld. The resulting equations

$$!_{X} (k_{p}) !_{p} \frac{i}{2} + g j_{X}^{ss} j_{X}^{2} s_{X}^{ss} + {}_{R} c_{C}^{ss} = 0 \quad (4)$$
$$!_{C} (k_{p}) !_{p} \frac{i}{2} c_{C}^{ss} + {}_{R} s_{X}^{ss} = F_{p}; \quad (5)$$

are the generalization of the state equation. W hile the oscillation frequency of the condensate wavefunction in an isolated gas is equal to the chem ical potential and therefore it is xed by the equation of state, in the present driven-dissipative system it is equal to the frequency !p of the driving laser and therefore it is an experim entally tunable parameter. As usual, stability of the solutions of Eqs. (4-5) has to be checked by linearizing Eq. (3) around the stationary state. For  $!_p > !_{LP}$  (k<sub>p</sub>), the relation between the incident intensity  $I_p / f_p f$  and the intemalone shows the typical S-shape of optical bistability (see Fig.1b and Refs.[18, 19, 20]). Note that nice hysteresis loops due to polariton bistability have been recently experimentally demonstrated [18] in the case  $k_p = 0$ . In the opposite case  $!_p < !_{LP} (k_p)$  (not shown), the behaviour of the system would instead be the typical one of an optical limiter [20].

In the stability region, the response of the system to a weak perturbation can be determ ined by means of the linearized theory. In the eld of quantum uids, this approach is called Bogoliubov theory [2]. By dening the slow ly varying elds with respect to the pump frequency as  $_{i}(x;t) = _{i}(x;t) \exp(i!_{p}t)$ , the motion equation of the four-component displacement vector  $~(x;t) = _{x}(x;t); _{c}(x;t); _{x}(x;t); _{c}(x;t)$ <sup>T</sup> reads

$$i\frac{d}{dt} \sim = L \sim + f_{d}; \qquad (6)$$

 $\mathbf{f}_d$  being the source term due to the weak perturbation and the spectral operator L being de ned as

$$L = \begin{bmatrix} 0 \\ 1_{x} + 2gj_{x}^{ss}j \\ 1_{y} + 2gj_{x}^{ss}j \\ 1_{y} + \frac{1}{2} \\ 1_{y}$$



 $g_{X}^{ss^{2}}e^{2ik_{p}x} = 0 \qquad 1$   $0 \qquad 0 \qquad C_{A} : (7)$   $g_{X}^{ss^{2}}e^{2ik_{p}x} = 0 \qquad 1$   $g_{X}^{ss^{2}}e^{2ik_{p}x} = 0 \qquad 1$ 

no signi cant m ixing between the LP and UP branches. Since we are interested in nearly-resonant excitation close to the bottom of the LP dispersion curve, we can describe the system in terms of the LP eld  $_{\rm LP}$  =  $X_{\rm LP \ X} + C_{\rm LP \ C}$  only, being  $X_{\rm LP}$  and  $C_{\rm LP}$  the Hopeld coe cients quantifying the excitonic and photonic components. In the parabolic approximation,  $!_{\rm LP}$  (k) '  $!_{\rm LP}^{0} + \sim k^{2} = 2m_{\rm LP}$  and the self-coupling constant is  $g_{\rm LP} = g \chi_{\rm LP} \frac{1}{2}$ . The mean-eld shift of the polariton mode is then  $!_{\rm M \ F} = g_{\rm LP} j \frac{s_{\rm S}}{1} \frac{2}{2}$ . Under these assumptions, the spectrum of the LP Bogoliubov excitations can be approximated by the simple expression

$$!_{LP}' !_{p} + k y \frac{i}{2} (2 !_{MF} + k p)(k p);$$
(8)

where k = k  $k_p$ ,  $_k = \sim k^2 = 2m_{LP}$ , the ow velocity  $v_p = \sim k_p = m_{LP}$ , and the electrive pump detuning

$$_{p} = !_{p} !_{LP} (k_{p}) !_{MF} :$$
 (9)

FIG.2: Left panels: exact B ogoliubov dispersion for the LP branches calculated from Eq.(7). R ight panels: corresponding RRS emission pattern in k-space. The intensity has been norm alized to the transmitted intensity. The  $k_{\rm p}$  point (indicated by a white circle) saturates by far the gray scale. Panels (a-f): resonant case  $_{\rm p}$  = 0 respectively in the linear regime (a-b), with gj  $_{\rm X}^{\rm ss}$   ${\rm f}$  = 0.2 m eV (c-d), 1 m eV (e-f). Panels (g-h):  $_{\rm p}$  > 0 case with gj  $_{\rm X}^{\rm ss}$   ${\rm f}$  = 0.504 m eV. Panels (i-j):  $_{\rm p}$  < 0 case with gj  $_{\rm X}^{\rm ss}$   ${\rm f}$  = 0.6 m eV. Pump wavevector:  $k_{\rm p}$  = 0.314 m  $^1$  (a-h), 0.408 m  $^1$  (i-j). Same cavity parameters as in Fig.1.

The eigenvalues of the operator L correspond to the frequencies of the Bogoliubov eigenm odes of the system. For each k, the spectrum is composed of four branches  $!_{UP;LP}$  (k): for each polariton branch (LP or UP), two branches exist, which are the in age of each other under the simultaneous transformations k !  $2k_p$  k and ! !  $2!_p$  ! [21]. Numerical calculations are shown in Fig.2. For the sake of clarity, only the branches relative to the LP have been traced, the ones relative to the UP being far away on the scale of the gure. These numerical results can be understood through the sim pli ed analytical approximation that follows.

Provided the interaction energy  $gj_X^{ss}f$  is much smaller than the polaritonic splitting  $!_{UP}$  !  $!_{LP}$ , there is

In the resonant case (  $_{p} = 0$ ), the branches touch at  $k = k_p$ . The e ect of the nite ow velocity  $v_p$  is to tilt the standard Bogoliubov dispersion [2] via the term k y. While in the non-interacting case in Fig.2(a) the dispersion remains parabolic, in the presence of interactions [Fig 2 (c) and (e)] its slope has a discontinuity at  $k = k_p$ : on each side of the corner, the + branch starts linearly with group velocities respectively given by  $v_g^{r;l} = c_s \quad v_p, c_s$  being the usual sound velocity of hysteresis curve of F ig.1 (b), the condition  $_{p} = 0$  corresponds to the inversion point A. If one moves to the right of the point A along the upper branch of the hysteresis curve, the mean-eld shift  $!_{MF}$  increases and the e ective pump detuning p becomes negative. In this case, as it is shown in Fig.1 (i), the branches no longer touch each other at kp and a full gap between them opens up for su ciently large values of  $j_{pj}$  (not show n).

On the other hand, the elective pump detuning p is strictly positive on the lower branch of the bistability curve of Fig.1 (b). In this case, the argument of the square root in (8) is negative for the wavevectors k such that  $_{p} > _{k} > _{p} 2 !_{MF}$ . In this region, the branches stick together [15] (i.e.  $\text{Re}[!_{+}] = \text{Re}[!_{-}]$ ) and have an exactly linear dispersion of slope  $v_{p}$  (Fig.2g). The imaginary parts are instead split, with one branch being narrowed and the other broadened [15, 21]. For

 $!_{M F} > =2$ , that is on the right of point B in Fig.1 (b), the multimode parametric instability [21] sets in. In the eld of quantum uids, this kind of dynamical instabilities are generally known as modulational instabilities [22].

The dispersion of the elementary excitations of the system is the starting point for a study of its response to an external perturbation. In particular, we shall consider here a weak and static disorder as described by the potential  $V_{C,X}$  (x). In this case the perturbation source term  $f_d$  is time-independent

$$\mathbf{f}_{d} = V_{X} \quad {}_{X}^{ss}; V_{C} \quad {}_{C}^{ss}; \quad V_{X} \quad {}_{X}^{ss}; \quad V_{C} \quad {}_{C}^{ss} \quad ^{T} \quad (10)$$

as well as the induced perturbation  $\tilde{d} = L^{1} f_{d}$ . The static disorder resonantly excites those Bogoliubov modes whose frequency is equal to !p. In the left panels of F ig 2, the excited m odes are given by the intersections of the mode dispersion with the horizional dotted lines. For the specic example of a spatially localized defect acting on the photonic component, we have plotted in the right panels of Fig2 and in Fig3 the photonic intensity j  $_{\rm C}$  f in respectively the momentum and the real space for di erent param eter regim es. These quantities correspond to the experim entally accessible far-and neareld intensity pro les [12, 13, 16] of the resonant Rayleigh scattering of the pum p (i.e. the coherently scattered light at the pump frequency  $! = !_p$ ). A sim ilar k-space pattem is obtained in the presence of a disordered ensemble of defects.



FIG. 3: Real space resonant Rayleigh scattering emission pattern for a localized defect at x = y = 0 acting on the cavity photon. The defect potential has a lateral size of 0.8 m and a depth of 1 m eV. In panel (a), linear regime as in Fig.2 (a) and (b). In panel (b), super uid regime as in Fig.2 (e) and (f). In each panel, the emitted intensity normalized to the transmitted intensity.

In the linear regime, the k-space emission pattern shown in Fig2(b) contains a peak at the incident wavevector  $k_p$ , plus the RRS ring [12, 16]. In the real

space pattern shown in Fig.3(a), as the polariton uid is m oving to the right, the defect induces a propagating perturbation with parabolic wavefronts oriented in the left direction.

In the presence of interactions, the RRS circle is transformed into a 1 -like shape with the low-k lobe more intense than the high-k one. If  $_{\rm p}$  < 0, the two lobes are separated by a gap (Fig 2), while they touch at  $k_p$ if p = 0 (Fig.2d). In this resonant case, when  $!_{MF}$  is large enough for the sound velocity  $c_{\!\scriptscriptstyle S}$  in the polaritonic uid to be larger than the ow velocity  $v_p$ , the slope of the + branch on the low -k side of the corner (F ig 2e) becomes positive and there is no intersection with the horizontal dotted line any longer. In this regime, RRS is no longer possible, and the polaritonic uid behaves as a super uid in the sense of the Landau criterion [3]. Once norm alized to the incident one, the RRS intensity is strongly quenched with respect to the previous cases and no RRS ring is any longer present. The weak em ission still visible in Fig 2 (f) is due to non-resonant processes, which are allowed by the nite broadening of the polariton modes. As no propagating m ode is resonantly excited, the perturbation in real space rem ains localized around the defect, as shown in Fig.3 (b). On the other hand, on the bottom of the bistability curve (where  $_{p} > 0$ ), the polariton gas is not super uid. The RRS intensity is even enhanced with respect to the linear regime because of the reduced linew idth of the Bogoliubov modes in the regions where branches stick together, as shown in Fig 2(g-h). the

To sum m arize, the polariton uid has a super uid behaviour with respect to static in purities if the equation  $!_{LP}(k) = !_{P}$  has no solutions with  $k \in k_{P}$ . If the corresponding linear regime equation  $!_{LP}(k) = !_{P}$  has a set of solutions corresponding to the elastic RRS ring, the e ect of super uidity is dram atic as the RRS ring is suppressed. W ithin the parabolic approximation in Eq. (8), a simple su cient condition for super uidity is found: p = 0 and  $(m_{LP} v_{P}^{2} \sim !_{MF}) < \sim j_{P}j$ .

In conclusion, we have shown the strict connection between the dispersion of the elementary excitations in a quantum uid ofm icrocavity polaritons and the intensity and shape of the resonant R ayleigh scattering on defects. In particular, we have pointed out some experimentally accessible consequences of polaritonic super uidity for realistic m icrocavity parameters. M ore in general, thanks to the coupling to externally propagating light, m icrocavity polaritonic systems appear to be promising candidates for the study of novel e ects in low-dimensional quantum uids.

W e acknowledge G.C.La Rocca with whom the original idea of the work was conceived. W e are grateful to Y.Castin and J.Dalibard for continuous discussions. LKB-ENS and LPA-ENS are two "Unites de Recherche de l'Ecole Normale Superieure et de l'Universite Pierre et Marie Curie, associees au CNRS". E lectronic address: Iacopo Carusotto@ lkb ens.fr

- [1] D. Pines and P. Nozieres, The theory of quantum liquids Vols.1 and 2 (Addison-Wesley, Redwood City, 1966).
- [2] L. Pitaevskii and S. Stringari, Bose-Einstein condensation (Oxford University Press, 2003).
- [3] A.J.Leggett, Rev.M od.Phys.71, S318-S323 (1999).
- [4] C.W eisbuch et al, Phys. Rev. Lett. 69, 3314 (1992).
- [5] P.G.Savvidis et al, Phys. Rev. Lett. 84, 1547 (2000).
- [6] M .Saba et al, Nature (London) 414, 731 (2001).
- [7] C.Ciutiet al, Phys. Rev. B 62, R 4825 (2000).
- [8] J.J.Baum berg et al, Phys. Rev. B 62, R16247 (2000).
- [9] G.Messin et al, Phys. Rev. Lett. 87, 127403 (2001).
- [10] For a recent review, see Sem icond. Sci. Technol. 18, Special Issue on M icrocavities, G uest Editors J. Baum berg and L. Vira, Publisher Sarah Quin (Bristol, UK, 2003).
- [11] H.Stolz et al, Phys.Rev.B 47, 9669 (1993).
- [12] R.Houdre et al, Phys.Rev.B 61, 13333R (2000).

- [13] R.Houdre et al, Phys. Rev. Lett. 85, 2793 (2000).
- [14] W . Langbein and J. M. Hvam, Phys. Rev. Lett. 88, 047401 (2002), and references therein.
- [15] For a review, see C. Ciuti, P. Schwendin ann, and A. Quattropani, Sem icond. Sci. Technol. 18, S279-S293 (2003) and references therein.
- [16] W .Langbein, proceedings of ICPS 26 (Edinburgh, UK, 2002).
- [17] S. Savasta, O. D i Stefano, and R. G irlanda, Phys. Rev. Lett. 90, 096403 (2003).
- [18] A.Baas et al, Phys.Rev.A 69, 023809 (2004).
- [19] N.A.G ippius et al., preprint cond-m at/0312214.
- [20] R.W. Boyd, Nonlinear Optics (A cadem ic Press, London, 1992).
- [21] C.Ciuti, P.Schwendim ann, A.Quattropani, Phys.Rev. B 63,041303(R) (2001); D.M.W hittaker, Phys.Rev.B 63,193305 (2001).
- [22] B.Wu and Q.Niu, Phys. Rev. A 64, 061603 (R) (2001).