On log{norm aldistribution on a comb structure

E.Baskin and A.Iomin Department of Physics and Solid State Institute, Technion, Haifa, 32000, Israel

We study specic properties of particles transport by exploring an exact solvable model, a socalled comb structure, where di usive transport of particles leads to subdi usion. A performance of Levy { like process enriches this transport phenomenon. It is shown that an inhom ogeneous convection ow, as a realization of the Levy{like process, leads to superdi usion of particles on the comb structure. A frontier case of superdi usion that corresponds to the exponentially fast transport is studied and the log{norm al distribution is obtained for this case.

PACS num bers: 05.40.-a, 05.40 Fb

Fractionalkinetics attractsm uch attention and various aspects of these phenom ena are rejected in the recent reviews [1, 2, 3, 4, 5]. It should be underlined that while subdi usion described in the fram ework of the fractional equations has num erous applications [2, 3, 6, 7, 8], there are only few examples on fractional superdi usion equations [4, 9, 10] related to Levy like process in dynam ical chaos [4, 9] and consideration of Levy process for the Langevin equations [10]. As it was mentioned in [5] \the superdiusion is far from being com pletely understood". Indeed, it is di cult to im agine the realphysical process with the in nite variance jump length during the nite time (Levy ight). In the Letter we present an example of real physical process which is absolutely analogous to the Levy ights. In this example we discuss a behavior of observable quantities, namely a position of the center of m ass of traveling contam inant and the form of the packet (the tail of a distribution function). The physical mechanism of this Levy { like e ect is an inhom ogeneous convection, and the space{tim e evolution of the contam inant in the presence of the inhom ogeneous convection ow is studied. To emphasize this relation between the inhom ogeneous convection and the Levy process, we consider the transport phenom enon on a subdi usive substrate, e.g. on a comb structure.

Our objective is to explain some complications and a possible mechanism of superdiusion on the comb structure depicted in Fig. 1. It is an analogue of a subdi usive m edia w here subdi usion has been already observed [11]. A com b m odel is known as a toy m odel for a porous medium used for exploration of low dimensional percolation clusters [12] and electrophoresis process [13]. It should be underlined that conditions or changes that are necessary to perform in the Liouville equation in order to observe superdiusion in the comb model are important for understanding the nature of this process from the general point of view. We show that a perform ance of Levy { like process along, say x { direction, is due to the inhom ogeneity of the corresponding x { com ponent of the convection current in the Liouville equation. The di usion process in such media (modeled by the comb structure) are anom alously slow with the subdi usive mean squared displacement of the order of hx^2 (t) i t; < 1.





There are external forces leading to convection. In general case, the velocity of the convection ow is space dependent, i.e. convection is inhom ogeneous. The question under investigation is how the observable shape of the initial packet changes, when the space{tim e evolution of the packet corresponds to the convection ow . It should be underlined that this problem arises in a variety of applications starting from transport of external species (pollution) in water ows through porous geological form ations [14, 15], problem s of di usion and reactions in porous catalysts [16] and fractal physiology [17, 18, 19]. We nd the conditions for the inhom ogeneous convection velocity, when the observable packet's shape corresponds to the norm ald i usion with = 1 for the mean squared displacement, or even superdiusion when hx² (t)i t; > 1. The inhom ogeneity will be taken in the form of a power law function ygx^s. In this case, the in nitely long ights take place during a nite xed time. We will show that the Levy process takes place when s > 0. Our primary objective is the consideration of a frontier case when s = 1. The corresponding solution of the Liouville equation for the distribution function of the transport particles is a so{called log{normaldistribution.

First, we describe the comb model. A special behavior of the di usion on the comb structure, depicted in Fig. 1, is that the displacement in the x (direction is possi-

ble only along the structure axis (x-axis, say at y = 0). Therefore the di usion coe cient as well as a mobility is highly inhom ogeneous in the y-direction. N am ely, the di usion coe cient is D_{xx} = D[°] (y), while the di usion coe cient in the y{direction along the teeth is constant D_{yy} = D. The Liouville equation

$$\frac{\partial G}{\partial t} + \operatorname{div} j = 0 \tag{1}$$

describes a random walk on this structure with the distribution function G = G(t;x;y), and the current $j = (D'(y)\frac{\partial G}{\partial x}; D\frac{\partial G}{\partial y})$. Therefore, it corresponds to the following Fokker{P lanck equation

$$\frac{\partial G}{\partial t} \quad D^{*} \quad (\mathbf{y}) \frac{\partial^{2} G}{\partial \mathbf{x}^{2}} \quad D \frac{\partial^{2} G}{\partial \mathbf{y}^{2}} = 0$$
(2)

with the initial conditions G(0;x;y) = (x)(y) and the boundary conditions on the in nities G(t; 1; 1) = 0and the same for the rst derivatives respect to x. Performing the Laplace and the Fourier transform s with respect to time and the x{coordinate, correspondingly, one obtains the solution for the subdi usion along the structure axis [11]

G (t;x;0) =
$$\frac{D^{1=2}}{2} \frac{z}{D^{2}t^{3}} d \exp \frac{x^{2}}{4D^{2}} \frac{D^{2}}{t}$$
: (3)

The total number of particles on the structure axis decreases with time

hGi=
$$\int_{1}^{2} G(t;x;0)dx = \frac{p}{2} \frac{1}{Dt}$$
: (4)

Therefore this solution of Eq. (3) describes the subdi usion when the number of particles is not conserved. In what follows consideration, this point will be bearing in m ind [20]. Therefore, this solution corresponds to the subdi usion with the second m om ent or the m ean squared displacem ent along the structure axis of the form

$$hx^{2}(t)i = \frac{hx^{2}(t)G(t;x;0)i}{hG(t;x;0)i} = D^{*} \frac{t}{D}^{1=2} : (5)$$

First, let us consider a transport problem on the comb structure due to the constant convection velocity $v_x = v_0$ (y). Here and in what follows one bears in m ind that the dimension of $v_s x^s$ is D'x. The di usive term in (2) is om itted, for simplicity. Then, we have

$$\frac{\partial G}{\partial t} + v_0 \quad (y) \frac{\partial G}{\partial x} \qquad D \frac{\partial^2 G}{\partial y^2} = 0:$$
(6)

Now applying the Laplace and Fourier transform, this equation is solved exactly with the solution in the form

G (t;x;0) =
$$\frac{D^{1=2}x}{t^{3}v_{0}^{2}} \exp(Dx^{2}=v_{0}^{2}t);$$
 (7)

and G = 0 for x < 0, since the distribution function m ust be positive. M oreover, the is no transport in the negative direction due to the initial conditions. It corresponds to the norm aldi usion with the second m om ent

$$hx^{2}(t)i = \frac{v_{0}^{2}}{D}t$$
: (8)

Therefore, to obtain superdi usion on the comb structure, it is not enough to add a fast process e.g. due to the constant convection. In what follows we discuss som e mechanism for the superdi usion, where we suggest an inhom ogeneous convection.

A superdi usive Levy { like process corresponds to innite ights during a nite xed time. Such a process could be a realization of an inhom ogeneous current in the x-direction. For example, the following two-dimensional convective current

 $j = (v_s x^s G (t; x; y) (y); D @G (t; x; y)=@y)$ (9)

after substituting in the Liouville equation (1) gives some modi cation of Eq. (6) in the form

$$\frac{\partial G}{\partial t} + \hat{V}_{x}G \quad (y) \qquad D \frac{\partial^{2}G}{\partial y^{2}} = 0; \qquad (10)$$

where s > 0 and

$$\hat{V}_{x}G = v_{s}x^{s}\frac{\partial G}{\partial x} + sv_{s}x^{s-1}G$$
 : (11)

A fier the Laplace transform with respect to the time, one presents the solution in the form

$$\begin{array}{c} h & i \\ G(p;x;y) = f(p;x) \exp (p=D)^{1-2} jy j; \quad (12) \end{array}$$

where we used that

$$\frac{e^2}{e^2} \exp \left(\frac{r}{D} \frac{p}{D} \frac{1}{y} \right) = \frac{p}{D} \left(2 \frac{p}{D} \frac{p}{D} \right) \exp \left(\frac{r}{D} \frac{p}{D} \frac{1}{y} \right) = \frac{e^2}{D} \exp \left(\frac{r}{D} \frac{p}{D} \frac{1}{y} \right)$$

Substituting Eq. (12) in Eq. (10), we obtain then the following equation for f = f(p;x)

$$v_s x^s f^0 + s v_s x^{s-1} f + 2 [pD]^{1-2} f = (x)$$
 (13)

with the following boundary conditions on the in nity f(p;1) = 0 and f(p;x) = 0 for x < 0. For s < 1, the solution of Eq. (13) has the following form

$$f = \frac{(x)}{v_s x^s} \exp - \frac{2(D p)^{1=2} x^{1-s}}{v_s (1-s)} ; \qquad (14)$$

where (x) is a step function: (x) = 1 for x = 0 and it is zero for x < 0. It corresponds to some kind of superdi usion along the structure axis where all moments are determined by the following distribution function

G (t;x;0) = (x)
$$\frac{D^{\frac{1}{2}}x^{1} x^{2}}{v_{s}^{2}(1 s) t^{3}} \exp \frac{D x^{2} x^{2}}{v_{s}^{2}(1 s)^{2} t^{3}}$$
 (15)

For example, hx^2 (t) i / $t^{\frac{1}{1-s}}$. When s = 0, Eq. (15) coincides with Eq. (7). To avoid de ciency at x = 0 where the solution is in nite, one needs to introduce a di usion process, by m eans the second derivative with respect to x. In this case, the equations (10), (11) and (13) describe the space time evolution of the initial pro le of particles on the asymptotically large scale x 1. Therefore, the solutions (14) and (15) are the asymptotic ones. W hen 1, it is more convenient to look for a solution on S a class of generalized functions. The case with s > 1needs a special care that is deserved a separate consideration [21]. When s = 1, it is a frontier case, with a strong Levy { like process. The so { called log { norm ald istribution could be realized here as well.

We consider the case of inhom ogeneous transport along the structure axis due to the inhom ogeneous drift with s = 1 and inhom ogeneous di usion of the form $D_{xx}(x;y) = dx^2$ (y). To this end we add the di usion dx^2 (y)@G=@x to the x{component of the current in

Eq. (9). The Fokker{Planck equation (2) reads now

$$\frac{\partial G}{\partial t} + \hat{V}_{x}G \quad (y) \qquad D \frac{\partial^{2}G}{\partial y^{2}} = 0; \qquad (16)$$

where di usion along the structure axis is

$$\hat{V}_{x}G = \hat{\alpha}x^{2}\frac{\theta^{2}G}{\theta x^{2}} \quad (2\hat{\alpha} \quad \underline{v})x\frac{\theta G}{\theta x} + vG : \quad (17)$$

A gain, perform ing the Laplace transform with respect to the time and presenting the solution of Eqs. (16) and (17) in the form of Eq. (12) we obtain the following equation for f = f(p;x)

$$d\tilde{x}^2 f^{(0)}$$
 (2 \tilde{a} y)x f^0 + (2 [pD]¹⁼² + v₁)f = (x) (18)

with the same boundary conditions as in Eq. (2). The following solution is obtained:

$$f(p;x) = \frac{(x)}{2[pD]^{1-2}} + \frac{C_1}{jxf};$$
(19)

where $r = \frac{1}{2} [(1 \ b) + \frac{p}{(b+1)^2 + 8} (dp)^{1-2}]$, and we denote $b = v_1 = d$ and $d = D = d^2$. Them odulus here indicates that the equation is invariant with respect to the inversion x ! x. We also used here that $x (0 \ x) = 0 x = (x)$ and $x^2 (0^2 \ x) = 0 x^2 = 2 \ (x)$ on a class of constant probe functions. This {function solution for the inhomogeneous equation can also be obtained by the Fourier transform. In this case the speci c solution in the Fourier space is $f_k = (2^2 \ pD)^{-1}$, whose the inverse Fourier transform gives the rst term in (19). The constant C_1 could be specified from the initial conditions. It does not carry any important information for the large scale asymptotics, and, without loosing of the generality, it could be equaled to the unity: $C_1 = 1$.

To perform the inverse Laplace transform it is convenient to use the long time asymptotics t ! 1 (p ! 0). In this case r 1 + a p w ith $a = 2d^{1-2} = (p + 1)$. It is also convenient to present the second term in (19) in the form of the exponential: $j_{xj} j^{r} = e^{r \ln j_{xj}}$. Hence the inverse Laplace transform becomes a standard procedure [22], and we obtain the following asymptotic solution:

G (t;x;0) =
$$\frac{x}{2}$$
 + $\frac{a \ln \dot{x} j}{2 \dot{x} j}$ exp $\frac{a^2 \ln^2 \dot{x} j}{4t}$: (20)

We would like to adm it here that jxj 1 and the logarithm ic function is positive. The rst term corresponds to the sm oothly tim e{decaying pining distribution, while the second term, which is the most important for the asym ptotic transport, corresponds to a som e kind of the log-norm ald istribution. It should be underlined that the asymptotic solution of Eq. (20) is independent of the coe cient d, which determines the diusion along the axis structure. It is natural, since an asymptotic space{ time evolution of the initial distribution is determined by the drift component of the current, but not by di usion. It means that one could consider the convective process analogous to Eqs. (6) and (10) instead of the more generalconsideration in the fram ework of Eqs. (16) and (17). In this case, of course, G (t;x;y) = 0 for x < 0.

Let us calculate the second m om ent analogously to Eq. (5). We have then hG i = $[1 + 4(a + v_1)] = D$ t, and the second m om ent is

$$hx^{2}Gi = (8^{p} t = a^{2}) exp[4t = a^{2}]:$$

Finely, the displacem ent is

$$p \frac{1}{hx^{2} (t)i} = \frac{s \frac{1}{hx^{2}Gi}}{hGi} / e^{2t=a^{2}}; \qquad (21)$$

This exponentially fast spreading is a result of the Levy{ like process introduced inside the system by means of the inhom ogeneous displacement current. There is another interesting interpretation of the obtained result. It could be considered as the Poisson distribution respect to the squared logarithm. Namely, if we denote $z = \ln^2 \frac{1}{2} \frac{1}{2} \frac{1}{2} \frac{1}{2}$ then we have from (20)

P (z)dz =
$$\frac{a}{2} e^{a^2 z = 4t} dz$$
: (22)

Hence, the average value of z is

It could be understood as the fact that time approaches to the in nity with the same rate as the squared logarithm of the x {coordinate, namely 4t $\vec{a} \ln^2 \dot{x}$ j. Therefore, returning to the log{norm alpart of the distribution function of Eq. (20) we have, approximately, that its asymptotic behavior is

G (t 1; jx j 1; y = 0)
$$\frac{1}{jx j \ln^2 jx j}$$
: (24)

It is simply to see, that the ux is zero on the in nities. Finely, one obtains that alleven m om ents of x diverge on the large scale asymptotics. The essential di erence between the distributions for subdi usion of Eq. (3) which gives for jxjt 1; that G (t; jxj0) / exp $(x^4=t)^{1=3}$, and the superdi usion of Eq. (24) is obvious. Therefore, the main mechanism of the transition from subdi usion in Eq. (5) to superdi usion in Eq. (21) on the comb structure is the inhom ogeneous convection.

In conclusion, we would like to classify a possible response of a substrate with anom abus transport. As it is shown here it depends on the external forcing according to the power rate of the inhom ogeneous convection $j_x(x;t) = v_s x^s$ (y)G (t;x;0). As it follows from the solution (15), when s < 0 it is subditusion [3, 23]. As we shown here, when s > 0 it is superditusion, Eq. (15). The hom ogeneous convection with s = 0 corresponds form ally to the norm all it usion as in Eq. (8), but the elective different usion coelected to v_0^2 =D is determined by the external

- [1] M F.Shlesinger, G M.Zaslavsky, and J.K lafter, Nature 363, 31 (1993).
- [2] Fractional Calculus in Physics, ed.R.Hilfer (W orld Scienti c, Singapore, 2000).
- [3] R.M etzler and J.K lafter, Phys. Rep. 339, 1 (2000).
- [4] G M. Zaslavsky, Chaos, 4, 25 (1994); Phys. Rep. 371, 461 (2002).
- [5] IM. Sokolov, J. K lafter, and A. Blumen, Phys. Today 55, 48 (2002).
- [6] K B. O kham and J. Spanier, The fractional Calculus (A cadem ic Press, O rlando, 1974).
- [7] E W . M ontroll and G H . W eiss, J. M ath. Phys. 10, 753 (1969);
- [8] E W .M ontrolland M F.Shlesinger, in Studies in StatisticalM echanics, v.11, eds J.Lebow itz and E W .M ontroll, (N oth (H olland, Am sterdam, 1984).
- [9] P.G rigolini, Quantum Irreversibility and Measurement, (W orld Scientic, Singapore, 1993); P.G rigolini, in Chaos{the Interplay between Stochastic and Deterministic Behavior, Lecture notes in Physics, 457, (Springer{ Verlag, Berlin, 1995).
- [10] H.C. Fogedby, Phys. Rev. Lett. 73, 2517 (1994).
- [11] V E. Arkhincheev and E M. Baskin, Sov. Phys. JETP 73, 161 (1991).
- [12] G H.W eiss and S.Havlin, Physica A 134, 474 (1986).
- [13] E. Baskin and G. Zilberstein, Electrophoresis 23, 2626

forcing v_0 . The frontier case with s = 1 has two features. The rst one corresponds to the log{norm aldistribution of transport particles, where one deals not with a sum of independent random variables but with their multiplication [8]. As the result the exponentially fast transport takes place (see Eq. (21)). The second feature is that the solution satis es to the natural boundary conditions, when the ux of transporting particles equals zero on the in nities for any time. When s > 1, it corresponds to superdi usion [21], as well.

The nal remark is that the relaxation along the structure axis is \inevitable" process due to the item $D^{\circ}(y)@G=@x$. Therefore, anom alous transport is due to the convection ow only on the asymptotically large scale x 1 where the di usion process could be om itted. This point is studied in detain in [21], where we shown that this approximation corresponds to Liouville{G reen asymptotic solution [24].

(2002).

- [14] B.Berkow itz, J.K laffer, R.M etzler, and H.Scher, W ater Resour. Res. 38, 1191 (2002)
- [15] R.Hilfer in Advanced in Chem. Phys. vol. XC II, edited by I.Prigogine and SA.Rice (John W iley, 1996) p. 299.
- [16] J.S.Andrade Jr., D A. Street, Y. Shibusa, S.Havlin, and H.E. Stanley, Phys. Rev. E 55, 772 (1997).
- [17] B.J.West, W. Deering, Phys. Rep. 246, 1 (1994).
- [18] T.F.Nonnenmacher and R.Metzler, in [2].
- [19] S.D orfm an, A. Iom in, and L.D orfm an, On tum or developm ent: fractional transport approach, unpublished.
- [20] In fact, we can consider the total distribution function G'(t;x) = dyG(t;x;y), which corresponds to the continuous time random walk model [3] with the conservation of the total number of particles. Nevertheless, we prefer to consider G(t;x;0), since it is more relevant value for practical applications [13].
- [21] A. Iom in and E. Baskin, in preparation.
- [22] See, for example, G A.Kom and T M.Kom, M athem atical H andbook, (M cG raw H ill, New York, 1968).
- [23] B.O 'Shaughnessy and I.P rocaccia, Phys. Rev. Lett. 54, 455 (1985).
- [24] F W J. O lver, Introduction to asymptotics and special functions (A cadem ic P ress, N ew York 1974).