

# Levy flights as subordination process: first passage times

Igor M. Sokolov

Institut für Physik, Humboldt Universität zu Berlin, Newtonstraße 15, 12489 Berlin, FRG

Ralf Metzler

NORDITA { Nordic Institute for Theoretical Physics,  
Blegdamsvej 17, 2100 Copenhagen, Denmark<sup>y</sup>

We obtain the first passage time density for a Levy flight random process from a subordination scheme. By this method, we infer the asymptotic behavior directly from the Brownian solution and the Sparre Andersen theorem, avoiding explicit reference to the fractional diffusion equation. Our results corroborate recent findings for Markovian Levy flights and generalize to broad waiting times.

An important issue in the theory of stochastic processes is the problem of first passage [1, 2, 3, 4]: Its solution is a key to the understanding of chemical reactions, stability of states of dynamical systems under external perturbations, extinction of populations, and many other problems in natural sciences. In normal Gaussian diffusion, a standard method of solution of the first passage problems is the method of images due to Kelvin [3, 4, 5]. However, for Levy flights (LFs) it has recently been demonstrated that the images method leads to a result that contradicts the Sparre Andersen theorem, according to which the first passage time density (FPTD) of a random walk process asymptotically follows the universal  $f(t) \sim t^{-3/2}$  behavior for any symmetric distribution of jump lengths [4, 6, 7]. This statement is of great importance since it not only points out possible inapplicability of Kelvin's method to superdiffusive jump processes, but also poses intricate questions of a correct continuous limit for Levy flights and their description within fractional diffusion or Fokker-Planck equations.

LFs are a paradigm for anomalous stochastic processes with a wide range of applications such as chaotic dynamics, processes in plasma, transport in micelles, or even quantum systems [8, 9, 10, 11, 12, 13, 14]. They can be defined in terms of a random walk process with long-tailed jump length distribution  $\langle x^2 \rangle = \langle x \rangle^2 + \langle x^2 \rangle^+$  ( $0 < \alpha < 2$ ) [15]. Alternatively, LFs can be considered as appearing from a Langevin equation with  $\alpha$ -correlated Levy noise [16, 17, 18]. The characteristic function of an LF given by

$$P(k;t) = \exp[-K^{(\alpha)} |k|^\alpha t] \quad (1)$$

is of stretched Gaussian type, where  $P(k;t) = \int_{-\infty}^{\infty} P(x;t) \exp(ikx) dx$  is the Fourier transform of the probability density function (PDF)  $P(x;t)$  [2, 8, 9, 19]. In what follows, the generalized diffusion constant  $K^{(\alpha)}$  will be set to unity. In the limit  $\alpha = 2$ , the characteristic function (1) reduces to the Gaussian

$P(k;t) = \exp[-K^{(2)} k^2 t]$ , the characteristic function of a standard random walk with Gaussian limit distribution and finite variance  $\langle x^2(t) \rangle = 2K^{(2)} t$  [1, 20], and finite higher order moments. For the general case  $0 < \alpha < 2$ , only fractional moments of the form

$$\langle |x|^{2+\alpha} \rangle = \frac{2^{1+\alpha}}{\Gamma(1+\alpha/2)} \frac{(1+\alpha/2-1)}{\Gamma(\alpha/2)} K^{(\alpha)} t^{\alpha/2}; \quad (2)$$

exist with  $\alpha < 2$  which can most easily be obtained by using properties of Fox H-functions [21].

The fractional diffusion equation [10, 11, 17, 21]

$$\frac{\partial P(x;t)}{\partial t} = K^{(\alpha)} \frac{\partial}{\partial |x|^\alpha} P(x;t) \quad (3)$$

governing an LF reflects the inherent long jumps by the non-locality of the fractional Riesz-Weyl operator [10, 18, 21, 22]

$$\frac{\partial P(x;t)}{\partial |x|^\alpha} = \frac{1}{2 \cos(\alpha/2)} \frac{\partial^2}{\partial x^2} \int_{-\infty}^{\infty} \frac{P(x';t) dx'}{|x-x'|^{2-\alpha}}; \quad (4)$$

for  $1 < \alpha < 2$ , and an analogous expression for  $0 < \alpha < 1$ ; its Fourier transform  $\mathcal{F}\{P(x;t)\} = \mathcal{F}\{P(k;t)\}$  [18, 22]. In the limit  $\alpha = 2$ , the fractional operator (4) reduces to the standard second-order partial derivative, as it should.

In terms of the dynamical equation for the PDF, the formulation of the first passage problem in terms of an absorbing (Dirichlet) boundary condition requires a cut-off in the Riesz-Weyl integral (4) [7]. Contrasting the universal Sparre Andersen result for the FPTD, it was found that the probability density of first arrival at a site differs from the FPTD, and is explicitly dependent on the index of the LF [7]. In the present work, we present an alternative derivation of the FPTD for LFs based on a subordination to a regular random walk process of the kind developed in Refs. [23, 24], leading us to a new, a priori unexpected twist: Although the PDF of the free process (without boundaries) is described by the fractional diffusion equation, Eq.(3), this equation does not uniquely describe the whole process, i.e. the exact properties of its sample paths. Depending on how anomalous

<sup>E</sup>lectronic address: igor.sokolov@physik.hu-berlin.de

<sup>y</sup>Electronic address: metzler@nordita.dk

transport statistics are introduced, we will discuss the appropriateness or failure of the images method.

Let us consider a stochastic process subordinated to a discrete random walk with some operational time that is defined by a one-sided Levy law [23]. The notion of subordination implies that the corresponding random process can be understood as follows [3]. The motion of the random walker can be parameterized by the number of steps  $n$ , the PDF of the random walker's position after  $n$  steps being given by the PDF  $P_{RW}(x;n)$ . The corresponding characteristic function  $\phi_n(k) = \int_{-\infty}^{\infty} P_{RW}(x;n) \exp(ikx) dx$  is determined by the PDF of step lengths  $\phi(x)$  such that  $\phi_n(k) = [\phi(k)]^n$  in terms of the characteristic function  $\phi(k)$  of the step length PDF, compare, for instance, Ref. [2].

The number of steps  $n$  itself that may be considered as the internal operational time of the process, is a non-decaying random function of the physical time (clock time)  $t$ . Denoting by  $p_n(t)$  the probability to perform exactly  $n$  steps up to time  $t$ , we obtain

$$P(x;t) = \sum_n P_{RW}(x;n) p_n(t); \quad (5)$$

At long times we suppose that a continuous operational time  $T$  may be introduced instead of the discrete index  $n$ . Moreover, we assume that the continuous analog of  $P_{RW}(x;n)$  exists. In this limit, we find from (5)

$$P(x;t) = \int_0^\infty P_x(x;T) p_T(T;t) dT; \quad (6)$$

where  $P_x(x;T)$  is the PDF to be at  $x$  at operational time  $T$  and  $p_T(T;t)$  is the PDF to be at operational time  $T$  at clock time  $t$ . This continuous limit corresponds exactly to the mathematical notion of the subordination, see Sec. X.7 of Ref. [3]: If  $fX(T)$  is a Markov process with continuous transition probabilities and  $fT(t)$  a process with non-negative independent increments, then  $fX(T(t))$  is said to subordinate to  $fX(t)$  using the operational time  $T$ .

Instead of having the PDF  $P_{RW}(x;n)$  of a Brownian random walk, let us assume the more general case that  $P_{RW}(x;n)$  corresponds to an LF, so that  $P_{RW}(x;n)$  is given by a symmetric Levy distribution of the form [3]

$$P_{RW}(x;n) = l_0 n^{1-\alpha} L\left(\frac{x}{l_0 n^{1-\alpha}}\right); 0 \quad (7)$$

( $0 < \alpha < 2$ ), where  $l_0$  is a scaling factor with length as physical unit. Note that the limit  $\alpha = 2$  corresponds to a Gaussian profile for  $P_{RW}(x;n)$ . Let us additionally assume that the number of steps per unit of the physical (clock) time  $t$  is distributed according to some distribution with a power-law tail:  $p(n; t=1) \sim n^{-1-\beta}$ , with  $0 < \beta < 1$ . Then, according to the generalized central limit theorem, at long times  $p_n(t)$  tends to a continuous limit distribution corresponding to the one-sided Levy law

$$p_T(T;t) = \frac{0}{t}^{1-\beta} L\left(T \frac{0}{t}\right); ; \quad (8)$$

Here,  $0$  is a scaling factor with physical unit time. The one-sided character of (8) ensures that  $p_T(T < 0; t) = 0$  [3, 19].

To obtain the limit distribution  $P(x;t)$  based on relations (7) and (8), we first Fourier-transform Eq. (6) in respect to  $x$ . With expression (1) for the characteristic function of a symmetric Levy stable density, we find

$$P(k;t) = \int_0^\infty \exp(-|k| l_0 T) \frac{0}{t}^{1-\beta} L\left(T \frac{0}{t}\right) dT; \quad (9)$$

This exactly corresponds to the Laplace-transform of a one-sided Levy density with Laplace variable  $u = |k| l_0$ . The Laplace transform of a one-sided Levy stable density is known [2, 19]:  $p_T(u;t) = \exp(-u t^{1-\beta})$ , and thus

$$P(k;t) = \exp(-|k| l_0 \frac{0}{t}^{1-\beta} t); \quad (10)$$

This expression, in turn, is the characteristic function of a symmetric Levy stable density with index  $\alpha$ . The scaling factor  $K(\alpha) = l_0 = 0$  can be interpreted as the associated fractional diffusion coefficient that depends on the indices of the corresponding subprocesses only through their product  $\alpha = \alpha_1 \alpha_2$ .

The PDF (10) fulfills the fractional diffusion equation (3) with order  $\alpha$ . This can also be shown on grounds of the subordination scheme developed in Refs. [23, 25]. However, it is remarkable that the dynamical equation (3) defines the PDF  $P(x;t)$ , but not the process (or better, the identity of the subprocesses) itself; unless we have to do with the limiting case  $\alpha = 2$  that necessarily corresponds to  $\alpha = 2$  and  $\beta = 1$ , defining the process uniquely. In other words, although the PDFs of all processes with identical  $\alpha$  are the same, these processes may still differ in the fractal dimension of their sample paths given by the set of jumps of random length corresponding to LFs with index  $\alpha$ ; and similarly they differ in the nature of the connection between operational time  $T$  and physical clock time  $t$  according to the one-sided Levy stable density with index  $\beta$ . This issue is of great importance for first passage problems, as we demonstrate now.

Our subordination procedure corresponds to a (random) change of the time variable of the process from operational time  $T$  to clock time  $t$ . This allows us to solve a number of problems connected to the underlying random process without explicitly referring to the fractional equation (3). We here consider the first passage across a boundary located at  $x = 0$ .

Let  $S_n(x_0)$  be the survival probability on the positive semi-axis (i.e. the probability of not crossing the boundary within the first  $n$  steps) after starting at  $x_0 > 0$  at  $n = 0$ . According to the Sparre Andersen theorem [3, 4, 6], the asymptotic form of this probability does not depend on the jump length distribution if only it is symmetric. For large number of steps, one invariably has  $S_n(x_0) \sim c(x_0) n^{-1/2}$ , where the prefactor  $c(x_0)$  depends

on the initial position  $x_0$ , as well as on  $\alpha$  and  $l_0$ . On subordinating the number of steps  $n$  to physical clock time  $t$  we see that the survival probability up to time  $t$  is

$$S(t; x_0) = \sum_n p_n(x_0) p_n(t); \quad (11)$$

where  $p_n(t)$  is the probability that exactly  $n$  steps occur within clock time  $t$ . Note that equation (11) corresponds to the spatial average of Eq. (5). Changing from  $n$  to the continuous operational time variable  $T$ , we get:

$$S(t; x_0) = \int_0^{\infty} p_T(T; x_0) p_T(T; t) dT; \quad (12)$$

This and  $p_n(x_0) \sim c(x_0)n^{1-2}$  gives rise to

$$S(t; x_0) \sim c(x_0) \frac{1}{1-2} \frac{1}{2} \frac{0}{t}^{1-2}; \quad (13)$$

as derived in the Appendix. We now obtain the FPTD,

$$f(t) = \frac{dS(t; x_0)}{dt} \sim \frac{c(x_0)}{1-2} \left( 1 + \frac{1}{2} \frac{0}{t^{1+1-2}} \right); \quad (14)$$

The following limiting cases can be distinguished:

(i) If the subordination from operational time  $T$  to clock time  $t$  through  $p_T(T; t)$  is narrow with  $\alpha = 1$ , i.e.,  $p_T(T; t) = \delta(T - t)$ , the universal  $f(t) \sim t^{3-2}$  behavior according to the Sparre Andersen theorem is recovered. In other words, in order to change this asymptotic behavior, one has to consider  $0 < \alpha < 1$  explicitly.

(ii) If we consider the process subordinated to Gaussian diffusion ( $\alpha = 2$ ), then  $\alpha = 2$  and

$$f(t) \sim \frac{2C(x_0)}{1-2} (1 + \alpha) \frac{0}{t^{1+1-2}}; \quad (15)$$

This result has the same scaling as the FPTD derived through the method of images in Ref. [7]. A few words on the interpretation of this seemingly paradoxical finding are in order. Result (15) corresponds to a random walk process with a Gaussian jump length density  $\psi(x)$ , so that the corresponding trajectory is that of a regular Brownian walk. It is therefore perfectly legitimate to use the images method for such a process, even though it is described by the fractional diffusion equation (3). In contrast, the genuine ('classical') LF discussed in Ref. [7] corresponds to a broad  $\psi(x) \sim |x|^{-1}$  with explicitly smaller than 2, but  $\alpha = 1$ . For this strongly non-local process with fractal trajectory, the FPTD follows the result (i). In this latter case, that is, the method of images fails and the FPTD follows  $f(t) \sim t^{3-2}$ .

(iii) In general, for a given  $\alpha$  one has to make sure that the inequalities  $\alpha = 2 < \alpha < 1$  are fulfilled, since simultaneously the two conditions  $\alpha = 1$  and  $\alpha = 2$  with  $0 < \alpha < 2$  have to be met. The FPTD for such a general  $\alpha$  therefore shows the following asymptotic behavior

$$f(t) \sim t^{-\alpha}; \quad \alpha = 2 \quad 1 + 1 = \alpha; \quad (16)$$

where  $0 < \alpha < 2$ . In this scheme, the Sparre Andersen decay with exponent  $3-2$  is the slowest one possible. This makes perfect sense since due to the Levy stable form of  $p_n(t) \sim n^{-1}$  a broad distribution of single jump events occurs in a finite time interval  $(0; t)$ , increasing the likelihood of crossing the boundary within any given finite time interval dramatically. The Levy nature of  $p_n(t)$  thus leads to an oversampling of the space in comparison to a process  $\alpha = 1$ .

The subordination scheme developed here allows one to express the FPTD of a random process solely on the basis of the properties of the Sparre Andersen universality and the subordination map from operational time  $T$  to physical clock time. Starting off from an LF in  $(n; t)$  coordinates, we introduce a broad distribution  $p_n(t) \sim n^{-1}$  of events per clock time interval  $t$ . We find that the resulting process is governed by the fractional diffusion equation (3) whose order  $\alpha$  is a product of the Levy index of the jump length distribution,  $\alpha$ , and the subordination distribution  $p_n(t)$ . The knowledge of  $\alpha$  alone is therefore insufficient to deduce the exact form of the jump length PDF and the trajectory it gives rise to.

As a direct consequence, the resulting FPTD in the limit  $p_n = \delta_{n,1}$  (the trivial subordination with  $\alpha = 1$ ) fulfills the Sparre Andersen universality for any process with symmetric jump length distribution. This case includes the case of genuine LFs as those discussed in Ref. [7], and is violated by the method of images for all  $0 < \alpha < 2$ . Conversely, for  $\alpha = 2$ , the subordination process has the trajectory of a normal Brownian random walk and is amenable to the images method to determine the FPTD. The result scales like  $f(t) \sim t^{1-1}$  as previously obtained. In the case of general  $\alpha$  and  $\alpha$ , the range spanned by the exponent in the FPTD  $f(t) \sim t^{1-\alpha}$  is  $3-2 \leq 1 + 1 = \alpha$ .

We believe that above findings help in interpreting the inadequacy of the images method for genuine LFs as found in [7] and moreover show that caution is necessary when generalizing well-known results from ordinary to anomalous diffusive processes.

We acknowledge helpful discussions with Aleksei Chechkin and Yossi Klatter.

#### Appendix A: CALCULATION OF THE $(1-2)$ -ORDER MOMENT

To obtain the final form of Eq. (13), we have to evaluate the integral

$$\int_0^{\infty} T^{1-2} \frac{0}{t}^{1-2} L\left(T \frac{0}{t}\right) dT; \quad ; \quad ;$$

which is equal to the  $(1-2)$ -order moment

$$M_{1-2}(\alpha) = \int_0^{\infty} T^{1-2} L\left(T \frac{0}{t}\right) dT$$

of the one-sided Levy stable density  $L(\cdot; \cdot; \cdot)$ . To calculate this expression, one can follow two alternative

routes. As such techniques are useful when dealing with Levy processes, we sketch both ways.

(i) The density  $L(\cdot; \cdot; \cdot)$  is the inverse Fourier transform of the characteristic function  $\hat{\rho}(k) = \exp(-jk)e^{-i/2}$  for  $k > 0$ , continued to the negative semi-axis by  $\hat{\rho}(k) = \hat{\rho}^*(k)$ . Then,

$$M_{1=2}(\cdot) = \int_0^{\infty} \frac{d}{1=2} \operatorname{Re} \int_0^{\infty} \frac{dk}{2} e^{-ik} \exp(-jk)e^{-i/2}$$

Interchanging both integrals, one can express the integral over  $k$  through Fresnel integrals to obtain  $\int_0^{\infty} e^{-ik} dk = \frac{1}{2}(1-i)k^{-1/2}$ . Changing then  $ik$  to  $u$  we find for the overall integral:

$$M_{1=2}(\cdot) = \int_0^{\infty} \frac{du}{2} \exp(-u) u^{1/2} :$$

Noting that the integrand is analytical everywhere except for  $u = 0$ , vanishes for  $|j| \geq 1$  in the lower right quadrant of the complex plane, and that the integral converges, we can change the contour of integration to the positive real axis, so that

$$M_{1=2}(\cdot) = \int_0^{\infty} du \exp(-u) u^{1/2} : \quad (A1)$$

Substituting  $u = u$ , we recover the integral representation of the  $H$ -function, so that Eq. (13) follows.

(ii) Alternatively, the density  $L(\cdot; \cdot; \cdot)$  is the inverse Laplace transform of  $\exp(-u) = {}_1H_{0;1}^{1;0}(u; 1=)$  expressed in terms of the

Fox  $H$ -function [21, 26]. By standard methods [26, 27], the inverse Laplace transform can be performed, yielding

$$L(\cdot; \cdot; \cdot) = (\cdot)^{-1} {}_1H_{1;1}^{1;0} \left( \frac{1}{(\cdot); 1=} \right) :$$

After a substitution, the resulting integral

$$M_{1=2}(\cdot) = \frac{1}{2} \int_0^{\infty} z^{1/2} {}_1H_{1;1}^{1;0}(z; 1=) dz$$

can be easily evaluated, by noting that it corresponds to the Mellin transform  $\hat{g}(s) = \int_0^{\infty} t^{s-1} g(t) dt$  of  $H_{1;1}^{1;0}(z)$  at  $s = 1/2$ . The result can be directly identified with the definition of the  $H$ -function [21, 26], so that  $M_{1=2}(\cdot) = {}_1(1=2) = (1=2)$ , reproducing Eq. (13).

- 
- [1] N. van Kampen, Stochastic Processes in Physics and Chemistry (North-Holland, Amsterdam, 1981).
  - [2] B. D. Hughes, Random Walks and Random Environments, Volume 1: Random Walks (Oxford University Press, Oxford, 1995).
  - [3] W. Feller, An Introduction to Probability Theory and Its Applications (John Wiley & Sons, New York, NY, 1968).
  - [4] S. Redner, A guide to first-passage processes (Cambridge University Press, Cambridge, UK, 2001).
  - [5] H. S. Carslaw and J. C. Jaeger, Conduction of heat in solids (Clarendon Press, Oxford, UK, 1948).
  - [6] E. Sparre Andersen, Math. Scand. 1, 263 (1953); ibid. 2, 195 (1954).
  - [7] A. V. Chechkin, R. Metzler, J. Klafter, V. Gonchar, and L. V. Tanatarov, J. Phys. A 36, L537 (2003).
  - [8] J. P. Bouchaud and A. Georges, Phys. Rep. 195, 127 (1990).
  - [9] M. F. Shlesinger, G. F. Zaslavsky, and J. Klafter, Nature 363, 31 (1993).
  - [10] R. Metzler and J. Klafter, Phys. Rep. 339, 1 (2000).
  - [11] I. M. Sokolov, J. Klafter, and A. Blumen, Phys. Today 55(11), 48 (2002).
  - [12] G. M. Zaslavsky, Phys. Rep. 371, 461 (2002).
  - [13] A. V. Chechkin, V. Y. Gonchar, and M. Szydlowski, Phys. Plasma 9, 78 (2002).
  - [14] H. Katori, S. Schlipf, and H. Walther, Phys. Rev. Lett. 79, 2221 (1997).
  - [15] J. Klafter, A. Blumen, and M. F. Shlesinger, Phys. Rev. A 35, 3081 (1987).
  - [16] B. J. West, P. Grigolini, R. Metzler, and T. F. Nonnenmacher, Phys. Rev. E 55 99 (1997)
  - [17] H. C. Fogedby, Phys. Rev. Lett. 73, 2517 (1994); S. Jespersen, R. Metzler, and H. C. Fogedby, Phys. Rev. E 59, 2736 (1999).
  - [18] A. V. Chechkin, V. Yu. Gonchar, J. Klafter, R. Metzler, and L. V. Tanatarov, J. Stat. Phys., at press.
  - [19] P. Levy, Theorie de l'addition des variables aleatoires (Gauthier-Villars, Paris, 1954). B. V. Gnedenko and A. N. Kolmogorov, Limit Distributions for Sums of Random Variables (Addison-Wesley, Reading, MA, 1954).
  - [20] P. Levy, Processus stochastiques et mouvement Brownien (Gauthier-Villars, Paris, 1965).
  - [21] R. Metzler and T. F. Nonnenmacher, Chem. Phys. 284, 67 (2002).
  - [22] S. G. Samko, A. A. Kilbas, and O. I. Marichev, Fractional Integrals and Derivatives, Theory and Applications (Gordon and Breach, New York, NY, 1993).
  - [23] I. M. Sokolov, Phys. Rev. E 63, 011104 (2001).
  - [24] D. Brockmann and I. M. Sokolov, Chem. Phys. 284, 409 (2002).
  - [25] I. M. Sokolov, J. Klafter and A. Blumen, Phys. Rev. E 64, 021107 (2001).
  - [26] H. M. Srivastava, K. C. Gupta, and S. P. Goyal, The  $H$ -functions of one and two variables with applications (South Asian Publishers, New Delhi, 1982).
  - [27] W. G. Glockle and T. F. Nonnenmacher, J. Stat. Phys. 71, 755 (1993).