

In uence of retardation e ects on 2D magnetoplasmon spectrum

M . V . Cheremisin

A .F .Io e Physical-Technical Institute, St.Petersburg, Russia

(Dated: December 24, 2021)

Within dissipationless limit the magnetic field dependence of magnetoplasmon spectrum for unbounded 2DEG system found to intersect the cyclotron resonance line, and, then approaches the frequency given by light dispersion relation. Recent experiments done for macroscopic disc-shape 2DEG systems confirm theory expectations.

PACS numbers: 73.20.Mf, 71.36.+c

Plasma oscillations in two-dimensional electron gas(2DEG) were first predicted in the middle 60th [1], and, then observed experimentally in liquid helium system [2] and silicon inversion layers [3, 4]. The recent pioneering observation [6] of retardation effects in the magnetoplasmon (MP) spectrum, discussed more than three decades ago, recom m ences the interest to the above problem . With retardation effects accounted we analyze 2D MP spectrum first derived in Ref.[5], and, then demonstrate the relevance of theory predictions with respect to experimental data [6].

Let us assume unbounded 2D electron gas imbedded in a dielectric in the presence of the perpendicular magnetic field. Following Ref.[7], the Maxwell equations for in-plane components of the electrodynamic potentials A ; χ yield

$$\begin{aligned} \square \chi &= 4\pi j_{\chi} A = \frac{4\pi j}{c} \chi; \\ \text{div} A + \frac{1}{c} \frac{\partial \chi}{\partial t} &= 0; \\ j_{\chi} &= -\frac{1}{c} \frac{\partial A}{\partial t} \chi; \end{aligned} \quad (1)$$

where $\square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$ is the d'Alembert operator, the conductivity tensor. Assuming the magnetoplasmon $e^{(i q r - i t)}$ propagated in 2DEG, and, then separating longitudinal and transverse in-plane components of the vector potential [7], 2D magnetoplasmon dispersion relation yields:

$$\frac{1}{2} + \frac{i_{xx}}{1} = \frac{1}{2} \frac{i_{xx}}{c^2} + \frac{y_{xx}}{c^2} = 0; \quad (2)$$

where $\chi = \frac{q^2}{c^2} \frac{w^2}{c^2}$. This result is exactly that obtained by Chui [5]. Within dissipationless limit the components of the conductivity tensor $i_{xx} = i_{yy} = \frac{i n e^2}{m (1 + \frac{c^2}{\omega^2})}$, $y_{xx} = y_{yy} = \frac{i c}{1} i_{xx}$ allow us to simplify Eq.(2) as follows

$$(Q^2 - 2)^{\frac{1}{2}} = \frac{1 + \frac{2}{c^2}}{4} + \frac{1 + \frac{2}{c^2}}{2}; \quad (3)$$

where we introduce the dimensionless wave vector $Q = \frac{q c}{\omega}$, frequency $\omega = \frac{1}{1_p}$ and cyclotron frequency $\omega_c =$




FIG .1: Zero- field plasma dispersion (bold line) asymptotes: light dispersion (dashed line), short-wavelength plasma limit (dotted line). Thin line represents the $\omega_c(Q)$ when the condition $\omega = \omega_c$ is satisfied. The calculated zero- field plasma frequency (open dots) and MP-CR intersection frequency (closed dots) are presented for two different samples ($Q = 3.94$ and $Q = 1.52$, see the text) used in Ref[6].

$\frac{1}{1_p}$. Then, $\omega_p = \frac{2 n e^2}{m c^2}$ is the plasma frequency. In absence of the magnetic field Eq.(3) reproduces the conventional [1, 7] zero- field longitudinal plasma dispersion relation $\omega^2 = 2 i_{xx} \omega$ as follows $Q^2 = 2 + \frac{4}{Q^2}$. In the short-wavelength limit $Q \gg 1$ one obtains square-root plasma dispersion (dotted line in Fig.1) $\omega = \frac{2 n e^2}{m} q$. The opposite long-wavelength limit case $Q \ll 1$ corresponds to light dispersion relation $\omega = \frac{c q}{1}$ shown in Fig.1 by the dashed line. Note, the authors [6] demonstrate the excellent agreement between the simplest zero- field 2D plasma theory [1, 9] and experimental results. For different disc-geometry quantum well samples the wave vector reported to relate to 2DEG disc radius via $q = \frac{1}{R}$; $R = 1.2$ in consistent with theory $R = 3.8$ [8].

The question we attempt to answer is whether retardation effects should modify 2D magnetoplasmon spectrum. In actual fact, Eq.(3) demonstrate that at fixed Q the plasma frequency grows with magnetic field (see Fig.2), and, then intersects the CR line. Note, this behavior is unexpected within the edge magnetoplasmon




FIG. 2: 2D magnetoplasmon spectra vs dimensionless cyclotron frequency Ω_c for different MP wave vector $Q = 2; 5; 10; 30$. Arrows represent the light frequency $\omega = \frac{c}{R}$. Cyclotron resonance is represented by dotted line.

formalism [10, 11]. Substituting $\omega = \Omega_c$ into Eq.(3) we derive the dependence (Fig.1, thin line) of the MP-CR intersection frequency vs wave vector $\Omega_{cr}(Q)$. It to be noted that low-eld magnetoplasmon remains longitudinal at $\Omega_c < \omega$, and, then becomes transverse one when $\Omega_c > \omega$.

It is instructive to compare the lowest angular momentum MP spectrum reported in [6] with that provided by the present theory. For high-density GaAs/AlGaAs heterostructure parameters ($n = 2.54 \cdot 10^{11} \text{ cm}^{-2}$, $m = 0.067m_0$; $R = 0.5 \text{ mm}$, $\omega = 12.8$) one obtains $\omega_p = 51 \text{ GHz}$ and $Q = 3.94$ (see Fig.1). The calculated zero-eld plas-

mon frequency $f_0 = \frac{\omega_p}{2} = 16 \text{ GHz}$ and $f_{cr} = \frac{\omega_{cr}}{2} = 24 \text{ GHz}$ are well comparable with experimental values 20 GHz and 32 GHz respectively. For higher 2D density sample ($n = 6.6 \cdot 10^{11} \text{ cm}^{-2}$, $R = 0.5 \text{ mm}$) we obtain $\omega_p = 132 \text{ GHz}$, $Q = 1.52$, and then $f_0 = 22 \text{ GHz}$ and $f_{cr} = 26 \text{ GHz}$ compared to the respective experimental values 27 GHz (36 GHz). In contrast, low-density sample ($n = 2.54 \cdot 10^{11} \text{ cm}^{-2}$; $R = 0.1 \text{ mm}$, $Q = 109$; $f_0 = 14.6 \text{ GHz}$) should exhibit intersection of the MP spectrum with CR line at $B = 2.52 \text{ T}$, i.e. within magnetic field range, which is higher than that used in Ref.[6].

Further increase of the magnetic field results in saturation of the magnetoplasmon spectrum (Fig.2) at certain frequency given by the light dispersion relation $\omega = \omega_p$. Experimentally, irrespective to 2DEG density the larger disc-mesa samples ($R = 0.5 \text{ mm}$) demonstrate [6] MP spectra (with the lowest radial and angular momentum) cut-off at certain frequency $f = 50 \text{ GHz}$. The latter coincides with frequency deduced from light dispersion relation $f = \frac{1}{2} \frac{c}{R} = 51 \text{ GHz}$. Note, in contrast to predicted MP spectrum saturation in strong fields, the experiments [6] demonstrate intriguing zigzag behavior which remains unexplained within our formalism.

In conclusion, we demonstrate the strong influence of retardation effects on magnetoplasmon spectrum in unbounded 2DEG system. The magnetic field dependence of MP spectrum found to intersect the cyclotron resonance line, and, then approaches the frequency given by light dispersion relation. The recent MP experiments in large disc-shape 2DEG system confirm theory predictions.

This work was supported by RFBR (grant 03-02-17588) and LSF (HPR I-CT-2001-00114, Weizmann Institute)

[1] F. Stern, Phys. Rev. Lett. 18, 546 (1967).
 [2] C.C. Grimes and G.A. Adams, Phys. Rev. Lett. 36, 145 (1976).
 [3] S.J. Allen, D.C. Tsui, R.A. Logan, Phys. Rev. Lett. 38, 98 (1977).
 [4] T.N. Theis, J.P. Kotthaus, P.J. Stiles, Solid State Comm. 26, 603 (1978).
 [5] K.W. Chui and J.J. Quinn, Phys. Rev. B 9, 4724 (1974).
 [6] I.V. Kukushkin et al, Phys. Rev. Lett. 90, 156801 (2003).
 [7] V.I. Falko and D.E. Khmel'nitskii, Sov. Phys. JETP 68,

1150 (1989).
 [8] R.P. Leavitt and J.W. Little, Phys. Rev. B 34, 2450 (1986).
 [9] In both 3D and 2D cases the edge plasmon frequency known to differ with respect to bulk case (see K.W. Chui and J.J. Quinn, Phys. Rev. B 5, 4707 (1972) and [10, 11])
 [10] A.L. Fetter, Phys. Rev. B 32, 7676 (1985).
 [11] V.A. Volkov and S.A. Mikhailov, Sov. Phys. JETP 67, 1639 (1988).