

Ab initio results for the electronic structure of $C_{50}Cl_{10}$.

Steven W.D. Bailey¹ and Colin J. Lambert¹

¹ *Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK*

In this paper we use ab initio density functional theory (DFT) to calculate the electronic properties of $C_{50}Cl_{10}$. In comparison with the unstable C_{50} which has a small $t_{lu}(LUMO) - h_u(HOMO)$ energy gap and a high total free energy compared with C_{60} , the belt of chlorine atoms stabilize the $C_{50}Cl_{10}$ fullerene by increasing the energy gap to approximately that of C_{60} and lowering the total free energy. We also examine the effects of inter-cage separation on the band structure for infinite periodic $C_{50}Cl_{10}$ chains where a high degree of dispersion is found to persist for separations beyond the predicted $C_{60} - C_{60}$ distance of closest approach.

A new example of the fullerene family, $C_{50}Cl_{10}$ has recently been produced as a stable solid [1]. The fullerenes smaller than the C_{60} buckyball are very difficult to produce in a stable form as the structure breaks the isolated pentagon rule (IPR) [2]. In C_{50} there are five sets of adjacent pentagons and this together with the high curvature of the molecular surface means it is highly unstable [3–5]. Xie et al have overcome this instability by attaching chlorine atoms around the C_{50} equator as seen in Fig. 1(a). In what follows, we examine the electronic properties and stability of $C_{50}Cl_{10}$, along with the electronic properties of chains of $C_{50}Cl_{10}$ molecules. To model $C_{50}Cl_{10}$ we use a double zeta basis set with a cutoff radius of 150 Ry to describe the carbon and Chlorine atoms within the SIESTA [6] implementation of DFT, where nonlocal norm-conserving pseudopotentials describe the core electrons and a linear combination of atomic orbitals the valence electrons. The relaxed, isolated molecule is shown in Fig. 1(a). In the case of a chain we consider molecules aligned along the z axis as shown in Fig. 1(b).




FIG. 1. Relaxed atomic coordinates of $C_{50}Cl_{10}$. The chlorine atoms form a fan around the equator of the C_{50} molecule (a). The proposed chain of $C_{50}Cl_{10}$ in the z direction (b)

TABLE 1. compares the Fermi energy E_F (eV), the total free energy E_{tot} (eV) and the $t_{lu}(LUMO) - h_u(HOMO)$ energy gap E_Δ (eV) between the C_{60} , C_{50} molecules and $C_{50}Cl_{10}$ as a linear chain with varying inter-cage separations in (Å), up to the isolated molecule.

TABLE 1.

Case	a	b	c	d	e	f
	C_{60} mol	C_{50} mol	$C_{50}Cl_{10} 2.5\text{\AA}$	$C_{50}Cl_{10} 2.9\text{\AA}$	$C_{50}Cl_{10} 3.4\text{\AA}$	$C_{50}Cl_{10}$ mol
E_F (eV)	-4.61	-7.04	-5.41	-5.41	-5.47	-5.67
E_{tot} (eV)	-9201	-7670	-11751	-11751	-11751	-11750
E_Δ (eV)	1.80	0.16	—	—	—	1.80

The table shows that the unstable C_{50} has a high E_{tot} of -7670eV in comparison to that of -9201eV for the C_{60} molecule but that the belt of ten Cl atoms stabilizes the $C_{50}Cl_{10}$ to produce a lower E_{tot} of -11751eV . This is reflected in the result for the $t_{lu}(LUMO) - h_u(HOMO)$ energy gap, (E_Δ), where the energy gap of 0.16eV for C_{50} opens to 1.80eV , matching that of C_{60} , for the $C_{50}Cl_{10}$ molecule. It is noted that the value of $E_\Delta = 1.80\text{eV}$ for C_{60} is in very good agreement with the experimental values of E_Δ between $(1.6 - 1.8)\text{eV}$ [7]

Fig. 2 shows the energy levels of the isolated molecules in cases (a,b and f) and the electronic band structures of $C_{50}Cl_{10}$ chains with the different inter-molecule separations in cases (c - e), of Table 1. In each case the energy is measured from the E_F for each relaxed structure studied. In cases(c - e) the band structure is shown for a wavevector kz in the z direction as defined in Fig. 1. The $C_{60} - C_{60}$ distance of closest approach is predicted to be approximately

3.0Å [8] therefore the range of cage separations for cases (c - e) were selected to cover this interval and beyond. The band structures for the infinite periodic chains of $C_{50}Cl_{10}$ for cases (c - e) show the dispersion of the $t_{lu}(LUMO)$ and $h_u(HOMO)$ bands persists up to and beyond a 3.4Å separation and that the $C_{50}Cl_{10}$ chain is an insulator. In conclusion we have used DFT calculations to investigate the electronic properties of the newly produced, solid $C_{50}Cl_{10}$ fullerene. We find that the belt of chlorine atoms around the equator of the C_{50} stabilize the structure by dramatically lowering the total free energy of $C_{50}Cl_{10}$ and increasing the $h_u(HOMO)$ - $t_{lu}(LUMO)$ gap to match that of C_{60} . The $t_{lu}(LUMO)$ and $h_u(HOMO)$ bands show a marked dispersion even for separations of the $C_{50}Cl_{10}$ cages in excess of the predicted closest $C_{60} - C_{60}$ approach of 3.0Å. This points to some interesting properties arising from carbon nanotube peapods such as (12, 12)@ $C_{50}Cl_{10}$.




FIG. 2. For the structures (a - f) defined in Table 1, the energy levels of the isolated molecules are seen in cases (a,b and f) and the electronic band structures for wavevectors kz in the z direction as defined in Fig. 1, in cases (c - e).

-
- [1] Su - Yuan Xie et al, Science. **304**, 5671 (2004).
 - [2] K. M. Kadish and R. S. Ruoff, Fullerene: Chemistry, Physics and Technology (Wiley, new York, 2002).
 - [3] T. Guo et al., Science **257**, 1661 (1992).
 - [4] H. W. Kroto, Nature **329**, 529 (1987).
 - [5] J. R. Heath, Nature **393**, 730 (1998).
 - [6] S. D. S-Portal, P. Ordejon, E. Artacho, J. Soler Int. J. Quantum Chem. **65**, 453 (1997).
 - [7] M.S. Dresselhaus, G. Dresselhaus and P. C. Eklund, Science of Fullerenes and Carbon Nanotubes (Academic, San Diego, 1996).
 - [8] A. Rochefort, Phys. rev. B, **67**, 115401, (2003).