

## Spectra of pinned charge density waves with background current

V. Gurarie and J. Levinsen

Physics Department, University of Colorado, Boulder, Colorado 80309-0390

(Dated: March 10, 2022)

We develop techniques which allow us to calculate the spectra of pinned charge density waves with background current.

## I. INTRODUCTION

The problem of a quantum mechanical particle whose motion is governed by a random time-independent Hamiltonian has been discussed in the literature for several decades. Because of its relevance to the electrons' motion in disordered conductors, the Hamiltonian which consists of a kinetic energy and a random scalar potential is probably the most famous example of such a problem. It is now well established that some of the wave functions of such Hamiltonians are localized, the phenomenon usually referred to as Anderson localization [1]. A wealth of new types of behavior was discovered once random Hamiltonians constrained by certain symmetries were studied. For example, the most general random Hamiltonian would be complex hermitian. Constraining it to be real (in other words, in posing time reversal invariance) changes the localization properties of its wave functions. An electron moving in a disordered conductor is described by a real random Hamiltonian. Turning on a magnetic field makes the Hamiltonian hermitian. This made the study of the crossover between real and hermitian random Hamiltonians easily accessible experimentally, and a subject of intense theoretical investigations [2].

Furthermore, it was discovered that a number of other symmetries can also be imposed on random Hamiltonians, which change their behavior yet again. Altogether, there are ten symmetry classes of random Hamiltonians, distinguished by nine different constraints imposed on them [3, 4].

Yet other types of random Hamiltonians arise when they are constrained not by a symmetry but by certain requirements their spectrum must satisfy. The most prominent example of that would be system with bosonic excitations [5]. To be specific, consider an energy functional  $E[\psi(x)]$ , where  $\psi(x)$  is some function, which has the form

$$E[\psi(x)] = \int_0^L dx \left[ \frac{1}{2} \left( \frac{d\psi(x)}{dx} \right)^2 + h(\psi(x); x) \right] \quad (1)$$

Here  $h(\psi(x); x)$  is a random function whose form will be specified below. Suppose  $\psi_0(x)$  is a minimum (local or global) of this functional. We would like to study the normal modes  $\psi_n$  of oscillations around that minimum, described by the equation

$$\frac{d^2}{dx^2} \psi_n(x) + \partial^2 h(\psi_0(x); x) \psi_n(x) = \epsilon_n \psi_n(x) \quad (2)$$

Eq. (2) is a random Schrodinger-like equation. Yet it is not equivalent to a particle moving in an arbitrary random potential.  $\partial^2 h(\psi_0(x); x)$  is a random function of  $x$ , but it is not an arbitrary random function. It is clear, for example, that  $\epsilon_n \geq 0$  for all  $n$ , which tells us that  $\partial^2 h(\psi_0(x); x)$ , although random, has to be constrained in such a way that all its eigenvalues are positive.

In most applications of Eq. (1),  $h(\psi; x)$  is chosen to be a smooth function of  $\psi$  and a rough function of  $x$ . For example,  $h(\psi; x) = A(\psi) \cos(\phi(x))$ , where  $A(\psi)$  and  $\phi(x)$  are random functions of  $x$  uncorrelated at different  $x$ , and  $\phi(x)$  is uniformly distributed between 0 and  $2\pi$ . This leads to the two point correlation function

$$\langle h(\psi; x) h(\psi_0; x_0) \rangle = \langle A(\psi) A(\psi_0) \rangle \cos(\phi(x) - \phi(x_0))$$

Under this definition of  $h(\psi; x)$ , the problem described by Eqs. (1) and (2) has been extensively studied in the context of pinned charge density waves [6]. Knowing the spectrum of Eq. (2) allows, for example, to calculate the AC conductance of the charge density wave. It was first deduced in Refs. [7] and [8] that the density of states  $\rho(\epsilon)$  of Eq. (2) is given by  $\rho(\epsilon) = \epsilon^{-1/2}$  if  $\epsilon < \epsilon_c$  and if  $\psi_0(x)$  is a global minimum of the energy functional Eq. (1), where  $\epsilon_c = 2\pi^2$  is the crossover scale. If, on the other hand,  $\psi_0(x)$  is but a local minimum of Eq. (1), then  $\rho(\epsilon) \sim \epsilon^{-1/2}$  for  $\epsilon < \epsilon_c$ .

A more detailed approach to the problem specified by Eq. (2) was developed in Ref. [9]. It was shown in that work that the potential  $\partial^2 h(\psi_0(x); x)$  of the "Schrodinger" equation Eq. (2) can always be represented as

$$\partial^2 h(\psi_0(x); x) = \frac{dV(x)}{dx} + V^2(x);$$

where  $V(x)$  is some new random function. As a result, Eq. (2) is equivalent to

$$H \psi_n(x) = \epsilon_n \psi_n(x); \quad (3)$$

where

$$H = -\frac{d}{dx} \left( \frac{d}{dx} + V(x) \right) \frac{d}{dx} + V(x); \quad (4)$$

and  $\epsilon_n = \epsilon_n^2$ . Now  $H$  is an example of a Hamiltonian constrained by a symmetry and can be solved by techniques developed in that context. The symmetry of  $H$  is usually referred to as chiral symmetry. It is expressed

by the relation  $\mathcal{H}_3 = \mathcal{H}$ , which holds true for any random  $V(x)$  [3]. Here  $\mathcal{H}_3$  is the usual Pauli matrix.

The full solution of Eq. (2), with the help of the mapping to Eq. (3), demonstrated that there indeed exists the crossover energy scale  $\epsilon_c$ . At  $\epsilon < \epsilon_c$  all the eigenfunctions  $\psi_n(x)$  are localized with the localization length  $\ell_n \sim \epsilon_c^{-1/2}$ , which is independent of  $n$ . At  $\epsilon > \epsilon_c$ , the eigenfunctions are still localized, but with the localization length which increases with  $n$  as  $\ell_n \sim n^{1/2}$ . The density of states at  $\epsilon < \epsilon_c$  is given by  $\rho(\epsilon) \sim \epsilon^{-1/2}$ . Finally, the density of states at  $\epsilon > \epsilon_c$  is given by  $\rho(\epsilon) \sim \epsilon^{-3/2}$  if  $\psi_0(x)$  is a global minimum of Eq. (1) and by  $\rho(\epsilon) \sim \epsilon^{-1/2}$  if  $\psi_0(x)$  is a local minimum of Eq. (1).

In this paper we would like to show that the same techniques which proved useful in solving Eq. (2) can also be used to solve another related problem, which we formulate below. Consider an equation

$$j \frac{d^2}{dx^2} - \frac{d}{dx} + \mathcal{E} h(\epsilon; x) = 0; \quad (5)$$

where  $j$  and  $\mathcal{E}$  are some parameters. If  $\mathcal{E} = 0$ , then this equation is equivalent to the minimization condition of the energy Eq. (1). If, on the other hand,  $\mathcal{E} > 0$ , then this defines a new problem. Eq. (5) was suggested in Ref. [10] to describe pinned charge density waves with background current. Normal modes of oscillations of such pinned charge density wave are given by

$$j \frac{d^2}{dx^2} - \frac{d}{dx} + \mathcal{E}^2 h(\psi_0(x); x) \psi_n(x) = \epsilon_n \psi_n(x); \quad (6)$$

where  $\psi_0(x)$  is a solution to Eq. (5). In this paper we are going to present the solution to the problem defined by Eq. (5) and Eq. (6).

The operator in the square brackets of Eq. (6) is non-hermitian. As a result, the eigenvalues  $\epsilon_n$  do not have to be real. It is well known, however, that the eigenvalues of equations of the type Eq. (6) always lie along one dimensional curves in the complex plane [11]. In this paper, we show that these curves take the shape depicted on Fig. 1. The fork point  $\epsilon_f$  cannot be found exactly. However, it is still possible to define the crossover scale  $\epsilon_c$ . All the states with energy less than  $\epsilon_c$  have the same localization length. As the energy is increased past  $\epsilon_c$ , the localization starts to grow and eventually diverges at  $\epsilon_f$  (which is always bigger than  $\epsilon_c$ ). The states corresponding to complex values of energy are delocalized.

We introduce the technique which allows to compute  $\epsilon_c$  and  $\epsilon_f$ . We calculate them in the regime where  $\mathcal{E} = (j)^{1/3} \gg 1$  and find them to be  $\epsilon_c = \mathcal{E}^2 = 4j$  and  $\epsilon_f = 2\mathcal{E}^2 = 8j$ . We also calculate the density of states along the part of the curve depicted on Fig. 1 which lies on the real axis and find it to be  $\rho(\epsilon) \sim \epsilon^{-1/2}$ .

The rest of this paper is organized as follows. In section II we map the problem defined by Eq. (6) to a random chiral Hamiltonian. In section III the localization length of normal mode oscillations will be derived in the limit of large background current and in section IV this result will




FIG. 1: The eigenvalues  $\epsilon_n$  of the Eq. (6) lie on this curve in the complex plane.

be compared with the usual Larkin length of the problem. Section V contains the calculation of the density of the low lying states. Finally, in Section VI we discuss the response of a pinned charge density wave with background current to an electric field using the formalism developed here.

## II. MAPPING TO A CHIRAL HAMILTONIAN

The first step of the solution is to map Eq. (6) to a more tractable form of a random chiral Hamiltonian. Following the general guidelines of Ref. [9], we treat Eq. (6) as the equation of motion of a particle with coordinate  $x$ , moving in time  $x$ , in the presence of viscous forces given by the  $\frac{d}{dx}$  term. We then pass from the Lagrangian to Eulerian description of the motion. Introducing the velocity function  $u(\psi_0(x)) = \mathcal{E} \psi_0(x)$  the equation of motion Eq. (6) becomes

$$j \partial_x u + j u \partial_x u + u = \mathcal{E} h(\psi_0(x); x); \quad (7)$$

The equation thus obtained is similar to the equation of motion of a one dimensional fluid without pressure, usually referred to as Burgers equation. However, it also features the  $u$  term, which can be interpreted as a kind of viscous friction. The Burgers equation with this term is novel and has not been considered in the literature before.

Performing a derivative on Eq. (7) with respect to results in

$$j \partial_x \mathcal{E} u + j (\mathcal{E} u)^2 + j u \mathcal{E}^2 u + \mathcal{E} u = \mathcal{E}^2 h; \quad (8)$$

We define the gradient of the velocity along the solution to Eq. (5),  $\partial_x$ , as  $F(x) = \mathcal{E} u(\psi_0(x); x)$ .  $F(x)$  can be related to  $h(\epsilon; x)$  with the help of Eq. (8)

$$j \frac{dF}{dx} + j F^2 + F = \mathcal{E}^2 h(\psi_0(x); x); \quad (9)$$

Thus Eq. (6) becomes

$$\frac{d^2}{dx^2} - \frac{d}{j dx} + \frac{dF}{dx} + F^2 + \frac{F}{j} \psi_n = \frac{n}{j} \psi_n; \quad (10)$$

or in a more symmetric form

$$\frac{d}{dx} + F + \frac{1}{j} \frac{d}{dx} + F - \frac{1}{j} \psi_n = \frac{1}{j} \psi_n : \quad (11)$$

The operator on the left-hand-side of Eq. (11) is not Hermitian. This may be remedied by using a trick due to Hatano and Nelson [12]. Writing

$$\psi_n(x) = e^{-\frac{1}{2j}x} \tilde{\psi}_n(x) \quad (12)$$

the equation becomes

$$\frac{d}{dx} + F + \frac{1}{2j} \frac{d}{dx} + F + \frac{1}{2j} \tilde{\psi}_n = \frac{1}{j} \tilde{\psi}_n : \quad (13)$$

Thus we mapped our equation into an eigenvalue problem for a Hermitian operator. This operator can also be rewritten in the form similar to Eq. (3),

$$H \frac{\tilde{\psi}_n(x)}{\tilde{\psi}_n(x)} = \frac{1}{j} \frac{\tilde{\psi}_n(x)}{\tilde{\psi}_n(x)} ;$$

where  $H$  is still given by Eq. (4),  $\frac{1}{j} = \frac{1}{2j}$ , and

$$V(x) = F(x) + \frac{1}{2j} :$$

However, as argued by Hatano and Nelson, the transformation Eq. (12) is valid only as long as  $\tilde{\psi}_n$  decays asymptotically as  $\exp(-|x|j)$ , with the localization length  $\tilde{\sim} < 2j = \frac{1}{2}$ . For  $\tilde{\psi}_n$  whose localization length obeys this condition, Eq. (11) is equivalent to Eq. (13), and therefore,  $\psi_n$  is real and positive. For other eigenfunctions  $\tilde{\psi}_n$  whose localization length is larger than  $2j = \frac{1}{2}$ , Eq. (13) is no longer equivalent to Eq. (11).

Notice that the localization length  $\tilde{\sim}$  of  $\tilde{\psi}_n$  is related to the localization length  $\psi_n$  of  $\psi_n$  as in

$$\psi_n = \frac{1}{\tilde{\sim}} \frac{1}{2j} ; \quad (14)$$

as long as  $\tilde{\sim} < 2j = \frac{1}{2}$ .

### III. THE LOCALIZATION LENGTH

Random chiral Schrödinger equations of the form (13) have been investigated by A. Comtet, J. Debois, and C. Monthus [13]. It was determined that the important parameter for these equations is  $\frac{1}{j} = \frac{1}{2} \int hV(x) dx$ , and correspondingly  $\frac{1}{j} = \frac{1}{2}$ . For all wavefunctions whose energy  $\frac{1}{j} < \frac{1}{j}$ , the localization length is constant and is given by  $\tilde{\sim} = 1 = \frac{1}{2} \int hV(x) dx$ . For  $\frac{1}{j} > \frac{1}{j}$ , the localization length quickly increases, being asymptotically proportional to  $\frac{1}{j}$ .

The crucial test for our theory is, therefore, whether  $\frac{1}{j} \int hV(x) dx$  is bigger or smaller than  $\frac{1}{2}$ . As we will see below,  $\frac{1}{j} \int hV(x) dx > \frac{1}{2}$ . As a result, Eq. (13) is equivalent

to Eq. (11) and consequently, to Eq. (6), at  $\frac{1}{j} < \frac{1}{j}$ . Thus at  $\frac{1}{j} < \frac{1}{j}$  all the eigenvalues of Eq. (6) are real and positive. On the other hand, there exist some  $\frac{1}{j} > \frac{1}{j}$  where the localization length of  $\tilde{\psi}_n$  becomes equal to  $2j = \frac{1}{2}$ . At that point, the wave functions  $\tilde{\psi}_n$  become delocalized, in accordance with Eq. (14). At  $\frac{1}{j} > \frac{1}{j}$ , the eigenvalues of Eq. (6) are no longer real and come in complex conjugate pairs. This justifies the picture presented on Fig. 1.

In what follows, we proceed to calculate  $\frac{1}{j}$ . In terms of  $V(x)$ , Eq. (9) becomes

$$\frac{dV}{dx} + V^2 - \frac{1}{4j^2} = \frac{\partial^2 h(\psi_0(x); x)}{j} : \quad (15)$$

This equation has the form of a Langevin equation with random force given by  $\partial^2 h$ , whose correlator is given by (see Ref. [5] for justification of  $\partial^2 h$  as a random white noise)

$$\partial^2 h(\psi_0(x); x) \partial^2 h(\psi_0(y); y) = \delta(x - y) :$$

In accordance with the theory of Langevin equations, if

$$\frac{dV}{dx} + g(V) = f(x)$$

with  $f(x)$  being uncorrelated at different values of  $x$ ,  $\langle h(x)f(x^0) \rangle = \delta(x - x^0)$ , then the probability  $P(v; x) = \langle h(v - V(x)) \rangle$  of observing  $V(x) = v$  at the position  $x$  obeys the Fokker-Planck equation

$$\frac{dP}{dt} = \frac{\partial}{\partial v} \frac{\partial}{\partial v} + g(v) P(v; x) :$$

In the present case, we are interested in the probability of observing a particular value for  $V(x)$  together with the probability that a particular solution  $\psi_0(x)$  of Eq. (5) is chosen. The relevant quantity describing this joint probability is

$$P(v; x) = \frac{\langle \psi_0(x) V(x) \rangle}{\langle \psi_0(x) \rangle} ;$$

where  $\langle \psi_0(x) \rangle$  is the density of solutions  $\psi_0(x)$  which in turn obeys the continuity equation

$$\frac{d}{dx} + V = 0 :$$

As a result, we find the Fokker-Planck equation

$$\frac{dP}{dx} = \frac{\partial}{\partial v} \frac{\partial}{\partial v} + V^2 - \frac{1}{4j^2} P + vP :$$

Here  $\frac{1}{j} = 1$ , however, we will keep the more general notation of  $\frac{1}{j}$  for convenience at a later point in the calculations.

A common approach to the Fokker-Planck equations is to map them into the Schrödinger equation with the help of

$$P = \exp \left[ -\frac{v^3 j^2}{3} + \frac{v^2}{4} \right] ;$$

This gives

$$\frac{d}{dx} = -\frac{2j^2}{2j^2} \frac{\partial}{\partial v^2} + U(v) ; \quad (16)$$

Here

$$U(v) = \frac{j^2}{2} v^4 - \frac{1}{4} v^2 - v(1 + \epsilon) + \frac{1}{32} j^2 ; \quad (17)$$

The Feynman path integral formulation of the Schrödinger equation Eq. (16) allows us to find the average of  $V$  using

$$\begin{aligned} \langle V \rangle &= \frac{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} D V(x) e^{S[V]} \int_{-\infty}^{\infty} D V(x) e^{S[V]} dx}{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} D V(x) e^{S[V]} dx} \\ &= \frac{d}{d} \log \int_{-\infty}^{\infty} D V(x) e^{S[V]} ; \quad (18) \end{aligned}$$

for  $\epsilon \ll 1$ . Here  $S[V]$  is the imaginary time action which corresponds to the quantum mechanics Eq. (16),

$$S[V] = \int_0^{\beta} dx \left[ \frac{j^2}{2} \left( \frac{dV}{dx} \right)^2 + U(V) \right] ; \quad (19)$$

At large  $\beta$  the path integral Eq. (18) is dominated by its ground state energy. The average  $V$  may now be calculated as

$$\langle V \rangle = \frac{dE_0}{d} ; \quad (20)$$

where  $E_0$  is the quantum mechanical ground state energy for a particle whose classical action is given by Eq. (19) and whose Schrödinger equation reads

$$-\frac{d^2}{2j^2 dv^2} + U(v) = E$$

If  $\epsilon = 0$ , then the ground state energy, together with its derivative with respect to  $j$ , can be estimated simply from dimensional analysis as  $E_0 = j^2 = j^2$ , to give  $E_0 = j^2 = j^2$ . This agrees with Ref. [9]. If  $\epsilon > 0$ , then dimensional analysis is of no help, since  $E_0$  can now depend on the dimensionless ratio  $\epsilon = (j^2)^{1/3}$ . The only way to find  $E_0$  and thus the localization length of the low lying states is by finding the ground state energy  $E_0$ .

By a suitable rescaling  $v = y \epsilon^{1/3} = j^{2/3}$ ,  $E = E \epsilon^{2/3} = j^{2/3}$ , we can bring the Schrödinger equation to the form

$$\frac{1}{2} \frac{d^2}{dy^2} + \frac{1}{2} y^2 - \epsilon^2 (1 + \epsilon) y = E ; \quad (21)$$

where

$$c = \frac{1}{2(j^2)^{1/3}} ;$$

In general, it is not possible to find the ground state energy of this Schrödinger equation exactly. We are going to find it only in the limit when  $c \ll 1$ . The potential in Eq. (21) has two minima. The minimum at positive  $x$  is the global minimum and is located at

$$y_{\min} = c + \frac{(1 + \epsilon)}{4c^2} + O(\epsilon^5) ; \quad (22)$$

The simplest approximation to the ground state energy is to set it equal to the value of the potential at the minimum. This is given by

$$E_{0,0} = (1 + \epsilon) c \frac{(1 + \epsilon)^2}{8c^2} + O(\epsilon^5) ;$$

To add quantum fluctuations to the problem, we have to consider the Schrödinger equation (21) evaluated around the minimum Eq. (22). Approximating the potential of Eq. (21) by a quadratic potential around the point  $x_{\min}$  we find the oscillator ground state energy to be

$$E_{0,qf} = c + \frac{3(1 + \epsilon)}{8c^2} + O(\epsilon^5) ;$$

In principle, the cubic and quartic term in the expansion of the potential around the minimum also contribute to the ground state energy, but these only have dependence upon  $\epsilon$  at the  $O(\epsilon^5)$  level and thus do not contribute to  $\langle V \rangle$  at lower orders.

Gathering the terms, it is seen that as a function of

$$\begin{aligned} \langle V \rangle &= \frac{dE_0}{d} = \frac{d}{d} E_{0,0} + E_{0,qf} \frac{1}{j^{2/3}} \\ &= \frac{1}{2j} + \frac{1}{2} \frac{1}{j^{2/3}} + O(\epsilon^5) ; \end{aligned}$$

Finally, substituting  $\epsilon = 1$ , we find

$$\langle V \rangle = \frac{1}{2j} + \frac{1}{2} + O(\epsilon^5) ;$$

Therefore, at large  $\beta$  the energy scale  $E_c$  is given by

$$E_c = \frac{1}{2j} ; \quad (23)$$

We see that  $\langle V \rangle > E_c = 1/(2j)$ , and the localization length of the low lying hermitean states  $\xi_n$  is consequently smaller than  $2j = 1/E_c$ . This allows us to deduce that Eq. (6) is indeed equivalent to Eq. (13) at  $\beta < \beta_c$ . The localization length of  $\phi_n$  at  $\beta < \beta_c$  can be deduced from Eq. (14) to be

$$\xi_n = \frac{1}{2j} = \frac{1}{2j} + O(\epsilon^5) ; \quad (24)$$

The calculation presented here justifies Fig. 1 at large  $\beta$ . In order to see that the picture of low lying localized




FIG. 2:  $\langle y \rangle$  in the ground state of Eq. (21) as a function of  $c$ . The dots represent  $\langle y \rangle$  evaluated numerically at various values of  $c$ . The solid line is  $\langle y \rangle$  calculated in the harmonic approximation to the potential in Eq. (21). The dash-dotted line represents the first two terms of the perturbative calculation  $c + \frac{1}{8c}$ . Finally, the dashed line is a straight line of the slope 1, which demonstrates that  $\langle y \rangle > c$ .

states persist at all values of  $\alpha$  we need to show that  $\langle y \rangle$  is always greater than  $\alpha = 2j$ . This amounts to showing that in the Schrodinger equation Eq. (21) the following relation always holds true

$$\langle y \rangle = \int_{-\infty}^{\infty} dy j_0(y) j_1(y) = \frac{dE_0}{d\alpha} \Big|_{\alpha=1} > c; \quad (25)$$

Here  $j_0(x)$  is the ground state wave function of Eq. (21), at  $\alpha = 1$ . We do not know how to show this analytically. We investigated Eq. (21) numerically. Fig. 2 shows the values of  $dE_0/d\alpha = \langle y \rangle$ , at  $\alpha = 1$ , plotted versus  $c$ . We see that Eq. (25) does seem to hold.

#### IV. THE LARKIN LENGTH

In the analysis of Eq. (5) it is customary to introduce the notion of Larkin length  $L$ .  $L$  is defined as size of a box in which the average  $\langle y^2 \rangle$  becomes of the order of 1. Thus at distances bigger than  $L$ ,  $j_0(x)$  becomes rough, while at shorter distances it can be thought of as smooth. We are now going to see that in our problem the localization length of low lying states of Eq. (6) is of the order of Larkin length.

To calculate the Larkin length, we follow the procedure described in Ref. [10]. We consider the equation Eq. (5). Let us assume that  $\alpha$  is small and neglect the dependence of  $h$ . In Fourier space this equation of motion is

$$(i k + j k^2) \phi_k = \alpha h(0);$$

Thus

$$\frac{D}{j k j^2} E = \frac{\hbar \alpha h(0) \phi h(0) i}{2k^2 + j^2 k^4};$$

Now the Larkin length is defined as the value of  $L$  for which

$$1 = \int_0^L \langle y^2(x) \rangle dx$$

$$= \int_{-\infty}^{\infty} \frac{dk}{2} \frac{1}{2k^2 + j^2 k^4}$$

$$= \frac{L}{2} \int_{-\infty}^{\infty} \frac{dk}{3} \tan^{-1}(L k) = j):$$

From the asymptotic form of  $\tan^{-1}$  in the large- $L$  limit it is found that

$$L = \frac{2}{j} + \frac{1}{2} j^{-1} j^2 k^4 + O(j^{-7});$$

At leading order  $L$  turns out to be proportional to the localization length obtained above Eq. (24). This coincides with the behavior of the pinned charge density waves without the background current [9]. However, the localization length and the Larkin length have different functional dependence on  $\alpha$  in lower orders.

#### V. DENSITY OF STATES

In this section we calculate the density of low lying states for our problem. Due to the form of the potential Eq. (17) which does not contain a cubic term, the density of low lying states may be easily calculated using the methods of ref. [5]. This is not immediately obvious, so in this section we will repeat the argument.

At energies below an energy of the order of  $\langle y \rangle$  the energy eigenvalues are real. Above, the eigenvalues trace out a one-dimensional spectrum in the complex plane, Fig. 1. Let us concentrate on low lying states  $\epsilon_n < \langle y \rangle$ , which are all real. To calculate their density of states  $\rho(\epsilon)$  we use the equivalence of Eq. (6) and Eq. (13) in this regime.

The integrated density of low lying states can be found using the discussion in Ref. [5] as

$$N(\epsilon) = \int_{x_1}^{x_2} dx V(x) + a; \quad (26)$$

$$a = \int_{x_1}^{x_2} \log(|\epsilon|);$$

with the average being over realizations of  $V(x)$ . The density of states is then obtained from  $\rho(\epsilon) = dN/d\epsilon$ .

Through a standard representation of the  $\delta$ -function

$$N(\epsilon) = \int_{-\infty}^{\infty} \frac{d\epsilon}{2} e^{i \int_{x_1}^{x_2} dx V(x) + a};$$

and defining  $S(\epsilon)$  by

$$\exp(-S(\epsilon)) = \exp(-i \int_{x_1}^{x_2} dx V(x)); \quad (26)$$

with  $\epsilon = x_2 - x_1$  the integrated density of states is

$$N(\epsilon) = \int_{-\infty}^{\infty} \frac{d\epsilon}{2} e^{i a - S(\epsilon)};$$

At large  $\epsilon$ ,  $S(\epsilon)$  is independent of  $\epsilon$  as will be argued below. We assume in the following that a certain value



lying states has also been obtained, giving the power law  $(\lambda)^{-1/3}$ . This result differs from the result  $(\lambda)^{-1/4}$  of Refs. [7, 9] due to the fact that turning on background current removes the notion of a global ground state.

Acknowledgments The authors wish to thank Leo Radzihovsky for useful discussions. J.L. also wishes to acknowledge support from the Danish Research Agency and the Danish-American Fulbright Commission.

- 
- [1] P.W. Anderson, Phys. Rev. 109, 1492 (1958).
  - [2] K.B. Efetov, Supersymmetry in Disorder and Chaos, Cambridge University Press, 1997.
  - [3] A.A. Itland and M.R. Zimbauer, Phys. Rev. B 55, 1142 (1997).
  - [4] M.R. Zimbauer, J. Math. Phys. 37, 4986 (1996).
  - [5] V. Gurarie and J.T. Chalker, Phys. Rev. B 68, 134207 (2003).
  - [6] H. Fukuyama and P.A. Lee, Phys. Rev. B 17, 535, (1978).
  - [7] I.L. Aleiner and I.M. Ruzin, Phys. Rev. Lett. 72, 1056 (1994).
  - [8] M. Fogler, Phys. Rev. Lett. 88, 186402 (2002).
  - [9] V. Gurarie and J.T. Chalker, Phys. Rev. Lett. 89, 136801 (2002).
  - [10] L. Radzihovsky and J. Toner, Phys. Rev. Lett., 81, 3711 (1998).
  - [11] I.Ya. Golsheid, B.A. Khonuzhenko, cond-mat/9707230.
  - [12] N. Hatano and D.R. Nelson, Phys. Rev. Lett. 77, 570 (1996).
  - [13] A. Comtet, J. Debois, and C. Monthus, Ann. Phys. 239, 312 (1995).
  - [14] S. Brazovskii, T. Nattermann, cond-mat/0312375