

The spin and charge gaps of the half-filled N-leg Kondo ladders

J. C. Xavier

Instituto de Física Gleb Wataghin, Unicamp, C.P. 6165, 13083-970 Campinas SP, Brazil

(Dated: 14th April 2024)

In this work, we study N-leg Kondo ladders at half-filling through the density matrix renormalization group. We found non-zero spin and charge gaps for any finite number of legs and Kondo coupling $J > 0$. We also show evidence of the existence of a quantum critical point in the two dimensional Kondo lattice model, in agreement with previous works. Based on the binding energy of two holes, we did not find evidence of superconductivity in the 2D Kondo lattice model close to half-filling.

PACS numbers: 75.30.Mb, 71.10.Pm, 71.30.+h, 75.10.-b

Recently, a great deal of excitement has been generated with the discovery of several heavy fermion compounds which exhibit a quantum critical point (QCP)^{1,2}. Several anomalous behaviors have been observed close to the QCP and one of the main problems of this area is to try to understand this puzzle. The Kondo lattice model (KLM) is the simplest model believed to describe heavy fermion materials³. This model incorporates the interaction between the localized spins and the conduction electrons via an exchange interaction of strength J . This interaction is believed to be the most important in heavy fermion compounds at low temperatures. At small values of J , due to the indirect Ruderman-Kittel-Kasuya-Yosida (RKKY) interaction, antiferromagnetic long-range order (LRO) is expected, whereas for large J a paramagnetic state emerges. The competition between these two states leads to the QCP, as first pointed out by Doniach⁴. Many theoretical works have concentrated on the one-dimensional (1D) model, which can be treated more easily. In particular, at half-filling, several numerical calculations show that the 1D KLM is always a gapped spin liquid with no magnetic quasi-LRO^{5,6,7}. Away from half-filling, there is power law decay of spin correlations, for small values of J (see, however, Ref. 8). On the other hand, for the two-dimensional (2D) KLM at half-filling, several studies support the existence of a QCP^{9,10,11} (see also Ref. 12).

In this work, we present the spin and charge gaps of the N-leg Kondo ladders at half-filling calculated by the density matrix renormalization group (DMRG) technique¹³. The N-leg Kondo ladders consist of N Kondo chains of length L coupled by the hopping term. The 2D system is obtained by taking both N and L to infinity. As far as we know this is the first study of the N-leg Kondo ladders. We will also show evidence of the existence of a QCP in the 2D Kondo model at half-filling in agreement with previous studies. Moreover, our results for the binding energy of two holes do not show any evidence for an effective hole-hole attraction in the 2D KLM for any Kondo coupling. All the results we will report here correspond to the conduction electron density $n = 1$. It is important to note that for this density, techniques other than the DMRG can be applied^{9,10,11}. However, away from half-filling, methods like Quantum Monte Carlo¹¹

are plagued by the “sign problem”. Therefore, our results at half-filling should serve as benchmarks studies before get results at more general fillings. DMRG results away from half-filling are in progress and will be presented elsewhere¹⁴.

We considered the N-leg Kondo Hamiltonian at half-filling with $M = N \times L$ sites

$$H = \sum_{\langle ij \rangle} (c_{ij}^\dagger c_j + H.c.) + J \sum_j S_j \cdot s_j$$

where c_j annihilates a conduction electron in site j with spin projection $\frac{1}{2}$, S_j is a localized spin $\frac{1}{2}$ operator, $s_j = \frac{1}{2} \sum_{\alpha\beta} c_{j\alpha}^\dagger c_{j\beta} \sigma_{\alpha\beta}$ is the conduction electron spin density operator and $\sigma_{\alpha\beta}$ are Pauli matrices. Here $\langle ij \rangle$ denote near-neighbor sites, $J > 0$ is the Kondo coupling constant between the conduction electrons and the local moments and the hopping amplitude was set to unity to fix the energy scale. We investigated the model with the DMRG technique with open boundary conditions. We used the finite-size algorithm for sizes up to $N \times L = 40$, keeping up to $m = 2000$ states per block. The discarded weight was typically about 10^{-4} to 10^{-8} in the final sweep. Although the DMRG is based on a one-dimensional algorithm, it has been applied to low dimensional systems. The procedure consists in mapping the low-dimensional model on a 1D model with long range interactions^{15,16}. Several important questions, such as the existence of striped phases and/or superconductivity in the 2D $t - J$ model, have been addressed with this procedure¹⁷. Here, we used this same method to study the N-leg Kondo ladders.

Our first concern is to check the applicability of the technique to the N-leg Kondo ladders. We can get some insight by looking at the ground state energies. We have checked that, for small slightly doped clusters with up to 12 sites, we can reproduce with the DMRG the same energies obtained independently by exact diagonalization (ED). As we increase the system size the Hilbert space increases exponentially and ED is no longer a suitable method to treat the system. This is the main advantage of the DMRG. In order to illustrate this advantage, we will show that we can obtain the ground state energies of the N-leg Kondo ladders in a controlled way¹⁸.

	2x10		4x4		
m	J=0.0	J=2.0	J=0.0	J=0.3	J=3.0
100	-27.56929	-39.15411	-21.85340	-22.12936	-41.11699
200	-27.62807	-39.15909	-21.88557	-22.19189	-41.25039
300	-27.63982	-39.15988	-21.88816	-22.25239	-41.28139
400	-27.64290	-39.16007	-21.88845	-22.29403	-41.29605
600	-27.64407	-39.16015	-21.88853	-22.33682	-41.31698
800				-22.36031	-41.32227
1200				-22.38726	-41.32606
1	-27.6455	-39.1602	-21.8885	-22.4413	-41.3401
exact	-27.64486		-21.88854		

Table I: The ground state energies of the Kondo lattice model for the clusters 2x10 and 4x4 as a function of the number of states m kept in the DMRG truncation for some Kondo couplings. The energies at $m = 1$ were obtained by an extrapolation, see text.

In table 1 we present the ground state energies as a function of the truncation m for the 2-leg (4-leg) Kondo ladders with length $L = 10$ ($L = 4$). The energies for $m = 1$ were obtained by an extrapolation assuming a dependence of the type $e(m) = e_1 + a/m + b/m^2$. As we can see, for the $J = 0$ (free case) with a small number m of states the relative error is smaller than 10^{-4} . This free case result does not appear to depend on the number of legs, as shown for the 2- and 4-leg ladders. However, a small coupling J makes the wave function much more complex and a larger number of states is necessary to keep the same accuracy. Furthermore, in this weak coupling regime, we need a much larger number of states (and consequently a longer computational time) to keep the algorithm stable as the number of legs is increased. An increase in computer effort, probably exponential, such as this has also been observed in other systems^{15,16}. This was the main reason why we restricted our calculations to $N \leq 6$ and large values of J . On the other hand, in the strong coupling limit, a smaller number of states is necessary to keep the same accuracy. For $J \rightarrow 0$ the conduction electrons form unbreakable mobile singlets with the localized spins, effectively behaving like free spinless fermions.

Having set these benchmarks, we are now able to calculate the spin and charge gaps of the N -leg Kondo ladders with $M = N \times L$ sites at half-filling. These are defined as $\Delta_s(L) = e(M; 1) - e(M; 0)$ and $\Delta_c(L) = e(M; 2; 0) - e(0; 0)$, respectively, where $e(N_c; s)$ is the ground state energy of the system with N_c conduction electrons in the subspace of total spin s ^{7,19}. In Fig. 1(a) we present the spin and charge gaps for the 2-leg Kondo ladder with length $L = 10$ (open symbols) and $L = 20$ (closed symbols) as a function of the Kondo coupling J . As we see, finite size effects are very small. In the strong-coupling limit the spin and charge gaps increase linearly with J , and can be estimated by second order perturba-




Figure 1: (a) The spin and charge gaps for the 2-leg Kondo ladder with $L = 10$ (open symbols) and $L = 20$ (closed symbols). The circles (squares) correspond to the spin (charge) gap. Inset: zoom of the region with $0 < J < 2$. (b) Same as (a), but for the 3-leg Kondo ladder with $L = 8$ (open symbols) and $L = 14$ (closed symbols).

tion theory with respect to $t \sim J^7$. At this order, the spin and charge gaps of the N -leg Kondo ladders are given, respectively, by $\Delta_s^{(2)} = J \frac{20}{3J} t^2$ and $\Delta_c^{(2)} = 3J \frac{2}{3} t^2 + N \left(\frac{4}{3} - 1 \right) \frac{t^2}{J}$; where $N = 1 \cos(N = N + 1)$. We used the infinite-size algorithm¹³ to determine the spin and charge gaps of the 2-leg Kondo ladders in the strong coupling limit²⁰. Our numerical calculations are in very good agreement with these results. For example, for $J = 40$ we obtained $\Delta_s = 39.752$ and $\Delta_c = 57.057$, very close to the perturbative results $\Delta_s^{(2)} = 39.750$ and $\Delta_c^{(2)} = 57.037$. On the other hand, for small values of J the spin gap has a different dependence on the coupling constant.

The results we have found for the 2-leg Kondo ladder are very similar to those of the 1D KLM⁷. In this case, for large J the spin gap goes linearly with J , while for small J it decays exponentially as $\Delta_s \sim \exp(-1/nJ)$, where $1/4$ ⁹. This kind of behavior does not seem to depend on the number of legs. The spin and charge

gaps of the 3-leg Kondo ladder show similar behavior, as shown in Fig. 1(b). Based on the spin and charge gap behaviors of the 1-, 2- and 3-leg Kondo ladders, we expect *non-zero spin and charge gaps* for the half-filled N -leg Kondo ladders. It is interesting to note that the spin gap has similar behavior whether the number of legs is even or odd. The same does not happen in N -leg Heisenberg ladders, where the spin gap is zero for odd N and different from zero for even N ²¹ (for a review see Ref. 22). This difference arises from the fact that the Kondo Ladders are coupled just with the hopping term, differently from the Heisenberg Ladders. So, the same analysis that is done in the N -leg Heisenberg ladders (see Ref. 22), does not apply in the N -leg Kondo ladders. Moreover, we might naively expect that the inclusion of the Coulomb interaction U in the N -leg Kondo ladders would lead to a spin gap with a distinct dependence on the number of legs (at least for $U \neq 0$), as in the Heisenberg ladders. However, most probably, even the presence of U will not lead to this distinct behavior. Note that the 1D Kondo-Hubbard model at half-filling has a non-zero spin gap¹⁹, differently from the 1D Heisenberg model.

We would still like to determine whether the spin gap remains finite or not when N goes to infinity (after $L \rightarrow \infty$). Approximate approaches^{9,10} and Quantum Monte Carlo results¹¹ support the existence of a QCP at $J_c \approx 1.45 \pm 0.05$. Below this value antiferromagnetic LRO appears with a zero spin gap. It is a very tough task to determine the presence of LRO directly by the value of the spin gap, since for small values of J the gap is very small, and as we increase the number of legs we lose accuracy in the energies.

Instead of looking at the spin gap to check for LRO, we can calculate the Fourier transform of the spin-spin correlation function $S(\mathbf{q}) = \frac{1}{M} \sum_{\mathbf{r}_1, \mathbf{r}_2} e^{i\mathbf{q} \cdot (\mathbf{r}_1 - \mathbf{r}_2)} \langle S_{\mathbf{r}_1} S_{\mathbf{r}_2} \rangle$. In Fig. 2(a), we show the spin structure factor $S(\mathbf{q})$ for the 2-leg Kondo ladder as a function of momentum q_x for $J = 0.5$. The peak at $\mathbf{q} = (\pi; 0)$ is a signature of some kind of antiferromagnetic order, not necessarily long-ranged. For small values of J we expect this peak to increase as we decrease J , since the RKKY interaction starts to dominate over Kondo compensation. Indeed, this is what we have seen. In Fig. 2(b), we present the spin structure factor $S(\mathbf{q} = (\pi; 0))$, for the clusters 2×10 (squares) and 3×8 (circles), as a function of the Kondo coupling J . Note the divergence as $J \rightarrow 0$. This kind of behavior was also observed in the magnetic susceptibility of the one-dimensional KLM⁵. Antiferromagnetic LRO exists if $\lim_{M \rightarrow \infty} S(\mathbf{q} = (\pi; 0))/M$ is non-zero. In Fig. 2(c), we present $S(\mathbf{q} = (\pi; 0))/M$ as a function of $M^{1/2}$ for several values of the Kondo coupling. As can be seen, even working with small clusters, our results suggest that for $J > 1.6$ $S(\mathbf{q} = (\pi; 0))/M$ tends to zero as we increase M , while for $J < 1.3$ it tends to a finite value. This result is compatible with $J_c = 1.45$ as suggested by other methods^{9,10,11}.

Finally, we would like to check for evidence of superconductivity in the 2D KLM close to the QCP. Several




Figure 2: The spin structure factor $S(\mathbf{q})$ for the N -leg Kondo ladders. (a) $S(\mathbf{q})$ vs momentum q_x for the 2-leg Kondo ladder with length $L = 10$ and $J = 0.5$. (b) $S(\mathbf{q} = (\pi; 0))$ vs the Kondo coupling J for the clusters 2×10 (squares) and 3×8 (circles). (c) $S(\mathbf{q} = (\pi; 0))/M$ vs $M^{1/2}$ for several Kondo couplings. The sizes of the clusters used were $M = 2 \times 2, 3 \times 4, 4 \times 4, 4 \times 6$, and 6×6 .




Figure 3: The binding energy Δ_B for the 1-, 2- and 3-leg Kondo ladders as a function of the Kondo coupling J .

heavy fermions compounds, like CeIn_3 and CePd_2Si_2 , can be tuned to an antiferromagnetic QCP by application of external pressure²³. Superconductivity was observed at low temperatures close to the QCP of these materials. Our aim will not be to find proof of superconductivity at finite hole doping close to a quantum critical point in the 2D KLM, which would entail the calculation of the pair correlation function. Instead, we can gain some insight by looking at the binding energy of two doped holes, defined as $\Delta_B = E(2) + E(0) - 2E(1)$, where $E(n)$ is the

ground state energy with n holes. If the holes are bound $\epsilon_B < 0$, and this is indicative that attractive effective forces are present²⁴. This is a necessary, though not sufficient, condition for a possible condensation of the holes. In Fig. 3, we present the hole binding energy for the 1-, 2- and 3-leg Kondo ladders. In the thermodynamic limit the binding energies vanish if holes do not form bound states and is asymptotically positive²⁴. As we see from this figure, this is exactly what we have observed. Increasing the cluster from 2×10 to 2×20 the binding energies decrease. As we see, the holes do not bind for *any* Kondo coupling. Similar results were also observed in the Kondo chain⁵. These results suggest that, at least close to half-filling, superconductivity will not emerge in the 2D Kondo lattice model. The fact that we have not found bind holes also

suggest the the KLM may not be the minimum model to describe correctly the Heavy Fermion compounds.

In summary, we presented, for the first time, a systematic study of the 2- and 3-leg Kondo ladders at half-filling by using the DMRG. Based on the behavior of the spin gap of the 1- to 3-leg Kondo ladders, we expect non-zero charge and spin gaps in the N -leg Kondo ladders for any $J > 0$. Using small clusters we were able to determine the presence of a QCP, in agreement with previous works. Our results also suggest that there is no superconductivity close to half-filling in the 2D Kondo lattice model.

The author thanks E. Miranda, E. Dagotto, P. G. Pagliuso, A. H. Castro-Neto, and R. R. dos Santos for useful discussions. This work was supported by FAPESP 00/02802-7 and 01/00719-8.

-
- ¹ S. R. Stewart, Rev. Mod. Phys. **73**, 797 (2001).
 - ² P. Coleman *et al.*, J. Phys.:Condens. Matter **13**, R723 (2001).
 - ³ A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, 1997).
 - ⁴ S. Doniach, Physica B **91**, 231 (1977).
 - ⁵ C. C. Yu and S. R. White, Phys. Rev. Lett. **71**, 3866 (1993).
 - ⁶ H. Tsunetsugu, Y. Hatsugai, K. Ueda, and M. Sigrist, Phys. Rev. B **46**, R3175 (1992); R. M. Fye and D. J. Scalapino, Phys. Rev. Lett. **65**, 3177 (1990).
 - ⁷ H. Tsunetsugu, M. Sigrist, and K. Ueda, Rev. Mod. Phys. **69**, 809 (1997).
 - ⁸ J. C. Xavier, R. G. Pereira, E. Miranda, and I. Affleck, Phys. Rev. Lett. **90**, 247204 (2003).
 - ⁹ Z. Wang, X. P. Li and D. H. Lee, Physica B **199-200**, 436 (1994).
 - ¹⁰ Z. P. Shi, R. R. P. Singh, M. P. Gelfand, and Z. Wang, Phys. Rev. B **51**, 15630 (1995).
 - ¹¹ F. Assaad, Phys. Rev. Lett. **83**, 796 (1999).
 - ¹² K. S. Beach, P. A. Lee, and P. Monthoux, cond-mat/0306336.
 - ¹³ S. R. White, Phys. Rev. Lett. **69**, 2863 (1993); Phys. Rev. B **48**, 10345 (1993).
 - ¹⁴ J. C. Xavier, E. Miranda, and E. Dagotto, unpublished.
 - ¹⁵ S. Liang and H. Pang, Phys. Rev. B **49**, 9214 (1994).
 - ¹⁶ S. R. White, Phys. Rev. Lett. **77**, 3633 (1996).
 - ¹⁷ R. M. Noack, S. R. White, and D. J. Scalapino, Phys. Rev. Lett. **73**, 882 (1994); S. White and D. Scalapino, Phys. Rev. Lett. **80**, 1272 (1998); G. B. Martins, C. Gazza, J. C. Xavier, A. Feiguin, and E. Dagotto Phys. Rev. Lett. **84**, 5844 (2000).
 - ¹⁸ In the DMRG it is believed that the energy error is proportional to the truncation error¹³. However, at the present moment, there is no exact relation between them. A possible way to estimate the relative error is the following. We define the relative error as $(e(m=1) - e(m_{last})) / e(m=1)$, where $e(m=1)$ is the extrapolated energy (see text) and $e(m_{last})$ is the energy with the biggest m available. In all results presented here, the relative error is smaller than $0.4 \cdot 10^{-4}$, $0.2 \cdot 10^{-3}$, $0.3 \cdot 10^{-3}$ and $0.7 \cdot 10^{-3}$ for the 2-, 3-, 4- and 6-leg Kondo ladders respectively.
 - ¹⁹ N. Shibata, T. Nishino, K. Ueda, and C. Ishii, Phys. Rev. B **53**, R8828 (1996).
 - ²⁰ The infinite-size algorithm is very stable for large J . For small values of J we were not able to get precise results using the same procedure.
 - ²¹ E. Dagotto, J. Riera, and D. Scalapino, Phys. Rev. B **45**, 5744 (1992); S. R. White, R. M. Noack and D. J. Scalapino, Phys. Rev. Lett. **73**, 886 (1994); G. Sierra, J. Phys. A: Math. Gen. **29**, 3299 (1996).
 - ²² E. Dagotto and T. M. Rice, Science **271**, 618 (1996).
 - ²³ N. D. Mathur, F. M. Grosche, S. R. Julian, I. R. Walker, D. M. Freye, R. K. W. Haselwimmer, and G. G. Lonzarich, Nature **394**, 39 (1998).
 - ²⁴ E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).