

A new numerical algorithm for the double-orbital Hubbard model | Hund-coupled pairing symmetry in the doped case

Shiro Sakai, Ryotaro Arita, Hideo Aoki

Department of Physics, University of Tokyo, Hongo, Tokyo 113-0033, Japan
(Dated: April 14, 2024)

In order to numerically study electron correlation effects in multi-orbital systems, we propose a new type of discrete transformation for the exchange (Hund's coupling) and pair-hopping interactions to be used in the dynamical mean-field theory + quantum Monte Carlo method. The transformation, which is real and exact, turns out to suppress the sign problem in a wide parameter region including non-half-filled bands. This enables us to obtain the dominant pairing symmetry in the double-orbital Hubbard model, which shows that the spin-triplet, orbital-antisymmetric pairing that exploits Hund's coupling is stable in a wide region of the band filling.

PACS numbers: 71.10.Fd, 74.20.Rp, 71.30.+h

In the physics of correlated electron systems, as highlighted by the high- T_c superconductivity, the prototypical transition-metal oxides have arrested attention because of a rich variety of physical properties, among which are anisotropic pairing symmetries in the cuprates and in Sr_2RuO_4 , the colossal magnetoresistance in manganites, and a complex phase diagram for $\text{La}_{1-x}\text{Sr}_x\text{MnO}_3$. [1] Since the d-orbitals are relevant, we are actually talking about multi-orbital systems, and, while some of the phenomena should be generically captured within single-band models, understanding of the effect of electron correlation in multi-orbital system remains a fundamental problem. [2]

Namely, the orbital degrees of freedom should couple to other degrees of freedom such as charge, spin, and lattice distortion, and we can expect even richer physical properties. Indeed, the colossal magnetoresistance results from Hund's coupling and the complex phase diagrams of Mn and Co compounds are considered to arise from a competition between Hund's coupling and the Jahn-Teller distortion. In perovskite-type oxides, the crystal field splits the ve 3d orbitals into three-fold degenerate (t_g) and two-fold degenerate (e_g) levels. When the degenerate levels are not fully filled, the degeneracy may be lifted by the Jahn-Teller effect, where the system is effectively mapped to a single-orbital model when the splitting is large enough. We take here the small splitting limit to concentrate on the physics specifically caused by the orbital degrees of freedom.

Both numerical and analytic methods have been developed to study correlated electron systems. The methods should preferably be non-perturbative if one wants to examine the effect specific to the electron correlation such as Mott's metal-insulator transition. The dynamical mean-field theory (DMFT) [3], which can fully include temporal fluctuations while spatial fluctuations are neglected, first succeeded in describing this transition both from metallic and insulating sides. In this method a lattice system (such as the Hubbard model) is mapped to an impurity model, which becomes an exact mapping in

the limit of infinite spatial dimension. A standard procedure is to solve the impurity problem with the (auxiliary-field) quantum Monte Carlo (QMC) method [4, 5], which involves no approximations except for the Trotter decomposition. So the DMFT+QMC method should be a desirable candidate for the multi-orbital cases. Unfortunately, it is difficult to extend the QMC method to multi-orbitals: First, it is impossible to express those (exchange and pair-hopping) interactions that are specific to multi-orbital cases in terms of the usual auxiliary fields. Second, even when we can accomplish this, the negative sign problem, a notorious problem in QMC calculations, is usually difficult to avoid for multi-orbitals.

This has motivated us to propose here a new auxiliary-field transformation that is applicable to Hund's and pair-transfer terms. The transformation, which is real and discrete, turns out to suppress the sign problem in a wide parameter region including non-half-filled bands. This enables us to examine the role of Hund's coupling and pair-transfer in a double-orbital model. One of the most intriguing questions for correlated electrons on multi-orbitals is what should be the symmetry of the superconducting pairing that arises from the electron-electron interaction. We have examined the dominant pairing symmetry in the double-orbital Hubbard model with the DMFT+QMC method. The result reveals that the spin-triplet orbital-antisymmetric even-frequency pairing, which exploits Hund's coupling, is stable in a wide region of the band filling.

So we take the two-fold degenerate Hubbard model [6] with a Hamiltonian,

$$\begin{aligned} H &= H_0 + H_1 + H_2; \\ H_0 &= \sum_{i,j} \sum_{\sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + \sum_{i,j} \sum_{\sigma} X_{ij}^\dagger c_{i\sigma}^\dagger c_{j\sigma}^\dagger; \\ H_1 &= U \sum_{i,j} n_{i\uparrow} n_{i\downarrow} + U^0 \sum_i n_{i1} n_{i2} \\ &\quad + (U^0 - J) \sum_i n_{i1} n_{i2}; \end{aligned} \quad (1)$$

$$H_2 = J \sum_{i,m \in m^0} \sum_{n \in m^0} (c_{im}^\dagger c_{in}^\dagger c_{im} c_{in} + c_{im}^\dagger c_{in}^\dagger c_{im} c_{in});$$

Here c_{im}^\dagger creates an electron of spin \uparrow in the orbital m ($\in 1;2$) at site i , and $n_{im} = c_{im}^\dagger c_{im}$. We only consider the nearest-neighbor hopping between the similar orbitals, electron-electron interactions are assumed to be intra-atomic with the intra-(inter-)orbital Coulomb interaction denoted as U (U^0), while the exchange and pair-hopping interactions as J . The Hamiltonian is rotationally invariant not only in spin, but also in real space if we fulfill the condition $U = U^0 + 2J$ (as is the case with d-orbitals). We have divided the interaction into the density-density interactions H_1 and the exchange and pair-hopping interactions H_2 .

The DMFT+QMC method has been used by many authors for the single-orbital Hubbard model. For the usual on-site Hubbard interaction, a decoupling is done with the discrete Hubbard-Stratonovich (HS) transformation [7],

$$e^{-a[n_\uparrow n_\downarrow - \frac{1}{2}(n_\uparrow + n_\downarrow)]} = \frac{1}{2} \int_{-1}^1 ds e^{s(n_\uparrow - n_\downarrow)} e^{a(s^2 - 1)}; \quad (2)$$

with $\log(e^{\frac{a}{2}} + e^{\frac{a}{2}}) = \frac{1}{2} \log(e^{2a} + 1)$, which transforms the two-body interaction into one-body interactions summed over an auxiliary field s . Applying this to each interaction term on the discretized imaginary time, we can decompose the partition function Z of the many-body system into a sum of the partition functions Z_{fsig} of one-body systems as $Z = \int_{fsig} Z_{fsig}$. The QMC samples the single-particle system s according to the weight Z_{fsig} . The negative sign refers to the fact that the weight is not positive-definite. The sign problem does not occur, as far as the DMFT is concerned, in the single-orbital Hubbard model because the impurity problem lacks the electron hopping terms.

While the auxiliary-field QMC method has been applied to some multi-orbital Hubbard models by neglecting the terms other than the density-density interactions (H_1 in eq.(1)), the QMC algorithm becomes a challenging problem for the exchange and pair-hopping interactions (H_2): the HS transformation (2) is obviously inapplicable to these terms. While a decoupling, $\exp(J c_{1m}^\dagger c_{2m}^\dagger c_{1m} c_{2m}) = (1/2) \int_{-1}^1 ds \exp[s J (c_{1m}^\dagger c_{2m}^\dagger - c_{1m} c_{2m})]$; is possible after breaking e^{-H_2} into a product of exponentials, it leads to a serious sign problem [8]. Another attempt by Motome and Imada [9] decouples H_2 with imaginary auxiliary fields to implement a QMC, but an electron-hole symmetry has to be assumed to avoid phase cancellation of the weights which become complex due to the imaginary auxiliary fields. Since the assumption dictates the half-filled band and $U = U^0$ (that violates $U = U^0 + 2J$), investigation of non-half-filled bands and/or the rotationally symmetric case is difficult with this method.




FIG. 1: The average sign plotted as a function of band filling at $\beta = 2$ for $U = U^0 = 1$; $J = 0.5$ and at $\beta = 10$ for $U = U^0 = 1$; $J = 0.2$ calculated with the present algorithm (circles) and with the one due to [9] (squares). In this particular plot we have set $U = U^0$ to facilitate comparison with Motome and Imada's method. [11] Curves are guides to the eye here and in subsequent figures.

We propose here a new type of discrete transformation for the exchange and pair-hopping interaction term s ,

$$e^{-H_2} = \frac{1}{2} \sum_{r=1}^X e^{r(f_\uparrow - f_\downarrow)} e^{a(N_\uparrow + N_\downarrow) + bN_\uparrow N_\downarrow}; \quad (3)$$

where

$$\begin{aligned} a &= \frac{1}{2} \log(e^{2J} + e^{4J} - 1); \\ b &= \log(\cosh(J)); \\ f &= c_1^\dagger c_2 + c_2^\dagger c_1; \quad N = n_1 + n_2 - 2n_1 n_2; \end{aligned}$$

This transformation is exact with the auxiliary field r being real and all the operators being hermitian. Although a term $N_\uparrow N_\downarrow$ which is forth order in n appears on the right hand side, we can apply the usual HS transformation to this term due to a property $N^2 = N$. The resulting terms in the form nn can be combined with the Coulomb terms, which are transformed with eq.(2), so that we need in total only two auxiliary fields for H_2 . Furthermore the interaction parameters $U; U^0; J$ can be varied independently, which means that we can treat the rotational symmetric cases of $U = U^0 + 2J$. We note that recently Han [10] proposed another type of HS transformation for the Hamiltonian (1). He adopts a continuous HS transformation in combination with the discrete one (2), and avoided the sign problem in a wide parameter region at half-filling. In contrast all the transformation are discrete in the present method.

One valuable property of the transformation (3) is that negative weights are indeed reduced significantly. An unexpected asset is that the suppression of the sign problem is almost independent of the band filling. This is shown in Fig.1, where the average sign $\langle \text{sign} \rangle$




FIG. 3: Temperature dependence of the pairing susceptibilities for $n = 1.8$, $U^0 = 0.7$, $J = 0.4$, $U = U^0 + 2J = 1.5$. The symbols denote pairing symmetries (1: spin-singlet, 3: triplet; $S^{a,b,c}$: orbital-symmetric (eq.(4)), A: antisymmetric; E: even-, O: odd-frequency).




FIG. 4: Band-filling dependence of the pairing susceptibilities for $n = 60$, $U^0 = 0.7$, $J = 0.4$, $U = U^0 + 2J = 1.5$ with the same abbreviation for pairing symmetries as in Fig.3.

enhancement of such a pairing should be due to Hund's coupling, which tends to align electron spins across different orbitals. The result is consistent with Han's[10] for the semi-elliptical density of states. While in the work of Han the susceptibility diverges at $T=W = 0.06$ for $J=W = 0.15$; $U^0=W = 0.45$, we do not detect the divergence up to $T=W = 1=180$. This should be because electrons are less correlated in the hypercubic lattice than in the Bethe lattice (which the semi-elliptical density of states would represent), since the Gaussian density of states for the hypercubic lattice, with high-energy tails, has an effectively larger band width.

Figure 4 shows the band-filling dependence of the pairing susceptibilities at $T=W = 1=120$ for $U^0 = 0.7$, $J = 0.4$,

and $U = U^0 + 2J = 1.5$. A new finding here is that the spin-triplet, orbital-antisymmetric (3AE) pairing induced by Hund's coupling remains dominant in a rather wide range ($n = 1.6 - 2$) of hole-doping at this temperature. It declines, however, at low fillings ($n = 1 - 1.4$). This result may be related to a mechanism of superconductivity proposed by Capone[14] for multi-orbital systems close to the Mott transition, where the electron repulsion U is envisaged to assist a pairing even for s-waves.

In summary, we have constructed a new discrete transformation (which reduces the sign problem and is applicable to doped bands) for the exchange and pair-hopping terms in double-orbital models, and implemented this in a DMFT+QMC calculation. Superconducting susceptibilities of the s-wave pairings calculated with this method shows that the spin-triplet, orbital-antisymmetric even frequency pairing, enhanced by Hund's coupling and being dominant at half-filling, is robust against hole doping. We note that it is difficult to investigate ferromagnetism in this model even with the present transformation, since the ferromagnetism appears only in very strongly correlated regime[15], where the sign problem becomes serious again, so this is a future problem.

Numerical calculations were partly performed on SR8000 in ISSP, University of Tokyo.

-
- [1] Y. Tokura and N. Nagaosa, *Science* 288, 462 (2000).
 - [2] L.M. Roth, *Phys. Rev.* 149, 306 (1966).
 - [3] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Rev. Mod. Phys.* 68, 13 (1996).
 - [4] J.E. Hirsch, *Phys. Rev. B* 31, 4403 (1985); J.E. Hirsch and R.M. Fye, *Phys. Rev. Lett.* 56, 2521 (1986).
 - [5] M. Jarrell, *Phys. Rev. Lett.* 69, 168 (1992); A. Georges and W. Krauth, *Phys. Rev. Lett.* 69, 1240 (1992).
 - [6] A.M. Oleś, *Phys. Rev. B* 28, 327 (1983).
 - [7] J.E. Hirsch, *Phys. Rev. B* 28, 4059 (1983); erratum, *Phys. Rev. B* 29, 4159 (1984).
 - [8] K. Held and D. Vollhardt, *Eur. Phys. J. B* 5, 473 (1998).
 - [9] Y. Motome and M. Imada, *J. Phys. Soc. Jpn.* 66, 1872 (1997); *ibid* 67, 3199 (1998).
 - [10] J.E. Han, cond-mat/0401104 (2004).
 - [11] hsigni is larger in Motome-Imada's method at half-filling ($n = 2$) due to the electron-hole symmetry, but it becomes smaller for $U = U^0 + 2J$.
 - [12] M. Jarrell, Th. Maier, C. Huscroft, and S. Moukouri, *Phys. Rev. B* 64, 195130 (2001).
 - [13] E. Müller-Hartmann, *Z. Phys. B* 74, 507 (1989); V. Zlatić and B. Horvatic, *Solid State Commun.* 75, 263 (1990).
 - [14] M. Capone, M. Fabrizio, C. Castellani, and E. Tosatti, *Science* 296, 2364 (2002); cond-mat/0401090 (2004).
 - [15] T. Morimoto and K. Kubo, *Phys. Rev. B* 58, 567 (1998).