

Explicit Gauge Fixing for Degenerate Multiplets: A Generic Setup for Topological Orders

Yasuhiro Hatsugai *

*Department of Applied Physics, Univ. of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, JAPAN*

(Received)

We supply basic tools for the study of the topological order of a multiplet which is an eigenspace of a finite-dimensional normal operator with continuous parameters. We allow intrinsic degeneracies within the multiplet where a well-known standard procedure does not work. As an important example, we give novel expressions for a spin Hall conductance for unitary superconductors with equal spin pairing. Generic topological orders will be treated in this unified manner particularly with nontrivial topological degeneracies.

KEYWORDS: Chern numbers, degeneracy, topological orders

It has been gradually clarified that many physically important phenomena have origins in their topological orders.^{1,2} Some of them include (fractional and integer) quantum Hall effects,^{3–12} Haldane spin chains,¹³ solitons in polyacetylenes,¹⁴ anisotropic superconductors and superfluids,^{15–20} chirality order in an itinerant magnetism,²¹ spin transport (spintronics) as a realistic application of Thouless pumping,^{22,23} polarizations in insulators,²⁴ and the exotic electronic states of graphite.²⁵ Strong correlations between electrons cause exotic mean field states^{26,27} and effective quasiparticles such as composite fermions²⁸ which can also be discussed in terms of the topological orders. In many cases, the topological order itself is hidden in bulk systems but exhibits apparent physical consequences at the boundaries of the systems, such as in edge states of the quantum Hall effects,^{6,29,30} local moments near impurities in the Haldane spin chains (Kennedy triplets),³¹ vortices and zero-bias conductance peaks in anisotropic superconductivities and boundary local moments in carbon nanotubes.²⁵

In many cases, nontrivial topological orders appear by restricting their physical space in a manner in which a type of gauge structure naturally emerges.^{32–34} To characterize the quantum state of a specific system, one must explicitly determine gauge invariant quantities for the physical states. The (first) Chern number^{8,35,36} is such a candidate and it has been used for several characterizations of topologically nontrivial states.^{7,30,37,38}

In this paper, we present a generic setup for the discussion of the topological order explicitly, particularly focusing on gauge fixing. A standard procedure for fixing gauge was reported by Kohmoto.⁸ This is well known today. However, the procedure does not work when degeneracy exists. If degeneracy is accidental, that is, it exists at certain special parameter values, it is negligible. However, in several interesting situations such as in unitary superconductors, degeneracy is due to an intrinsic symmetry, that is, the standard procedure cannot be applied for any values of the parameters (see below). In such cases, the present generic gauge fixing procedure is

essentially important, particularly for numerical calculations of Chern numbers. We extend the standard procedure to general situations which allow intrinsic degeneracies of eigenstates. A typical situation where our method is crucial is the calculation of spin Hall conductances for numerically obtained BCS Hamiltonians, where the order parameters are given numerically by minimizing the mean field free energy. Then quasiparticle states are obtained by diagonalizing a Bogoliubov-de Gennes equation. When the order is unitary, it has an intrinsic degeneracy³⁹ which prevents direct applications of the standard procedure to the calculation of the spin Hall conductance. Also, when a physical ground state has a nontrivial topological order and it lives on a genus $g(>0)$ Riemann surface, a fundamental topological degeneracy can occur with degeneracies q^g with some integer q .² A typical situation is the fractional quantum Hall effect with the filling factor $\nu = 1/q$.⁴⁰ In such a degenerate case, the present extension is indispensable. Further generic expressions in the present paper can be applicable to a wide range of physically interesting situations.

Multiplet and Unitary Equivalence: Let us consider taking a normal operator \mathbf{L} , ($\mathbf{L}^\dagger \mathbf{L} = \mathbf{L} \mathbf{L}^\dagger$), in an N -dimensional ($N < \infty$) linear space. This implies that \mathbf{L} is diagonalizable by a unitary matrix, $\mathbf{U} = (\psi_1, \dots, \psi_N)$, as $\mathbf{L}\mathbf{U} = \mathbf{U}\mathbf{\mathcal{E}}$, $\mathbf{\mathcal{E}} = \text{diag}(\epsilon_1, \dots, \epsilon_N)$. Note that normal operators include hermite, skew-hermite, unitary and skew-unitary operators. (Correspondingly, the eigenvalues ϵ_i 's are real, pure imaginary, and on the unit circle on the complex plane). Also, we assume that the operator \mathbf{L} is labeled by a set of continuous parameters as $\mathbf{L}(x)$, $x = (x_1, \dots, x_{d=\dim V})$, where V is a d -dimensional parameter space.⁴¹ Various physical realizations of the operator \mathbf{L} are (i) momentum-dependent Hamiltonians in the quantum Hall effect^{7,8} and an anisotropic superconductivity,^{17,42} (ii) parameter-dependent Hamiltonians $\mathbf{L}(x) = \mathbf{H}(x)$ in the discussion of Thouless pumping²² and the Berry phase,^{33,34} and (iii) a time evolution operator, $\mathbf{L}(x) = T \exp(-(i/\hbar) \int dt H(t))$.

Now construct an M -dimensional multiplet (a linear space) $W(x)$ with the parameters which we considered

*email:hatsugai@pothos.t.u-tokyo.ac.jp

($\dim W = M \leq N$). Take M linearly independent orthonormalized bases ψ_i , $i = 1, \dots, M$ for $W(x)$ (M column vectors of dimension N) and form an $N \times M$ matrix (the basis of the multiplet W) as

$$\Psi(x) = (\psi_1, \psi_2, \dots, \psi_M).$$

Supplementing the basis Ψ_C of the orthogonal multiplet W_C ($W \oplus W_C = \mathbb{R}^N$), they form an orthonormalized complete basis for \mathbb{R}^N as $(\Psi, \Psi_C) = \mathcal{U}$.

Here, we must be cautious. We require the multiplet $W(x)$ to be uniquely specified by the parameter x . However, this does not necessarily mean that the basis $\Psi(x)$ is uniquely specified by x , which gives us the freedom to change the basis. This leads to ambiguity in specifying the basis of $W(x)$. That is, we may take a different basis, $\Psi'(x) = \Psi(x)\omega(x)$, where ω is an M -dimensional unitary matrix. The ambiguity of the gauge freedom ω can be clarified, for example, when one tries to construct the multiplet from the ψ_i 's obtained numerically. This provides a gauge freedom for a connection we shall define. This was first observed by Wilczek-Zee in a study on non-Abelian Berry phases.³⁴

To specify the multiplet $W(x)$ uniquely using x , the multiplet should include all of the degenerate eigen spaces as

$$\epsilon_i(x) \neq \epsilon_j(x),$$

$$i \in I^{in} = \{1, \dots, M\}, j \in I^{out} = \{M+1, \dots, N\}. \quad (*)$$

We describe this condition the existence of a *generic energy gap*. The degeneracy within the multiplet is allowed, $\epsilon_i = \epsilon_j$, $i, j \in I^{in}$, which exists, for example, in unitary superconductivity⁴² and spin degenerate cases. The degeneracy between the multiplet W we considered and the supplementary W_C is not allowed.

The parameter space and the multiplet are specified by the concrete topological orders we shall study. Let us list some examples of V s with corresponding multiplets W s: (1) V : the Brillouin zone, (1-i) W : a collection of Landau level wave functions in the quantum Hall effects,⁷ (1-ii) W : quasi-particle states in anisotropic superconductors with and without equal spin pairing,¹⁷ and (2) V : a set of external parameters in the study of Thouless pumping²² and the Berry phase,^{33,34} W : generically degenerate ground states, to be specific, for example, V : (2-i) a collection of fluxes passing through the systems, Aharonov-Bohm fluxes and the strength of vortices in the type II superconductivity and (2-ii) parameters specifying the axes of spins in magnetic systems.

Connection and First Chern Number: Define a non-Abelian connection one-form \mathcal{A} which is an $M \times M$ matrix as $\mathcal{A} = \Psi^\dagger d\Psi$. Correspondingly, $\mathcal{A}' = \Psi' d\Psi' = \omega^{-1} \mathcal{A} \omega + \omega^{-1} d\omega$. This is a *gauge transformation* in our problem. Also, define a field strength two-form $\mathcal{F} = d\mathcal{A} + \mathcal{A}^2$ which is transformed by gauge transformation as $\mathcal{F}' = \omega^{-1} \mathcal{F} \omega$. Then $\text{Tr } \mathcal{F}$ is gauge invariant and an integral of $\text{Tr } \mathcal{F}$ over the two-dimensional orientable compact surface S ($\partial S = 0$) in V gives the first Chern number^{35,36} $C_S = \frac{1}{2\pi i} \int_S \text{Tr } \mathcal{F} = \frac{1}{2\pi i} \int_S \text{Tr } d\mathcal{A}$. For example, consider the three-dimensional Brillouin zone $V = T^3(k_x, k_y, k_z)$ as the full parameter space and

the two-dimensional Brillouin zone, $S = T_{xy}^2(k_z)$, with the fixed third momentum k_z for the surface S .⁴³⁻⁴⁵

A global connection over the full surface S is not allowed to exist in a system with a nontrivial topological order. Then let us divide the integral region S into several patches S_R ($S = \cup S_R$, $R = 0, 1, 2, \dots$), where the connection \mathcal{A}_R is locally defined within S_R as $\mathcal{A}(x) = \mathcal{A}_R(x)$.^{8,35} Furthermore, we assume each S_R , $R = 1, 2, \dots$, does not share any boundaries with $\partial S_0 = -\sum_{R \geq 1} \partial S_R$. When the connection \mathcal{A}_R is related to \mathcal{A}_0 by the gauge transformation ω_{0R} as $\mathcal{A}_R = \omega_{0R}^{-1} \mathcal{A}_0 \omega_{0R} + \omega_{0R}^{-1} d\omega_{0R}$, the Chern number C_S is written using the Stokes theorem as $C_S = \frac{1}{2\pi} \sum_{R \geq 1} \int_{\partial S_R} \text{Im Tr } \omega_{0R}^{-1} d\omega_{0R}$.

Explicit Gauge Fixing: Topological invariants are usually given by gauge-dependent quantities. To evaluate the expression, one must fix the gauge. Without fixing it, we cannot have any well-defined derivative. Now let us explicitly fix the gauge for the multiplet. Although the basis Ψ has a gauge freedom, a projection operator into the multiplet $P = \Psi \Psi^\dagger$ is a gauge invariant. Define an unnormalized basis, Ψ_Φ^U , from a generic basis, Φ (an $N \times M$ matrix), as $\Psi_\Phi^U = P\Phi = \Psi \eta_\Phi$ ($\eta_\Phi = \Psi^\dagger \Phi$). The overlap matrix of the basis $O_\Phi = \Psi_\Phi^{U\dagger} \Psi_\Phi^U$ is generically semipositive definite. Then, only if the determinant of the matrix O_Φ is nonzero, we can define a normalized wavefunction, $\bar{\Psi}_\Phi = \Psi_\Phi^U O_\Phi^{-1}$, where $o_\Phi \equiv U_\Phi^{1/2} \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_M}) U_\Phi$ with $O_\Phi = U_\Phi \text{diag}(\lambda_1, \dots, \lambda_M) U_\Phi$. This o_Φ is hermite and positive definite. Now we define the connection \mathcal{A}_Φ with the gauge fixing by Φ as $\mathcal{A}_\Phi = \bar{\Psi}_\Phi^\dagger d\bar{\Psi}_\Phi$. This is well defined unless $\det O_\Phi = 0$.^{8,37}

Define regions S_R^Φ , $R = 1, 2, \dots$, as (infinitesimally) small neighborhoods of zeros x_R^Φ 's of $\det O_\Phi(x)$ and S_0^Φ as a rest of S as

$$S = \bigcup_{R \geq 0} S_R^\Phi, \quad \det O_\Phi(x) \begin{cases} \neq 0 & \forall x \in S_0^\Phi \\ = 0 & \text{at } \exists x_R \in S_R^\Phi, R = 1, 2, \dots \end{cases}$$

We use this gauge for the region S_0^Φ , and for the region S_R^Φ , we use a different gauge by $\tilde{\Phi}$, with $\det O_{\tilde{\Phi}} \neq 0$ everywhere in S_R^Φ . The transformation matrix between $\bar{\Psi}_{\tilde{\Phi}}$ and $\bar{\Psi}_\Phi$ is obtained as $\omega = o_\Phi \eta_\Phi^{-1} \eta_{\tilde{\Phi}} o_{\tilde{\Phi}}^{-1}$. Since o_Φ and $o_{\tilde{\Phi}}$ are strictly positive definite at the boundaries ∂S_0^Φ , we have $\text{Im Tr } \log \omega = -\text{Im Tr } \log \tilde{\Phi}^\dagger P \Phi$. Finally, we obtain an expression for the first Chern number with explicit gauge fixing as

$$C_S = -N_\Omega^T(S) = -\sum_{R \geq 1} n_\Omega^R(S_R^\Phi)$$

$$n_\Omega^R(S_R^\Phi) = \frac{1}{2\pi} \oint_{\partial S_R^\Phi} d'\Omega, \quad \Omega = \Omega(\tilde{\Phi}, \Phi) = \text{Arg det } \tilde{\Phi}^\dagger P \Phi,$$

where $N_\Omega^T(S)$ is the total number of signed vortices with the vorticity $n_\Omega^R(S_R^\Phi)$ inside the region S_R , $R \geq 1$ ($\partial S_0^\Phi = -\cup_{R \geq 1} \partial S_R^\Phi$).⁴⁶ Since $\Omega = \text{Arg det } \eta_{\tilde{\Phi}}^\dagger \det \eta_\Phi$, all the vortices of $\Omega(\tilde{\Phi}, \Phi)$ are given by zeros of $\det \eta_{\tilde{\Phi}}$, ($x^{\tilde{\Phi}}$) and $\det \eta_\Phi$, (x^Φ). The Chern number is obtained by summing up the vorticity only at x^Φ . This form of Chern number is not found in literature.

Since we assume that the two-dimensional surface S is compact and Ω is regular except at x_1^Φ, \dots and $x_1^{\tilde{\Phi}}, \dots$, a union of curves $(\cup_R S_R^\Phi) \cup (\cup_R S_R^{\tilde{\Phi}})$ is contractible to a point within a region where Ω is well defined. This implies that $N_{\Omega(\tilde{\Psi}, \Psi)}^T(S) = N_{\Omega(\Psi, \tilde{\Psi})}^T(S)$. That is, the vector field Ω depends on the gauge (choice of Φ and $\tilde{\Phi}$) but the total vorticity $N_\Omega^T(S)$ is a gauge invariant of the multiplet W . None of the vortices has any direct physical meaning. Only the total number of vortices $N_\Omega^T(S)$ has a physical significance.

The projection \mathbf{P} is essential for carrying out the present gauge fixing procedure. It has also an integral representation, $\mathbf{P} = \frac{1}{2\pi i} \oint_\Gamma G_T(z)$, where $\mathbf{G}_T = (\mathbf{I}_N - \mathbf{L})^{-1}$ and the closed curve Γ encloses all of the eigenvalues $\epsilon_1, \dots, \epsilon_M$ inside, but not those $\epsilon_{M+1}, \dots, \epsilon_N$ on the complex plane.⁴⁷ From this form of projection, the stability condition of the generic gap (*) for obtaining a well-defined multiplet is clear. The first Chern number has an apparently gauge-independent form given by $\text{Tr } \mathcal{F} = -\text{Tr } d\mathbf{P} \mathbf{P} d\mathbf{P}$ as well.⁴⁸ Also, the Chern number for the multiplet is expressed as

$$C_S = \frac{1}{2\pi i} \sum_{k \in I^n} \int_S \langle d\mathbf{L}^\dagger \{ \mathbf{G}_C(\epsilon_k) \}^\dagger \{ \mathbf{G}_C(\epsilon_k) \} d\mathbf{L} \rangle_{\psi_k}, n$$

where $\mathbf{G}_C = (\mathbf{I} - \mathbf{P})\mathbf{G}_T$. This is equivalent to a Kubo formula in the case of the quantum Hall effect.^{7,48} This formula is particularly important since mathematical objects such as Chern numbers have a direct relation with a physical quantity such as a Hall conductance. Surprisingly, this is observable in a bulk system.

Sum Rules: Assume the multiplet W is a direct sum of orthogonal multiplets W_1 and W_2 as $W = W_1 \oplus W_2$, which is expressed by bases Ψ_1 and Ψ_2 (orthonormalized in each multiplet) as $\Psi = (\Psi_1, \Psi_2)$, where $\Psi_1^\dagger \Psi_1 = \mathbf{I}_{M_1}$, $\Psi_2^\dagger \Psi_2 = \mathbf{I}_{M_2}$, $\Psi_1^\dagger \Psi_2 = \mathbf{O}_{M_1 M_2}$, and $\Psi_2^\dagger \Psi_1 = \mathbf{O}_{M_2 M_1}$.

The connection is given as $\mathcal{A} = \begin{pmatrix} \Psi_1^\dagger \\ \Psi_2^\dagger \end{pmatrix} (d\Psi_1, d\Psi_2)$.

Thus, a trace of the connection is additive as $\text{Tr } \mathcal{A} = \text{Tr } \mathcal{A}_1 + \text{Tr } \mathcal{A}_2$, where $\mathcal{A}_1 = \Psi_1^\dagger d\Psi_1$ and $\mathcal{A}_2 = \Psi_2^\dagger d\Psi_2$. From this simple observation in the connection level, a *sum rule* for Chern numbers is as follows: $C_S(W_1 \oplus W_2) = C_S(W_1) + C_S(W_2)$. The sum rule in the field strength level was previously discussed.⁴⁹ A simple consequence of the present sum rule is the total sum rule, that is, the Chern number of the total multiplet W_T always vanishes,⁴² $\sum_i C_S(W_i) = C_S(\oplus_i W_i) = 0$, since $\mathbf{P}_{\oplus_i W_i} = \mathbf{I}_N$.

One-Dimensional Example (dim $W = 1$): When the multiplet is one-dimensional, such as $\Psi = \psi$ and $(\mathbf{P})_{ij} = \psi_i \psi_j^*$, we have $\tilde{\Phi}^\dagger \mathbf{P} \tilde{\Phi} = (\mathbf{P})_{N1} = \psi_N^* \psi_1$ by taking Φ and $\tilde{\Phi}$ as ${}^t\Phi = (1, 0, \dots, 0)$ and ${}^t\tilde{\Phi} = (0, \dots, 0, 1)$. Then the Chern number is given as

$$C_S = -\frac{1}{2\pi} \oint_{\partial S_\Phi^\Phi} d\text{Im} \log(\psi_1/\psi_N), \quad S_0^\Phi = S \setminus \bigcup_{R \geq 1} S_{R \geq 1}^\Phi,$$

where S_R^Φ includes a single zero of $\det \mathcal{O}_\Phi = |\psi_1|^2$. This is a well-known classic expression.⁸

Multiplet of Several Landau Levels: When one consid-

ers two-dimensional electrons on a lattice with the flux ϕ per plaquette, one-particle states are given by q bands when $\phi = p/q$ with the mutually prime p and q . Furthermore, the spectrum is given by the famous Hofstadter's butterfly. When the fermi energy E_F is in the j -th energy gap from below, the Hall conductance σ is given by the sum of the Chern numbers of the j bands.⁷ In this case, take a multiplet from a filled fermi sea ($W = \text{FS}$) and construct the basis of the multiplet from the j Bloch states $\psi_j(\mathbf{k})$ below E_F as $\Psi = (\psi_1, \dots, \psi_j)$, $M = j$, then the Chern number $C_{T^2}(\text{FS})$ naturally gives the Hall conductance σ_{xy} which is the sum of the Chern numbers of the filled bands.^{7,37,38,49}

Dirac Monopole: When the dimension of the total Hilbert space N is 2, only the nontrivial multiplet is one-dimensional $M = 1$. Then take an hermite Hamiltonian $\mathbf{H}(x) = \mathbf{R}(x) \cdot \boldsymbol{\sigma}$ for the normal operator \mathbf{L} where $\boldsymbol{\sigma}$'s are Pauli matrices and $\mathbf{R}(x)$ is a real three-dimensional vector ((R, θ, ϕ) is a polar coordinate of \mathbf{R}). As an example, consider the multiplet Ψ_- with the energy $-R$ as³³ ${}^t\Psi_-(x) = (-\sin \frac{\theta(x)}{2}, e^{i\phi(x)} \cos \frac{\theta(x)}{2})$. The projection is given as $\mathbf{P}_- = \Psi_- \Psi_-^\dagger = \begin{pmatrix} \sin^2 \frac{\theta}{2} & -e^{-i\phi} \sin \frac{\theta}{2} \cos \frac{\theta}{2} \\ -e^{i\phi} \sin \frac{\theta}{2} \cos \frac{\theta}{2} & \cos^2 \frac{\theta}{2} \end{pmatrix}$. Using a gauge by Φ and $\tilde{\Phi}$ as ${}^t\Phi = (\cos \frac{\chi}{2}, e^{i\xi} \sin \frac{\chi}{2})$ and ${}^t\tilde{\Phi} = (\cos \frac{\tilde{\chi}}{2}, e^{i\tilde{\xi}} \sin \frac{\tilde{\chi}}{2})$, $\det \mathcal{O}_\Phi = 0$ and $\det \mathcal{O}_{\tilde{\Phi}} = 0$ give $(\theta(x), \phi(x)) = (\chi, \xi)$ and $(\tilde{\chi}, \tilde{\xi})$, respectively. This clearly shows that the positions of the vortices defined by the vector field $\Omega = \text{Arg}(-\sin \frac{\theta}{2} \cos \frac{\chi}{2} + e^{+i(\phi-\tilde{\xi})} \cos \frac{\theta}{2} \sin \frac{\tilde{\chi}}{2})(-\sin \frac{\theta}{2} \cos \frac{\chi}{2} + e^{-i(\phi-\xi)} \cos \frac{\theta}{2} \sin \frac{\tilde{\chi}}{2})$ are gauge-dependent and do not have any direct physical meaning. One can chose the positions of the vortices as one wishes.

Unitary Superconductors: Let us first consider the simplest case, that is, the unit cell includes only one site. Then the Bogoliuvov-de Gennes equation for generic superconductivity is given in a momentum space by a 4×4 secular equation, $\mathbf{H}\psi = E\psi$, $\mathbf{H} = \begin{pmatrix} \epsilon \mathbf{I}_2 & \Delta \\ \Delta^\dagger & -\epsilon \mathbf{I}_2 \end{pmatrix}$. As for the unitary order, the order parameter matrix Δ is written as $\Delta \equiv |\Delta| \Delta_0$, $|\Delta| \geq 0$, $\Delta \Delta^\dagger = |\Delta|^2 \mathbf{I}_2$, where Δ_0 is a 2×2 unitary matrix. Then the eigenstates (quasiparticle) are doubly degenerate as ${}^t\psi_-(\mathbf{w}) = (-\sin \frac{\theta}{2} \mathbf{w}, \cos \frac{\theta}{2} \Delta_0^\dagger \mathbf{w})$, for example, for the $E = -R$ state where \mathbf{w} is a normalized arbitrary two-component vector, $\mathbf{w}^\dagger \mathbf{w} = 1$, $R = \sqrt{|\Delta|^2 + \epsilon^2}$, $\epsilon = R \cos \theta$ and $|\Delta| = R \sin \theta$. Now let us construct a multiplet for the degenerate $E = -R$ quasiparticle bands as $\Psi_- = (\psi_-(\mathbf{w}_1), \psi_-(\mathbf{w}_2))$, where \mathbf{w}_1 and \mathbf{w}_2 form an arbitrary two-dimensional orthonormalized complete set: $\mathbf{w}_i^\dagger \mathbf{w}_j = \delta_{ij}$, $\mathbf{w}_1 \mathbf{w}_1^\dagger + \mathbf{w}_2 \mathbf{w}_2^\dagger = \mathbf{I}_2$. Then the projections \mathbf{P}_- are given in a gauge invariant form as $\mathbf{P}_- = \begin{pmatrix} \mathbf{I}_2 \sin^2 \frac{\theta}{2} & -\Delta_0 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \\ -\Delta_0^\dagger \sin \frac{\theta}{2} \cos \frac{\theta}{2} & \mathbf{I}_2 \cos^2 \frac{\theta}{2} \end{pmatrix}$. Now let us fix the gauge by choosing ${}^t\Phi = {}^t(\mathbf{0}_2, \mathbf{I}_2)$ and ${}^t\tilde{\Phi} = {}^t(\mathbf{I}_2, \mathbf{0}_2)$. Then we have $\det \mathcal{O}_{-, \Phi} = \cos^4 \frac{\theta}{2}$, $\Omega = \text{Arg} \det \Delta_0 = \text{Arg} \det \Delta$. Since the overlap determinant $\det \mathcal{O}_{-, \Phi} = 0$ vanishes at $\theta = \pi$ for the multiplet W_- , the Chern num-

ber is given by

$$C_S^- = -\frac{1}{2\pi} \sum_p \oint_p d' \text{Arg det } \Delta,$$

where p 's are points on the surface S which are specified by $\theta = \pi$, that is, $|\Delta|(p) = 0$ and $\epsilon(p) = -R(p) = -E(p)$. This is a novel expression for the generic spin Hall conductance for the unitary superconductors with equal spin pairing. In the previous work,⁴² the Chern number was given as the sum of two integers using an eigenvalue equation for the unitary matrix Δ_0 . Here, we give a direct expression only using the order parameter matrix Δ . Furthermore, if one parameterizes the unitary 2×2 matrix Δ_0 as $\Delta_0 = e^{i\Theta} e^{i\hat{n} \cdot \vec{\sigma}}$, $|\hat{n}| = 1$, we have $C_S^- = -\frac{1}{\pi} \sum_p \oint_p d\Theta$. The present method is crucially important and efficient when the order parameter is given by numerically solving a BCS self-consistent equation with a large unit cell. Even in this generic situation, to evaluate the spin Hall conductance, we must determine arbitrary orthonormalized (degenerate) eigenstates by diagonalizing the mean field Hamiltonian (Bogoliubov-de Gennes equation).

Finally, we mention the higher order Chern numbers $C_n(S^n) = N_n \int_{S^n} \text{Tr } \mathcal{F}^n$, $n = 2, 3, \dots$, where N_n is a normalization constant and $S^n = S \times \dots \times S$, (n -times). In principle, they can also explicitly be evaluated using the present gauge fixing procedure as a sum of integrals over the $(2n-1)$ -dimensional spheres S_R^Φ enclosing $(2n-2)$ -dimensional regions P_R^Φ which are defined by the zero of $\det \mathcal{O}_\Phi$ in S^n . They should also help in the characterization of the topological order in complex situations.

We thank M. Kohmoto for useful discussions. Part of the work by Y.H. was supported by a Grant-in-Aid from the Japanese Ministry of Education, Culture, Sport, Science and Technology, and the JFE 21st Century Foundation.

- 1) A. Shapere and F. Wilczek, eds.: *Geometric Phases in Physics* (World Scientific, Singapore, 1989).
- 2) X. G. Wen: Phys. Rev. B **40** (1989) 7387.
- 3) K. V. Klitzing, G. Dorda and M. Pepper: Phys. Rev. Lett. **45** (1980) 494.
- 4) T. Ando: J. Phys. Soc. Jpn. **37** (1974) 622.
- 5) H. Aoki and T. Ando, Solid State Comm. **38** (1981) 1079.
- 6) R. B. Laughlin: Phys. Rev. B **23** (1981) 5632.
- 7) D. J. Thouless, M. Kohmoto, P. Nightingale and M. den Nijs: Phys. Rev. Lett. **49** (1982) 405.
- 8) M. Kohmoto: Ann. Phys. (N. Y.) **160** (1985) 355.
- 9) R. B. Laughlin: Phys. Rev. Lett. **50** (1983) 1395.
- 10) Q. Niu, D. J. Thouless and Y. Wu: Phys. Rev. B **31** (1985) 3372.
- 11) S. Girvin, A. H. MacDonald and P. M. Platzman: Phys. Rev.

Lett. **54** (1985) 581.

- 12) P. Stréda: J. Phys. C **15** (1982) L717.
- 13) F. D. M. Haldane: Phys. Rev. Lett. **50** (1983) 1153.
- 14) W. P. Su, J. R. Schrieffer and A. J. Heeger: Phys. Rev. Lett. **42** (1979) 1698.
- 15) T. Senthil, J. B. Marston and M. Fisher: Phys. Rev. B **60** (1999) 4245.
- 16) Y. Morita and Y. Hatsugai: Phys. Rev. B **62** (2000) 99.
- 17) Y. Hatsugai and S. Ryu: Phys. Rev. B **65** (2002) 212510.
- 18) G. E. Volovik: Pis'ma ZhETF, JETP Lett. **66** (1997) 493.
- 19) G. E. Volovik: *Exotic Properties of Superfluid ^3He* (World Scientific, Singapore, 1992).
- 20) N. Read and D. Green: Phys. Rev. B **61** (2000) 10267.
- 21) Y. Taguchi, Y. Oohara, H. Yoshizawa, N. Nagaosa and Y. Tokura: Science **291** (2001) 2573.
- 22) D. J. Thouless: Phys. Rev. B **27** (1983) 6083.
- 23) S. Murakami, N. Nagaosa and S.-C. Zhang: Science **5** (2003) 1348.
- 24) R. D. King-Smith and D. Vanderbilt: Phys. Rev. B **47** (1993) 1651.
- 25) S. Ryu and Y. Hatsugai: Phys. Rev. Lett. **89** (2002) 077002.
- 26) X. G. Wen, F. Wilczek and A. Zee: Phys. Rev. B **39** (1989) 11413.
- 27) I. Affleck and J. B. Marston: Phys. Rev. B **37** (1988) 1988.
- 28) J. Jain: Phys. Rev. Lett. **63** (1989) 199.
- 29) B. I. Halperin: Phys. Rev. B **25**, (1982) 2185.
- 30) Y. Hatsugai: Phys. Rev. B **48** (1993) 11851.
- 31) T. Kennedy: J. Phys. Condens. Matter **2** (1990) 5737.
- 32) P. Dirac: *Lectures on Quantum Mechanics* (Yeshiva Univ., New York, 1964).
- 33) M. V. Berry: Proc. R. Soc. **A392** (1984) 45.
- 34) F. Wilczek and A. Zee: Phys. Rev. Lett. **141** (1984) 2111.
- 35) T. Eguchi, P. B. Gilkey and A. J. Hanson: Phys. Rep. **66** (1980) 213.
- 36) B. Zumino: *Current Algebra and Anomaly* (World Scientific, Singapore, 1991), p.362.
- 37) Y. Hatsugai: Phys. Rev. Lett. **71** (1993) 3697.
- 38) Y. Hatsugai: J. Phys. C, Condens. Matter **9** (1997) 2507.
- 39) M. Sigrist and K. Ueda: Rev. Mod. Phys. **63** (1991) 239.
- 40) F. D. M. Haldane and E. H. Rezayi: Phys. Rev. B **31** (1985) 2529.
- 41) We require $\dim V = d \geq 2$ for the following discussions. However, one can freely supplement additional parameters. Then the case $\dim V = 1$ is also within the analysis.
- 42) Y. Hatsugai, S. Ryu and M. Kohmoto: Phys. Rev. B **70** (2004) 054502.
- 43) B. I. Halperin: Jpn. J. Appl. Phys. Suppl. **26** (1987) 1913.
- 44) M. Kohmoto, B. I. Halperin and Y. S. Wu: Phys. Rev. B **45** (1992) 13488.
- 45) The surface S can be a genus g Riemann surface Σ_g . The simplest and the most common case is the torus $T^2 = \Sigma_1$ which appears as a two-dimensional Brillouin zone and a couple of fluxes in the multiframe AB effects.
- 46) We have to note that integral theorems cannot be applied here since $\log \det \omega$ and $\text{Arg det } \tilde{\Phi}^\dagger P \Phi$ are not global single valued functions at the boundaries ∂S_0^Φ . We use d' to indicate it.
- 47) T. Kato: Prog. Theor. Phys. **4** (1949) 514.
- 48) J. E. Avron, R. Seiler and B. Simon: Phys. Rev. Lett. **51** (1983) 51.
- 49) J. E. Avron and R. Seiler: Phys. Rev. Lett. **54**: (1985) 259.