

REALISTIC OPERATION OF AN ELECTRON ENTANGLER : A DENSITY MATRIX APPROACH

OLIVIER SAURET¹, DENIS FEINBERG¹, THIERRY MARTIN²

¹ LEPES, CNRS, BP 166, 38042 Grenoble, FRANCE

² CPT, CNRS, Univ. Mediterranee, Case 907, 13288 Marseille, FRANCE

The detailed operation of an electron spin entangler is studied, using density matrix equations. The device is made of a superconductor, two quantum dots and two normal leads. The treatment takes into account coherent tunneling in a non-perturbative way, and analyzes the various parasitic effects, in addition to the main process (crossed Andreev reflection) : those include singlet pairs passing through a single dot, or cotunneling between dots through the superconductor. The optimum operation of the device is characterized.

Producing entangled electron pairs is a challenge for fundamental experiments (analogous to those performed with photons¹), as well as controlling quantum information² in solid state devices³. It has been proposed that a superconductor, connected to energy filters, can serve as a source of spin singlets^{4,5}. Here we address the operation of the "S-DD entangler" made of a superconductor (S) connected in parallel to a double quantum dot (DD) where Coulomb blockade prevents two electrons to pass through a single dot. The dots are small enough so that a single electron state is involved in each dot. The "crossed Andreev" (CA)^{6,7} process is thus favoured, where a spin singlet is emitted, shared by the DD. The feasibility of such a device crucially depends on the control of "parasitic" processes spoiling entanglement, mainly of two kinds : first, singlet pairs can pass through a single dot, either through a double occupation state (direct Andreev DA), or one by one (Fig. 1). Second, an electron can pass from one dot to the other by elastic cotunneling^{6,8} (CT) through S. The first, but not the second, was considered in Ref. 5. In addition, all processes are mixed together, making a consistent treatment difficult. Such a study is indeed possible by using the density matrix equations, which generalize the usual master equations to the inclusion of coherent processes. Those correspond to both Andreev transitions or to cotunneling. They are made of two virtual transitions, with a quasiparticle created in S then destroyed. On the contrary, single electron transitions between dots and the leads L,R are incoherent. The complete quantum master (QM) equations for the subsystem made of S and the two dots can be derived⁹ for instance following Ref. 10. Here the discussion is based on the analysis of the dot populations and averaged current flow. Further results are devoted to shot noise correlations and Bell inequalities¹¹. Notice that the S-DD entangler was recently studied¹² in series with a splitter detecting entanglement¹³. On the other hand, QM equations were employed¹⁴ for a different principle of entangler using another dot instead of a superconductor¹⁵. Also, QM equations were at the basis of the analysis of a device permitting teleportation of the electron spin in a dot array¹⁶.

In the ideal operation of the S-DD entangler, the Coulomb blockade in each dot is strong enough so as to rule out double occupancy. Starting from an empty state 00, CA reaction allows transitions to the singlet state, shared between the two dots, 11_s with a rate $\Gamma_A T$ (Γ_A is the geometrical factor^{6,7}). For a resonant CA process, the dot energy levels satisfy $\epsilon = E_1 + E_2 = 0$. The two electrons are evacuated from the dots into the reservoirs (with chemical potentials $\mu_{L,R} < E_1, E_2$) and the transitions to states 01, 10 occur with rates Γ_i ($i = L, R$).

If the Coulomb charging energy is not so strong, a coherent transition from 00 to a doubly occupied dot states 20 or 02 can occur via a direct Andreev (DA) process, which has a rate Γ_i which is larger than for the CA process (Fig. 1). Electrons can subsequently be detected into this reservoir, with rates Γ_i^0 . This conduction channel implies dot energies $E_i + U_i$ associated with double occupancy. One may also start from an initial state 10 or 01. DA can then

proceed through the empty dot, but the charging energy of states 21 or 12 equals $E_1 + E_2 + U_i$. Detection in the reservoir can either lead to 20 (02), or to singlet and triplet states 11_{st} . Another parasitic channel involves two electrons of a Cooper pair tunneling one by one towards the same reservoir (Fig. 1). This involves a singly occupied virtual state which costs an energy ϵ_S , the superconducting gap. Contrary to the DA process, the dot is emptied before the quasiparticle in S is annihilated. This process happens with a rate $\sim \Gamma_i = \Gamma_i^2 / \epsilon_S^2$. Last but not least, cotunneling (CT) allows a coherent transfer of an electron from one dot to the other via S. It couples states 01 and 10, but also 20 (02) and 11, 21 and 12. CT has a rate Γ_C which is reduced by a geometrical factor.




Figure 1: Full operation of the entangler, including the three Andreev channels and cotunneling. Real states are squared while virtual states are dashed squared. States with three electrons are omitted for clarity, spin is not represented. The Ω 's are the one-electron tunneling matrix elements. The crossed Andreev (CA) process is in the middle, the direct Andreev ones (DA) second from top or bottom, and the one-by-one processes are the top or bottom ones. Cotunneling (CT) connects the states vertically.

Let us exclude high energy states 22 with $N = 4$ electrons in the double dot (DD), keeping states 00, 0, 0 (with spin), 11 (singlet and triplets), 2 and 2 (2 means a local singlet). Transport of electron pairs through the DD is highly correlated. With CA alone, pairs pass one after the other through the DD. Including the parasitic processes, there is a strong mixing of processes and no simple perturbative calculation is possible. Yet, starting from the full one-electron Hamiltonian, one can derive a complete set of QM equations for the populations $p = \text{Tr}(\rho)$ and the "coherences" (diagonal and non-diagonal matrix elements of the density matrix ρ_{DD}). It takes the general form¹⁰:

$$\dot{\rho} = -i[H, \rho] + \sum_i \Gamma_i T_i(\rho) + \sum_j \Gamma_j T_j(\rho) \quad (1)$$

$$\dot{\rho} = -i(E - E) + i \sum_i \Gamma_i (T_i - T_i^\dagger) + \frac{\sum_j \Gamma_j}{2} (\rho + \rho^\dagger) \quad (2)$$

where the Γ 's are the coherent rates and the Γ 's the incoherent rates. One assumes in the derivation that virtual states with at most one quasiparticle in S are created. The obtained set of equations⁹ is valid up to second order in the matrix elements describing tunneling to the leads, and to any order in the coherent rates (which are of order two in the tunneling matrix elements between the superconductor and the dots).

In Ref. 5, a T-matrix calculation was performed, calculating separately the ideal (CA) current, and the DA and one-by-one parasitic currents. Here the optimum operation of the device can be settled on a finer basis, and a better understanding of the physics involved is obtained. First let us assume a symmetric device ($\Gamma_L = \Gamma_R$) and treat the processes separately, without cotunneling. Assuming $\frac{8}{A} T^2 \ll U^2$, the CA current in each lead is

$$I_L^{CA} = e \frac{8 \frac{A}{A} T^2}{8 \frac{A}{A} T^2 + U^2} \quad (3)$$

On the other hand, the DA current I_L^{DA} and the one-by-one current I_L^{obo} read

$$I_L^{DA} = e \frac{16T^2}{16T^2 + U^2}; \quad I_L^{obo} = e \frac{K T^2}{\frac{U}{S} + K T^2} \quad (4)$$

where K is a numerical constant. If $T \ll U/S$, one has

$$I_L^{DA} \approx e \frac{16T^2}{U^2}; \quad I_L^{obo} \approx 4e \frac{T^2}{\frac{U}{S}} \quad (5)$$

The general case can be treated, setting $U = E_1 + E_2$ and putting all processes together. As an illustration, an analytical formula can be given for the total current, up to first order in the parasitic processes

$$I_L = e \left[\Gamma_L + \Gamma_R + 4 \frac{T^2}{U^2} \left(\frac{1}{\Gamma_L} + \frac{1}{\Gamma_R} \right) \frac{c T^2}{E^2} + 2K \frac{A}{U} \left(\frac{T^2}{S} + \frac{1}{\Gamma_0} \right) \right] \quad (6)$$

$$8A \frac{1}{U^2} \left(\frac{T^2}{S} + \frac{1}{\Gamma_0} \right) + \frac{2}{U^2} \left(\frac{T^2}{S} + \frac{1}{\Gamma_0} \right) \left[\frac{1}{\Gamma_L} + \frac{1}{\Gamma_R} \right]$$

where $A = \frac{8}{2 + 8 \frac{A}{A} T^2 + U^2}$ and $\Gamma_0^{-1} = A + 1 + \Gamma_R = \Gamma_L + \Gamma_R$.




Figure 2: Populations of the DD states as a function of $U = \Gamma$ for $S = 9.5K$; $E_1 = E_2 = 0.5K$; $T = 0.1K$; $A = 0.2$. States b, c, d, e, f, g, h, k respectively correspond to charge states $(11_s), (10), (01), (20), (02), (21), (12)$ and (11_t) .

The optimization of the entangler requires that most pairs emitted by the superconductor leave separated into L and R. To fight DA and one-by-one processes, one may compare the currents, taken individually for each process, or alternatively adopt a dynamical argument: starting from state $|i\rangle = (00)$ at time $t = 0$, the probability p_i of singlet state $|i\rangle$ oscillates slowly (with frequency $A T$), but with a large amplitude. On the contrary, the probability of state $|j\rangle = (20)$ oscillates more rapidly (with frequency T), but with a small amplitude. The competition between the two processes crucially depends on the decay rate Γ . If it is small, CA is favoured, but if it is too large, DA process wins, state $|i\rangle$ has no time to form. A detailed

analysis gives the criterion $U; S \gg \max[T; =_A; =_A]$. A similar analysis can be made for the effect of cotunneling: once in the state (10) or (01), decay in L;R must be faster than the cotunneling frequency $_CT$, leading to the criterion $_CT \ll \max[E_1 - E_2;]$. This can be confirmed by a numerical solution for the probabilities of various states (Fig. 2).

In summary, density matrix (quantum master) equations can be derived from a microscopic Hamiltonian for a realistic entangler, and allow to integrate all processes in a coherent and non-perturbative way. A range of parameters for optimum operation is $_AT; _CT \ll _{L,R} \ll U; S$. More information can be obtained by further analysis of the current fluctuations (shot noise correlations)¹¹.

References

1. A. A. Spect, J. Dalibard, and G. Roger, Phys. Rev. Lett. 49, 1804 (1982); L. Mandel, Rev. Mod. Phys. 71, S274 (1999); A. Zeilinger, Rev. Mod. Phys. 71, S288 (1999).
2. D. Bouwmeester, A. Ekert, and A. Zeilinger, The Physics of Quantum Information (Springer-Verlag, Berlin, 2000); M. A. Chang and I. L. Nielsen, Quantum Computation and Quantum Information, Cambridge University Press, 2000).
3. D. Loss and D. P. DiVincenzo Phys. Rev. A 57, 120 (1998).
4. G. B. Lesovik, T. Martin, and G. Blatter, Eur. Phys. J. B 24, 287 (2001); N. M. Chitchev, G. Blatter, G. B. Lesovik and T. Martin, Phys. Rev. B 66, 161320 (2002).
5. P. Recher, E. V. Sukhorukov, and D. Loss, Phys. Rev. B 63, 165314 (2001).
6. G. Falci, D. Feinberg, and F. W. J. Hekking, Europhys. Lett. 54, 255 (2001).
7. M. S. Choi, C. Bruder, and D. Loss, Phys. Rev. B 62, 13569 (2000).
8. D. V. Averin and Yu. V. Nazarov, in Single Charge Tunneling, H. Grabert and M. H. Devoret eds. (Plenum, New York 1992). G. Bignon, M. Houzet, F. Pistolesi and F. Hekking, cond-mat/0310349 (2003).
9. O. Sauret, D. Feinberg and T. Martin, cond-mat/0402416 (Phys. Rev. B, to appear).
10. S. A. Gurvitz and Ya. S. Prager, Phys. Rev. B 53, 15932 (1996).
11. O. Sauret, T. Martin and D. Feinberg, in preparation.
12. P. Samuelsson, E. V. Sukhorukov and M. Buttiker cond-mat/0402368.
13. G. Burkard, D. Loss, and E. V. Sukhorukov Phys. Rev. B 61, R16303 (2000).
14. D. Saraga, D. Loss, Phys. Rev. Lett. 90, 166803 (2003).
15. W. D. Oliver, F. Yamaguchi, and Y. Yamamoto, Phys. Rev. Lett. 88, 037901 (2002).
16. O. Sauret, D. Feinberg and T. Martin, Eur. Phys. J. B 32, 545 (2003); *ibid.*, Phys. Rev. B 69, 035332 (2004).