

# Luttinger liquid versus charge density wave behaviour in the one-dimensional spinless fermion Holstein model

H. Fehske<sup>a,\*</sup>, G. Wellein<sup>b</sup>, G. Hager<sup>b</sup>, A. Weiße<sup>c</sup>, K. W. Becker<sup>d</sup>, A.R. Bishop<sup>e</sup>

<sup>a</sup>*Institut für Physik, Universität Greifswald, D-17487 Greifswald, Germany*

<sup>b</sup>*Regionales Rechenzentrum Erlangen, Universität Erlangen-Nürnberg, Germany*

<sup>c</sup>*School of Physics, The University of New South Wales, Sydney, NSW 2052, Australia*

<sup>d</sup>*Institut für Theoretische Physik, TU Dresden, D-01062 Dresden, Germany*

<sup>e</sup>*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

---

## Abstract

We discuss the nature of the different ground states of the half-filled Holstein model of spinless fermions in 1D. In the metallic regime we determine the renormalised effective coupling constant and the velocity of the charge excitations by a density-matrix renormalisation group (DMRG) finite-size scaling approach. At low (high) phonon frequencies the Luttinger liquid is characterised by an attractive (repulsive) effective interaction. In the charge-density wave Peierls-distorted state the charge structure factor scales to a finite value indicating long-range order.

*Key words:* non-Fermi liquid behaviour, quantum phase transition, Mott-Hubbard systems

---

The challenge of understanding quantum phase transitions in novel quasi-1D materials has stimulated intense work on microscopic models of interacting electrons and phonons such as the Holstein model of spinless fermions (HMSF)

$$H = -t \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) + \omega_0 \sum_i b_i^\dagger b_i - g\omega_0 \sum_i (b_i^\dagger + b_i)(n_i - \frac{1}{2}). \quad (1)$$

The HMSF describes tight-binding band electrons coupled locally to harmonic dispersionless optical phonons, where  $t$ ,  $\omega_0$ , and  $g$  denote the electronic transfer amplitude, the phonon frequency, and the electron-phonon (EP) coupling constant, respectively.

Despite of its simplicity the HMSF is not exactly solvable and a wide range of numerical methods has been applied in the past to map out the ground-state phase diagram in the  $g$ - $\omega_0$ -plane, in particular for the

half-filled band case ( $N_{el} = N/2$ ). There, the model most likely exhibits a transition from a Luttinger liquid (LL) to a charge-density wave (CDW) ground state above a critical EP coupling strength  $g_c(\omega_0) > 0$  [1].

In this contribution we present large-scale DMRG calculations, providing unbiased results for the (non-universal) LL parameters  $u_\rho$ ,  $K_\rho$ , and the staggered charge structure factor  $S_c(\pi)$ .

To leading order, the charge velocity  $u_\rho$  and the correlation exponent  $K_\rho$  might be obtained from a finite-size scaling of the of the ground-state energy of a finite system  $E_0(N)$  with  $N$  sites

$$\varepsilon_0(\infty) - (E_0(N)/N) = (\pi/3)(u_\rho/2)/N^2 \quad (2)$$

( $\varepsilon_0(\infty)$  denotes the bulk ground-state energy density) and the charge excitation gap

$$\Delta_{ch}(N) = E_0^{(\pm 1)}(N) - E_0(N) = \pi(u_\rho/2)/(NK_\rho) \quad (3)$$

(here  $E_0^{(\pm 1)}(N)$  is the ground-state energy with  $\pm 1$  fermions away from half-filling). The LL scaling relations (2) and (3) were derived for the pure electronic spinless fermion model only [2].

---

\* Corresponding Author: Phone: +49-3834-86-4760, Fax +49-3834-86-4701, Email: fehske@physik.uni-greifswald.de




Fig. 1. Finite-size scaling of the charge gap  $\Delta_{ch}(N)$  and the ground-state energy  $E_0(N)$ . Exact diagonalisation (ED) data is included for comparison.

Figure 1 demonstrates, exemplarily for the adiabatic regime, that they also hold for the case that a finite EP coupling is included. The resulting LL parameters are given in Tab. 1. Most notably, around  $\omega_0/t \sim 1$ , the LL phase splits in two different regimes: For small phonon frequencies the effective fermion-fermion interaction is *attractive*, while it is *repulsive* for large frequencies. In the latter region the kinetic energy is strongly reduced and the charge carriers behave like (small) polarons. In between, there is a transition line  $K_\rho = 1$ , where the LL is made up of (almost) non-interacting particles. The LL scaling breaks down just at  $g_c(\omega/t)$ , i.e. at the transition to the CDW state. We found  $g_c^2(\omega/t = 0.1) \simeq 7.84$  and  $g_c^2(\omega/t = 10) \simeq 4.41$  [3].

$g^2$	$\omega_0/t = 0.1$		$\omega_0/t = 10.0$	
	$K_\rho$	$u_\rho/2$	$K_\rho$	$u_\rho/2$
0.6	1.031	$\sim 1$	$\sim 1$	0.617
2.0	1.055	0.995	0.949	0.146
4.0	1.091	0.963	0.651	0.028

Table 1  
LL parameters at small and large phonon frequencies.

Figure 2 proves the existence of the long-range ordered CDW phase above  $g_c$ . Here the charge structure factor

$$S_c(\pi) = \frac{1}{N^2} \sum_{i,j} (-1)^j \langle (n_i - \frac{1}{2})(n_{i+j} - \frac{1}{2}) \rangle \quad (4)$$

unambiguously scales to a finite value in the thermodynamic limit ( $N \rightarrow \infty$ ). Simultaneously  $\Delta_{ch}(\infty)$  acquires a finite value. In contrast we have  $S_c(\pi) \rightarrow 0$  in the metallic regime ( $g < g_c$ ). The CDW for strong EP coupling is connected to a Peierls distortion of the lattice, and can be classified as traditional band insula-




Fig. 2. Scaling of the charge structure factor  $S_c(\pi)$  using periodic (PBC) and open (OBC) boundary conditions.

tor and bipolaronic insulator in the strong-EP coupling adiabatic and anti-adiabatic regimes, respectively.

The emerging physical picture can be summarised in the schematic phase diagram shown in Fig. 3. In the adiabatic limit ( $\omega_0 \rightarrow 0$ ) any finite EP coupling causes a Peierls distortion. In the anti-adiabatic strong EP coupling limit ( $\omega_0 \rightarrow \infty$ ), the HMSF can be mapped perturbatively onto the XXZ model and the metal-insulator transition is consistent with a Kosterlitz-Thouless transition at  $g_c^2(\infty) \simeq 4.88$ .




Fig. 3. Phase diagram of the 1D half-filled spinless fermion Holstein model. Here  $\alpha = \omega_0/t$ ,  $\varepsilon_p = \omega_0 g^2$ , and  $\lambda = \varepsilon_p/2t$ .

Work was supported by DFG under SPP 1073 and SFB 463, KONWIHR, HLRN Berlin, and the US DOE.

## References

- [1] See, e.g., R. B. Bursill, R. H. McKenzie, and C. J. Hamer, Phys. Rev. Lett. **80**, 5607 (1998), S. Sykora et al., cond-mat/0402184, and references therein.
- [2] J. L. Cardy, J. Phys. A **17**, L385 (1984); J. Voit, Rep. Prog. Phys. **58**, 977 (1995).

- [3] H. Fehske, M. Holicki, and A. Weiße, *Advances in Solid State Physics*, **40**, 235 (2000).