

EUROPHYSICS LETTERS

Europhys. Lett., (), pp. ()

The Many Electron Ground State of the Adiabatic Holstein Model in Two and Three Dimensions

B. Poornachandra Sekhar, Sanjeev Kumar and Pinaki Majumdar

Harish-Chandra Research Institute,
Chhatnag Road, Jhusi, Allahabad 201 019, India

(received ; accepted)

PACS. 71.38. { Polarons and electron-phonon interaction.
PACS. 71.45.L { Charge density wave systems.
PACS. 74.20. { Theory.

Abstract. {

We present the complete ground state phase diagram of the Holstein model in two and three dimension considering the phonon variables to be classical. We first establish the overall structure of the phase diagram by using exact diagonalisation based Monte Carlo (ED-MC) on small lattices and then use a new "travelling cluster" approximation (TCA) for annealing the phonon degrees of freedom on large lattices. The phases that emerge include a Fermi liquid (FL), with no lattice distortions, an insulating polaron liquid (PL) at strong coupling, and a charge ordered insulating (COI) phase around half-filling. The COI phase is separated from the Fermi liquid by a regime of phase coexistence whose width grows with increasing electron-phonon coupling. We provide results on the electronic density of states, the COI order parameter, and the spatial organisation of polaronic states, for arbitrary density and electron-phonon coupling. The results highlight the crucial role of spatial correlations in this strong coupling problem.

Introduction. { Electron-phonon (EP) interactions are ubiquitous in metals and dominate the finite temperature resistivity in most electron systems. While the perturbative, Fermi liquid, regime in EP systems is well understood [1], the electronic ground state is fundamentally reorganised when EP interactions are strong. The presence of a strong local interaction can generate a large lattice distortion, creating a potential well for the electron, and lead to a 'self trapped' polaronic state [2]. Strong electron-phonon coupling, sometimes in conjunction with other interactions, form a crucial component in the physics of several correlated systems, e.g, the manganites [3], the nickelates [4], and the traditional charge density wave systems [5].

The full polaron problem, considering the quantum dynamics of the phonons and strong EP coupling, is well understood for a single (or few) electrons [6, 7, 8, 9, 10]. The non perturbative nature of the problem and the exponential growth in Hilbert space with electron number, N_{el} , has made the finite density problem difficult to access. Usually, the physical interest is in the "adiabatic" regime since typical phonon frequencies in most materials are much smaller than the electron hopping scale. Although adiabaticity by itself does not lead to a simpler problem,

the adiabatic limit, where phonon dynamics is ignored, leads to a relatively tractable situation. This limit, of electrons coupled to classical phonons, has been explored within dynamical mean field theory (DMFT) [11, 12, 13]. DMFT, however, loses out on spatial correlations or the possibility of accessing non periodic phases, important at strong coupling. In this paper we use an approach that explicitly retains spatial correlations and set out the ground state of many electron systems, in two and three dimension, coupled to adiabatic phonons.

Let us define the Holstein model, whose adiabatic limit we explore:

$$H = \sum_{ij} t_{ij} c_i^\dagger c_j + \sum_i \mu c_i^\dagger c_i + \sum_i n_i x_i + \frac{1}{2M} \sum_i p_i^2 + \frac{K}{2} \sum_i x_i^2 \quad (1)$$

The t are nearest neighbour hopping on a d dimensional lattice, μ is the chemical potential, and $n_i = c_i^\dagger c_i$ is the electron density operator for spinless fermions. The phonon coordinate is x_i , and p_i is the momentum conjugate to x_i , with $[x_i; p_j] = i \delta_{ij}$. M and K are respectively the mass and stiffness of the phononic oscillators, and μ is the EP coupling. We set $t = 1$, as our reference scale, put $K = 1$ and also $\hbar = 1$. The adiabatic limit sets $M \rightarrow \infty$. For "quantum" phonons, with $\omega_{ph} = \sqrt{K/M}$, the last three terms in H can be written as:

$$\sum_i n_i (\alpha_i^\dagger + \beta_i) + \omega_{ph} \sum_i \beta_i^\dagger \beta_i, \text{ using } x_i = (\alpha_i + \beta_i^\dagger) / \sqrt{2M \omega_{ph}}, \text{ and } p_i = \sqrt{2M \omega_{ph}} (\alpha_i - \beta_i^\dagger).$$

The parameter space of the model is defined by: (i) the coupling parameter $\mu = E_p = 2dt$, with $E_p = \omega_{ph}^2 = 2K$, (ii) the 'adiabaticity' $\omega_{ph} = t$, and (iii) electron density, n , in addition to dimensionality, d . The parameter μ quantifies the competition between trapping and delocalisation, while ω_{ph} measures the 'slowness' of the phonons compared to electron dynamics.

The ground state of this model is understood only in a few limiting cases: (i) At weak to moderate coupling, traditional Migdal-Eliashberg (ME) theory [14] can be used for the many electron problem for $d = 1, 2$, and describes a Fermi liquid (FL) metal (or a superconductor, when electron spin is also considered). (ii) At strong coupling, and for arbitrary d , there is no controlled analytic theory even for a single electron coupled to the phonon system. There are, however, powerful numerical methods and the 'single polaron' ground state is essentially understood [6, 7, 8, 9, 10]. Little is known of the many electron problem beyond ME theory. (iii) Controlled theory of strong coupling finite density systems has been possible only in the classical limit, $\omega_{ph} \rightarrow 0$, in the limit $d \rightarrow \infty$. The original study by Millis et al. [11] highlighted the FL and polaronic insulator phases, within DMFT, and other DMFT studies have explored charge ordering [12, 13].

While contributing to the initial understanding, DMFT misses out on some crucial aspects of the physics. (i) The spatial correlation of lattice distortions, and the resulting correlations between polarons, when they form, is not accessible, and (ii) the possibility of phase coexistence and cluster formation is hard to access. We need to move beyond DMFT to recover these physical effects, and address current materials issues. The method we use, described next, can handle many electron systems strongly coupled to adiabatic phonons, at arbitrary disorder, explicitly in 1-3 dimension. This paper focuses on the ground state of 'clean' systems, subsequent papers discuss the finite temperature effects and the role of disorder [15].

Method. { At low temperature the adiabatic EP problem reduces to finding the phonon configuration that would minimise the total energy: $E_{fxg} = \sum_{ij} t_{ij} c_i^\dagger c_j + \sum_i n_i \mu + \sum_i n_i x_i^2 + (K=2) \sum_i x_i^2$: At strong coupling the electronic energy (in angular brackets) in a phonon background, $\langle f_{xg} \rangle$, is not analytically known, since the problem is equivalent to that of electrons in an arbitrary strongly fluctuating landscape. The only unbiased way to solve the problem is to generate Monte Carlo samples of the phonon configurations and accept or reject them using the fermion energy obtained by exact diagonalisation [16]. Such




Fig. 1. { Phase diagrams at $T = 0$. FL indicates Fermi liquid (band metal), the chessboard pattern represents a charge ordered insulator with f ; ;g modulation, and PL is a positionally disordered insulating polaron liquid. Shading indicates regimes of phase separation. The phase diagrams are obtained through a combination of TCA and analytic estimates: (a) result on 2d, TCA with $L_c = 4$ and system size 24^2 , (b) result on 3d with $L_c = 4$ and system size 8^3 . The coexistence width at weak coupling is estimated as explained in the text.

exact diagonalisation based Monte Carlo (ED-MC) is feasible only for electronic Hilbert space dimension H_N upto 100 since the computation time T_N grows as H_N^4 . Although the finite size effects are quite strong, the small sizes do provide a feel for the phase diagram. Our 'reference results' in 2d and 3d are based on ED-MC on 8^2 and 4^3 systems.

To overcome the severe size limitation of ED-MC we have used a 'travelling cluster' approximation [17] (TCA) for estimating the energy change due to a phonon move. Thus, instead of estimating the energy change of the update $x_i \rightarrow x_i^0$, at site R_i , as $E = E_{fx}^0 - E_{fxg}$, by diagonalisation of the full Hamiltonian, we obtain an estimate from a 'cluster' Hamiltonian constructed by considering L_c^2 (or L_c^3) sites around the reference site R_i . The energy of the system as a whole of course cannot be approximated by the energy, $E_{cl}fR_i g$, of the cluster, but the change in the energy of the system when a phonon update is made is accurately captured by the smaller subsystem. We have broadly benchmarked TCA against ED-MC in an earlier paper [17], and also provide several comparisons between the exact and TCA results in this paper. Our large size TCA results are obtained by using 4^2 clusters on 24^2 systems in two dimension, and 4^3 clusters on 8^3 in three dimension.

To allow effective annealing we study the system at low finite temperature, $T = 0.02$, with varying β . At equilibrium, at a given β , we calculate the following: (i) the electron density, $n(\beta)$, (ii) the structure factor $S(q) = (1/N^2) \sum_{ij} \langle n_i n_j \rangle e^{iq \cdot (R_i - R_j)}$, including $S(\pi; \pi; \dots)$, the order parameter for commensurate charge ordering ($\langle n_i \rangle$ is the quantum average of n_i in a MC configuration and the outer angular brackets indicate average over configurations), (iii) the electronic density of states (DOS), $N(\omega)$, (iv) the distribution of lattice distortions, as well as MC snapshots of the electron density. Fig.1 sets out the ground state phase diagram in 2d and 3d, while the detailed indicators, like $n(\beta)$, $S(\pi; \pi; \dots)$, and $N(\omega)$, from which the phase diagram is constructed, are shown in Figs.2-3. To access the "weak coupling" part of the phase diagram, $\beta < 1.5$ say, we have used an analytic method described later.

Phases. { The two panels in Fig.1 show the asymptotic large L phase diagram obtained through a combination of TCA and analytic estimates. Our simulations reveal that there




Fig. 2. { Panels (a) and (b) show the variation in carrier density with μ computed with ED-MC (symbols) and TCA (lines). Panel (a) is for 2d, with ED on 8^2 and TCA with 4^2 on 8^2 , while (b) is for 3d, with ED on 4^3 and TCA with 3^3 on 4^3 . Panels (c) and (d) show the COI 'order parameter' with varying electron density and coupling. Panel (c) is for 2d, panel (d) is for 3d. Symbols for ED, lines for TCA, same system size as in (a) (b). The insets to the lower panels show the COI order parameter at $n = 0.5$. TCA on large sizes does not significantly change the T=0 order parameter. Panel (a) and (c) share a common legend, as do panel (b) and (d).

are primarily three phases in the adiabatic limit, for $T \neq 0$, in both 2d and 3d. These are (i) the FL phase, i.e, the tight binding electron system, without any lattice distortions, (ii) an insulating polaron liquid (PL) with 'liquid like' short range positional correlations, and a gap in the DOS, and (iii) charge ordered insulating (COI) phases, with a gap in the DOS, and a peak at $q = \pi$ in $S(q)$. The COI is either a simple 'chessboard' phase at $n = 0.5$, or a phase with 'defects' at $n = 0.5$ (and strong coupling). There is also the possibility of other charge ordered phases, apart from π , in the vicinity of $n = 1/4; 1/8$, etc, at strong coupling, but their energy difference with respect to the PL phase is very small, making it difficult to access them in a low but finite T simulation. There are no 'metallic' phases with lattice distortions, in contrast to the DMFT results in the adiabatic limit [11]. We discuss the phase diagram in terms of the two broad regimes: (i) strong coupling polaronic phases, and (ii) weak coupling: the COI instability and phase separation.

(i) Strong Coupling: A naive balance of the 'polaron binding energy' $E_p = \frac{1}{2} = 2K$ (for a site localised electron), and the lower edge of the tight binding band, $E_b = 2dt$, suggests that polaron formation would occur at $t_c = \frac{1}{4K}$, implying $\frac{2d}{c} = 2.82$ and $\frac{3d}{c} = 3.46$. More accurate earlier estimates [7], and our simulations, indicate $\frac{2d}{c} = 2.6$ and $\frac{3d}{c} = 3.3$. The lowering arises because the polaron is not quite site localised, and taking into account the kinetic energy $\frac{1}{2} dt^2 = E_p$, the polaronic state becomes favourable slightly before the naive threshold. The polaron states which emerge for $t > t_c$ are compact, although not site localised,




Fig. 3. { Density of states computed with TCA on 24×24 in 2d, and $8 \times 8 \times 8$ in 3d. Symbols: circles for $n = 0.1$, triangles for $n = 0.3$, and squares for $n = 0.5$. (a) 2d, $\lambda = 2.2$, (b) 2d, $\lambda = 3.0$, (c) 3d, $\lambda = 2.6$, (d) 3d, $\lambda = 3.5$.

with wavefunctions decaying over a couple of lattice spacings. Due to the strong localisation, and the associated short range lattice distortion field, the finite density system is a short range correlated array of trapped electrons (the PL phase), keeping maximum mutual separation in order to maximize kinetic energy gain by virtual hopping. In the dilute limit the polaron system has a very large compressibility, $\partial n / \partial \mu$, since the polarons are effectively non interacting and is virtually 'pinned' to the single polaron energy as the carrier density is increased from zero. This dictates the sharp rise in the n curves, Fig.2.(a) (b). However, with n increasing towards half-filling the short range 'repulsion', U ($U = E_p$) for nearest neighbours, begins to be felt and the n characteristic is flat at $n = 0.5$, indicative of an incompressible half-filled state. The polaronic repulsion favours a "chessboard" CO phase at $n = 0.5$.

The ordered half-filling phase maintains the positional correlations of the $n = 0.5$ phase, but has vacancies (or defects), see e.g, Fig.4. The polarons tend to avoid nearest neighbour locations, preferring to be along the diagonal (where the 'repulsion' is weaker). As long as the system is sufficiently dense, these positional correlations 'percolate' sustaining long range order, with a clear peak in $S(\mathbf{k}; \omega)$. Our results suggest that the strong coupling ordered half-filling CO I phase survives down to a density $n \approx 0.35$ in the coupling regime shown (Fig.1). The order parameter for the CO I phases is shown in Fig.2.(c) (d), and the DOS in Fig.3. The order parameter computed with TCA on small sizes, see Fig.2 caption, accurately matches ED-MC results. TCA results on large sizes for $S(\mathbf{k}; \omega)$ (not shown) are not significantly different.

The critical coupling for the FL to PL transition is density dependent, decreasing signif-




Fig. 4. Monte Carlo snapshots of the electron density n_r with varying n and μ in 2d. Top panels $n = 2.2$, bottom panels $n = 3.0$. Density $\mu = 0.10; 0.30; 0.50$, left to right. Top left panel is FL. Both panels to the extreme right are 'perfect' COI, while the other three panels are PL.

icantly, Fig.1 (c) (d), from the single polaron threshold as n is increased. By the time $n = 0.3 - 0.4$, μ_c is reduced to $\sim 75\%$ of the 'single polaron' value, in both 2d and 3d, i.e. the critical E_p is almost halved. Qualitatively, electrons which are already localised affect an added particle via their strongly inhomogeneous distortion field, $\phi(x)$, which interplays with EP coupling leading self-consistently to the decreasing $\mu_c(n)$. Trapping at finite n involves a combination of (self-generated) "disorder" and polaronic tendency [18], akin to the interplay of Anderson localisation and single polaron formation. All the phases, COI or PL, which occur for $\mu_c(n)$ have a clear spectral gap, and are distinguished mainly by $S(q)$.

(ii) Weak Coupling: At weak coupling there is an instability in a bipartite tight binding lattice, at $n = 0.5$, due to Fermi surface nesting. The density response function $\chi(q; \omega = 0)$ diverges for $q \neq \pi$; $\omega = \pi$. For $\omega = 0$, where the analysis in terms of the non-interacting susceptibility is valid, this leads to a charge order instability, with ordering temperature $T_{CO} \propto t^{E_p}$. With increasing μ the CO phase continues to be the ground state, but as the charge modulation increases, and the band splitting grows, the FL to COI transition at $T = 0$ becomes first order, with respect n , since the electron density in the modulated x_1 background is significantly different from that in the homogeneous background. The n data in Fig.2. (a) (b) highlight the discontinuity in n near half-filling. It is difficult to access the instability and the coexistence width at weak coupling in a small system since $\chi(q; \omega = 0)$ has strong size dependence. Direct simulation can locate the CO phase only down to $\mu \sim 1.5$. At lower μ we variationally computed the COI order parameter at $n = 0.5$ on large lattices, and also the $\mu_c(\mu)$ at which the FL to COI transition occurs. It suggests that the FL to COI transition becomes (noticeably) first order, with a density discontinuity, for $\mu \sim 1.0$, in both 2d and 3d. The weak coupling COI phase at $n = 0.5$ connects continuously to the strong coupling "polaron ordered" phase [12, 13].

Discussion. Despite accurate handling of strong coupling and spatial correlations, a few physical effects in the adiabatic limit are still difficult to capture within a numerical simulation.

