

Towards a Hydrodynamic Theory of Infinite Neutral Nonrelativistic Matter

Girish S. Setlur

Harish Chandra Research Institute
Chhatnag Road, Jhusi, Allahabad, India 211 019.

March 22, 2022

Abstract

We recast the problem of infinite neutral nonrelativistic matter interacting via U(1) gauge fields in the hydrodynamic language. We treat the nuclei as being spinless bosons for simplicity (for example in He4). We write down the formal action in terms of a full set of independent hydrodynamic gauge invariant variables. The claim is that the results of this theory are nonperturbative and nuclei and electrons are treated on an equal footing.

1 The Theory

Consider the action for neutral matter composed of electrons and nuclei which we assume are all spin-zero point particles.

$$S = \sum_{\sigma} \int \psi_e^{\dagger}(\mathbf{x}\sigma) \left(i \frac{\partial}{\partial t} + \frac{\nabla^2}{2m_e} \right) \psi_e(\mathbf{x}\sigma) + \int \psi_N^{\dagger}(\mathbf{x}) \left(i \frac{\partial}{\partial t} + \frac{\nabla^2}{2M_N} \right) \psi_N(\mathbf{x}) \quad (1)$$

Here $\int = \int_0^{-i\beta} dt \int d^3x$. We want this to be invariant under gauge transformations.

$$\psi_e(\mathbf{x}\sigma) \rightarrow e^{-ie\theta(\mathbf{x})} \psi_e(\mathbf{x}\sigma) \quad (2)$$

$$\psi_N(\mathbf{x}) \rightarrow e^{iZ_N e\theta(\mathbf{x})} \psi_N(\mathbf{x}) \quad (3)$$

Thus we have to couple to gauge fields. The invariant action with gauge fields reads as follows.

$$S = \int \psi_e^{\dagger} \left(i\partial_t + e\phi - \frac{(-i\nabla + \frac{e}{c}\mathbf{A})^2}{2m_e} \right) \psi_e + \int \psi_N^{\dagger} \left(i\partial_t - Z_N e\phi - \frac{(-i\nabla - Z_N \frac{e}{c}\mathbf{A})^2}{2M_N} \right) \psi_N \\ - \frac{e}{m_e c} \int \mathbf{B} \cdot \psi_e^{\dagger} \mathbf{S} \psi_e - \int \frac{1}{4} F^2 \quad (4)$$

Thus in addition we have to perform the replacements,

$$A_0 \rightarrow A_0 + \partial_t \theta \quad (5)$$

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla \theta \quad (6)$$

Now we polar decompose the fields,

$$\psi_e = e^{i\Phi_e} e^{-i\Pi_e} \sqrt{\rho_e} \quad ; \quad \psi_N = e^{-i\Pi_N} \sqrt{\rho_N} \quad (7)$$

We further postulate that $\nabla \Phi_e = 0$, this additional phase reflects the fermionic character of ψ_e .

$$\begin{aligned} S = & \int \sqrt{\rho_e} \left(C_e - \partial_t \Phi_e + \frac{\nabla^2 - \mathbf{C}_e^2 + i(\nabla \cdot \mathbf{C}_e) + 2i\mathbf{C}_e \cdot \nabla}{2m_e} \right) \sqrt{\rho_e} \\ & + \int \sqrt{\rho_N} \left(-Z_N C_N + \frac{\nabla^2 - Z_N^2 \mathbf{C}_N^2 - iZ_N(\nabla \cdot \mathbf{C}_N) - 2iZ_N \mathbf{C}_N \cdot \nabla}{2M_N} \right) \sqrt{\rho_N} \\ & - \frac{e}{m_e c} \int \mathbf{B} \cdot \sqrt{\rho_e(\mathbf{x}\beta)} e^{i\Pi_e(\mathbf{x}\beta)} e^{-i\Phi_e(\mathbf{x}\beta)} \mathbf{S}_{\beta\alpha} e^{i\Phi_e(\mathbf{x}\alpha)} e^{-i\Pi_e(\mathbf{x}\alpha)} \sqrt{\rho_e(\mathbf{x}\alpha)} \\ & + \int \frac{1}{2} (\mathbf{E}^2 + \mathbf{B}^2) \end{aligned} \quad (8)$$

$$e\phi + \partial_t \Pi_e = C_e \quad (9)$$

$$\partial_t \Pi_N - Z_N e \phi = -Z_N C_N \quad (10)$$

$$-\nabla \Pi_e + \frac{e}{c} \mathbf{A} = \mathbf{C}_e \quad (11)$$

$$-\nabla \Pi_N - Z_N \frac{e}{c} \mathbf{A} = -Z_N \mathbf{C}_N \quad (12)$$

$$\mathbf{E} = -\nabla \phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} \quad (13)$$

$$\mathbf{B} = \nabla \times \mathbf{A} \quad (14)$$

$$-Z_N \nabla \Pi_e - \nabla \Pi_N = Z_N \mathbf{C}_e - Z_N \mathbf{C}_N \quad (15)$$

$$Z_N \partial_t \Pi_e + \partial_t \Pi_N = Z_N C_e - Z_N C_N \quad (16)$$

$$-Z_N \nabla \partial_t \Pi_e - \nabla \partial_t \Pi_N = Z_N \partial_t \mathbf{C}_e - Z_N \partial_t \mathbf{C}_N \quad (17)$$

$$Z_N \partial_t \nabla \Pi_e + \partial_t \nabla \Pi_N = Z_N \nabla C_e - Z_N \nabla C_N \quad (18)$$

$$\begin{aligned}
S = & \int \left(\rho_e C_e - \rho_e \partial_t \Phi_e - \frac{(\nabla \rho_e)^2}{4\rho_e} + \rho_e \mathbf{C}_e^2 \right) \\
& + \int \left(-\rho_N Z_N C_N - \frac{(\nabla \rho_N)^2}{4\rho_N} + \rho_N Z_N^2 \mathbf{C}_N^2 \right) \\
& - \frac{e}{m_e c} \int \mathbf{B} \cdot \rho_e(\mathbf{x}\alpha) \mathbf{S}_{\alpha\alpha} - \frac{e}{m_e c} \int \mathbf{B} \cdot \sqrt{\rho_e(\mathbf{x}\bar{\alpha})} e^{i(\Pi_e(\mathbf{x}\bar{\alpha}) - \Pi_e(\mathbf{x}\alpha))} e^{i(\Phi_e(\mathbf{x}\alpha) - \Phi_e(\mathbf{x}\bar{\alpha}))} \mathbf{S}_{\bar{\alpha}\alpha} \sqrt{\rho_e(\mathbf{x}\alpha)} \\
& + \int \frac{1}{2} (\mathbf{E}^2 + \mathbf{B}^2)
\end{aligned} \tag{19}$$

$$0 = \partial_t \mathbf{C}_e - \partial_t \mathbf{C}_N + \nabla C_e - \nabla C_N \tag{20}$$

$$\frac{e}{c} \mathbf{B} = \nabla \times \mathbf{C}_N \tag{21}$$

$$e \mathbf{E} = -\partial_t \mathbf{C}_N - \nabla C_N \tag{22}$$

$$\partial_t \Pi_e(\mathbf{x}\bar{\alpha}) - \partial_t \Pi_e(\mathbf{x}\alpha) = C_e(\mathbf{x}\bar{\alpha}) - C_e(\mathbf{x}\alpha) \tag{23}$$

We may now write in terms of the Matsubara frequencies,

$$\Pi_e(\mathbf{x}\bar{\alpha}, t) - \Pi_e(\mathbf{x}\alpha, t) = \sum_n e^{z_n t} \frac{C_e(\mathbf{x}\bar{\alpha}, z_n) - C_e(\mathbf{x}\alpha, z_n)}{z_n} \tag{24}$$

$$\mathbf{C}_e(\mathbf{x}\alpha, t) = \mathbf{C}_N(\mathbf{x}, t) - \sum_n \frac{e^{z_n t}}{z_n} (\nabla C_e(\mathbf{x}\alpha, z_n) - \nabla C_N(\mathbf{x}, z_n)) \tag{25}$$

Therefore the effective action in terms of the independent gauge invariants reads as follows :

$$\begin{aligned}
S = \int & \left(\rho_e C_e - \rho_e \partial_t \Phi_e - \frac{(\nabla \rho_e)^2}{4\rho_e} + \rho_e (\mathbf{C}_N(\mathbf{x}, t) - \sum_n \frac{e^{z_n t}}{z_n} (\nabla C_e(\mathbf{x}\alpha, z_n) - \nabla C_N(\mathbf{x}, z_n)))^2 \right) \\
& + \int \left(-Z_N \rho_N C_N - \frac{(\nabla \rho_N)^2}{4\rho_N} + \rho_N Z_N^2 \mathbf{C}_N^2 \right) \\
& - \frac{1}{m_e} \int \nabla \times \mathbf{C}_N \cdot \rho_e(\mathbf{x}\alpha) \mathbf{S}_{\alpha\alpha} \\
& - \frac{1}{m_e} \int \nabla \times \mathbf{C}_N \cdot \sqrt{\rho_e(\mathbf{x}\bar{\alpha})} e^i \sum_n e^{z_n t} \frac{C_e(\mathbf{x}\bar{\alpha}, z_n) - C_e(\mathbf{x}\alpha, z_n)}{z_n} e^{i(\Phi_e(\mathbf{x}\alpha) - \Phi_e(\mathbf{x}\bar{\alpha}))} \mathbf{S}_{\bar{\alpha}\alpha} \sqrt{\rho_e(\mathbf{x}\alpha)} \\
& + \int \frac{1}{2e^2} [(\partial_t \mathbf{C}_N + \nabla C_N)^2 + c^2 (\nabla \times \mathbf{C}_N)^2] \quad (26)
\end{aligned}$$

where the fermionic phase functional is given by

$$\Phi_e([\rho]; \mathbf{x}\alpha) = \sum_{\mathbf{q}, n} \rho_{-\mathbf{q}, -n, \alpha} C(\mathbf{q}, \alpha, n) e^{i\mathbf{q} \cdot \mathbf{x}} e^{z_n t} \quad (27)$$

$$\beta z_n C(\mathbf{q}, \alpha, n) = \frac{1}{2 \langle \rho_{\mathbf{q}\alpha, n} \rho_{-\mathbf{q}\alpha, -n} \rangle_0} - \frac{(2m)\beta z_n^2}{2N^0 \mathbf{q}^2} - \frac{\beta \mathbf{q}^2}{2N^0(2m)} \quad (28)$$

where $\langle \rho_{\mathbf{q}\alpha, n} \rho_{-\mathbf{q}\alpha, -n} \rangle_0$ is the density-density correlation function of the noninteracting Fermi (electron) theory.