

# Mean field dynamics of superfluid-insulator phase transition in a gas of ultra cold atoms

Jakub Zakrzewski

Instytut Fizyki im. Mariana Smoluchowskiego and Mark Kac Complex Systems Research Center,  
Uniwersytet Jagielloński, ulica Reymonta 4, PL-30-059 Kraków, Poland  
(Dated: 1st September 2022)

A large scale dynamical simulation of the superfluid to Mott insulator transition in the gas of ultra cold atoms placed in an optical lattice is performed using the time dependent Gutzwiller mean field approach. This approximate treatment allows us to take into account most of the details of the recent experiment [Nature 415, 39 (2002)] where by changing the depth of the lattice potential an adiabatic transition from a superfluid to a Mott insulator state has been reported. Our simulations reveal a significant excitation of the system with a transition to insulator in restricted regions of the trap only.

PACS numbers: 03.75.-b, 03.75.Kk, 03.75.Lm

## I. INTRODUCTION

A theoretical suggestion [1] of a possibility to realize one of the standard models for interacting particles – the Bose-Hubbard (BH) model [2, 3] in a cold gas placed in an optical lattice has been followed soon by a seminal experiment [4]. The reported realization of a quantum phase transition between superfluid (SF) and Mott insulator (MI) phases showed convincingly that it was possible to control experimentally parameters of the model practically at will. This triggered several studies involving Bose condensate [5, 6, 7, 8, 9, 10] as well as, more recently Fermi-Bose mixtures [11, 12, 13] placed on the optical lattices (the reference list must be not complete bearing in mind that more than 70 papers with "optical lattice" in the title are listed in the cond-mat archive last year only).

At the same time a number of groups [14, 15, 16, 17, 18] tried to understand the details of the very first experiment [4] to check the underlying physics. To imagine the difficulty in modeling the experiment let us recall that it involves about  $10^5$  interacting atoms (bosons) placed in the harmonic trap and the three dimensional (3D) lattice potential. Such a system is well described by a Bose-Hubbard model with position dependent chemical potential [1]. Even finding the ground state of the system for that number of particles and  $65 \times 65 \times 65$  lattice sites is a formidable task. State of the art quantum Monte Carlo (QMC) [14, 15, 17] calculations aimed at the ground state properties include up to 16 sites in 3D [14], more sites may be included in one (1D) or two (2D) dimensional models [15, 17]. These studies, while interesting on their own, can shed little light on the dynamics of the system when its parameters are varied. Except for special exactly solvable models, the efficient simulation of time-dependent properties of interacting many-body system remains an open problem although recently quite a progress has been obtained for 1D systems [20, 21].

It seems, therefore, that the only reasonable and tractable way of analyzing the dynamics of the discussed experiment is using approximate methods. To this end

we shall use an approach based on the time dependent variational principle with Gutzwiller ansatz. That will allow us to model the details of the 3D experiment [4]. The prize for it is similar to that paid in other approximate treatments – one may always question the extent to which the approximations allow to describe the properties of the systems studied. We hope to convince the reader that the numerical results are at least mutually consistent and thus may provide considerable insight into the dynamics of the experiment.

The discussion of the dynamics is postponed to Section III since we discuss in the next Section the static mean field solutions for the ground state for experimental parameters. Here a comparison with available exact QMC results is possible at least. This shall give us some confidence about the applicability of the mean field approach yielding, at the same time, the initial state for the dynamics studied later.

## II. STATIC MEAN FIELD FOR THE BOSE-HUBBARD MODEL

The Bose-Hubbard Hamiltonian describing the system takes the form [1]

$$H = \sum_{\langle i,j \rangle} J a_i^\dagger a_j + \frac{U}{2} \sum_i n_i (n_i - 1) + \sum_i W_i n_i \quad (1)$$

where  $n_i = a_i^\dagger a_i$  is an occupation number operator at site  $i$  (with  $a_i$  being the corresponding annihilation bosonic operator),  $U$  the interaction energy,  $J$  the tunneling coefficient and  $W_i$  the energy offset at site  $i$ .  $\sum_{\langle i,j \rangle}$  denotes a sum over nearest neighbors. Both  $J$  and  $U$  are functions of the lattice potential and may be easily expressed in terms of integrals of the Wannier functions of the lowest energy band for cold atoms implementation of the model [1].

Consider first the standard homogeneous situation in which all  $W_i$ 's are equal. The last term in (1) becomes proportional to the (conserved) number of bosons and

may be dropped. The only remaining parameter of the model is the ratio of  $U=J$ . When tunneling dominates the system in its ground state is superfluid while in the opposite case it becomes the Mott insulator. The borderline between the two phases depends on the chemical potential. The Mott insulator state is incompressible and is characterized by an integer mean occupation of sites. In effect starting from the superfluid at small  $U=J$  and a non integer ratio of  $N=M$  ( $N$  denotes the number of atoms while  $M$  the number of sites) and increasing  $U=J$  the ground state remains superfluid up to highest values of  $U=J$  at fixed boson density. On the other hand the range of values corresponding to the commensurate filling increases with  $U=J$ . In effect, the separation line between a MI and a SF forms characteristic lobes [2, 3, 16],

For a detailed discussion of the BH model see [2, 16]. As we are interested in the mean field approximation, let us just quote Zwerger [16] saying "In two and three-dimensional lattices, the critical value for the transition from a MI to a SF is reasonably well described by a mean-field approximation". In one dimension the mean field approximation is much worse [16].

In the presence of the additional potential, e.g. the harmonic trap, local energies  $W_i$  depend on the sites location. Then the effective chemical potential at each site becomes  $\mu_i = \mu + W_i$ . As pointed out already in [1] this will lead for large  $U=J$  to a shell like structure with MI phases with different integer occupations (highest in the middle assuming attractive binding additional potential) separated by SF regions. This picture has been nicely confirmed in quantum Monte Carlo calculations both in 1D [15] and in 3D [14].

The latter exact results are of particular interest for us since they allow for a comparison with the mean field approximation. In [14] a 3D  $16 \times 16 \times 16$  lattice is considered with different values of  $U$  and  $J$  parameters as well as the harmonic trap. To find the mean field ground state we minimize

$$\langle E \rangle = \langle G | \hat{H} | \hat{N} | G \rangle; \quad (2)$$

where  $\hat{N} = \sum_i n_i$  and  $|G\rangle$  is the Gutzwiller trial function

$$|G\rangle = \sum_{i=1}^M \sum_{n=0}^{X_m} (f_n^{(i)} |n\rangle_i); \quad (3)$$

The number of parameters  $f_n^{(i)}$  depends on the number of sites (here  $16^3$ ) as well as the maximal occupation at a given site  $n_m$ . The average maximal occupation at the center of the trap is 2 for the data considered in [14]. Therefore, it is sufficient to take  $n_m = 7$ . That yields a minimization procedure over 32768 parameters. Such a number of parameters must lead to a spurious local minimum, unless a very good estimate exists for the initial set of  $f_n^{(i)}$ 's (i.e. the initial  $|G\rangle$ ). Fortunately such a guess is quite obvious and is often termed a local mean




Figure 1: Mean field particle density distribution (on-site filling factors) as a function of the distance from the center of the trap measured in the units of the lattice constant  $a$ . Filled circles correspond to numerical results, lines are drawn to guide the eye. The parameters of the BH model match exactly those quoted in Fig. 1 of [14]. Comparison with the latter results obtained within "exact" quantum Monte Carlo reveals the accuracy of the mean field approximation.

field approximation. Namely at each site  $i$  one takes a solution for  $f_n^{(i)}$ 's corresponding to the homogeneous BH model with the effective chemical potential  $\mu_i = \mu + W_i$ . Provided  $W_i$  changes smoothly from site to site, such an approach should be an excellent approximation to a full mean field solution. And indeed it is, we have found for the data discussed below that the initial and final  $\langle E \rangle$  [see Eq. (2)] differ by at most 2%; the number of iterations of standard Numerical Recipes minimization packages [22] is slightly bigger than the number of parameters.

The results obtained are presented in Fig. 1 in the same form as the corresponding plot in [14] to make the comparison easier. The parameter values correspond precisely to those taken in [14]. Please note that the tunneling constant  $J$  is denoted as  $t$  in [14]. A comparison of both figures indicates that, as far as the average occupation at different sites is concerned, the mean field solution is in excellent agreement with the quantum Monte Carlo results.

Instead of the occupation at different sites one may take a look at the momentum distribution, i.e. the quantity closely related to that measured in the experiment (see [4] and the discussion below). The momentum dis-




Figure 2: The rescaled mean field momentum distribution (in arbitrary units) in the first Brillouin zone in the (0,0,1) direction derived from (4) for data presented in Fig. 1). The distributions are to be compared with the corresponding "exact" distributions in Fig. 2 of [14]. Observe that mean field result differs from the exact distribution for the case (d) only (see discussion in the text).

tribution is given by [14]

$$n_k = \sum_{i,j} \langle k | \hat{f}^X e^{ik \cdot (r_i - r_j)} \langle a_i^y a_j^y \rangle; \quad (4)$$

where  $k$  is the wavevector,  $\langle k |$  a Fourier transform of the Wannier site-function. The latter yields a broad bell-shaped background and provides merely information about the lattice. The relevant information about the atoms is contained in the Fourier transform of  $\langle a_i^y a_j^y \rangle$ . In the mean field approximation this factorizes for different  $i, j$ :  $\langle a_i^y a_j^y \rangle \approx \langle a_i^y \rangle \langle a_j^y \rangle$ . Such a factorization seems quite drastic and one may expect significant differences between the momentum distributions obtained from QMC and within the mean field approximation. It is really not so, however, for bosons in a harmonic trap as visualized in Fig. 2 for the mean field. That figure should be compared with QMC results presented in Fig. 2 of [14]. Observe that differences appear only for panel (d), the exact results yield significantly broader momentum distribution. As discussed in [14] almost no SF fraction is present in the QMC result corresponding to panel (d). Then the factorization must affect strongly the momentum distribution since in a vast majority of sites  $\langle a_i \rangle = 0$ . Clearly, however, as long as some SF fraction is present in the system the mean field momentum distribution closely resembles the exact quantum results.

This apparent quite close agreement between QMC results and the mean field approximation for  $16 \times 16 \times 16$  lattice and about  $10^3$  atoms is very encouraging in view of realistic experimental conditions [4]. Here both the external potential changes less rapidly (the size of the lattice is now  $65 \times 65 \times 65$ ) and number of atoms exceeds

$10^5$  thus one may expect that the mean field approximation works even better.

While the test described above have been taken for some chosen (by authors of [14]) values of  $U, J$ , as well as the trap frequency, to simulate the experiment we have to determine first the relevant range of parameters. From now on we shall measure the quantities of dimension of energy in the units of the recoil energy of  $^{87}\text{Rb}$  atoms for light with a wavelength  $\lambda = 2\pi/K = 852\text{nm}$ , i.e.  $E_r = \hbar^2 K^2 / 2M$ , where  $M$  is the mass of  $^{87}\text{Rb}$  atoms. The depth of the optical lattice  $V$  changes from 0 to  $22E_r$ . Finding Wannier functions for different values of  $V$  [23] we evaluate the corresponding  $U(V)$  and  $J(V)$  values. The energy offset at each site  $W_i$  has two components in the experiment [4]. One is the harmonic magnetic trap potential (time-independent), another is due to the Gaussian intensity profiles of lattice creating laser beams. The latter may be also approximated by a harmonic term [4] the corresponding frequency is then dependent on  $V$ .

To find the mean field ground state for different  $V$  values and the number of atoms of the order of  $10^5$  one needs to solve a minimization problem over  $2 \times 10^6$  parameters (with  $n_m = 7$  as before) which is hardly manageable. One may, however, use the symmetry of the problem (cubic lattice combined with spherically symmetric trap) to significantly reduce that number. Let  $i, j, k$  count the sites in  $x, y, z$  directions, respectively with each index taking the values from  $-32$  to  $n_s - 32$  (yielding 65 sites in each direction). Due to the ground state symmetry it is enough to consider only the sites with  $0 \leq i, j, k \leq n_s$  which reduces the number of minimized parameters to about 48 thousands. Needless to say we have checked on the smaller  $16 \times 16 \times 16$  problem that the symmetry




Figure 3: Solid lines represent mean field atom density distribution (on-site filling factor  $\bar{n}$ ) as a function of the distance from the center of the trap (measured in units of the lattice constant). Dashed lines represent the corresponding variances at different sites  $2\sigma_i^2 = 2(\langle n_i^2 \rangle - \langle n_i \rangle^2)$ . The total number of atoms and the SF factor  $S_F$  (see text) for each plot are (a):  $V = 9E_r, N = 99771, S_F = 0.95$ ; (b):  $V = 13E_r, N = 99502, S_F = 0.40$ ; (c):  $V = 16E_r, N = 95408, S_F = 0.11$ ; (d):  $V = 22E_r, N = 94172, S_F = 0.01$ .

reduced problem yields the same ground state as the full minimization.

The results obtained are presented in Fig. 3 and are practically indistinguishable from the initial guess i.e. a wavefunction coming from local mean field approximation discussed above. The chemical potential has been adjusted (for each case) to have the average number of atoms  $N = \langle \hat{N} \rangle = \sum_i \langle n_i \rangle$  around  $10^5$ . This leads to more than two atoms (on average) per site in the center of the trap. To characterize whether the state is closer to being superfluid or Mott insulator we define the superfluid factor  $S_F = 1 - N \langle a_i^2 \rangle / \langle a_i \rangle^2$ . This factor is zero for pure MI state (when  $\langle a_i \rangle = 0$  as each node is in a Fock state) and reaches unity for Poissonian statistics at each node. While obviously it is not a "proper" order parameter for the phase transition, it seems to be convenient for characterizing the states obtained. Using this factor we can quantify states shown in Fig. 1, noting first a general qualitative agreement with experimental findings [4]. The case  $V = 9E_r$  seems almost fully superfluid (with, however, strongly subpoissonian statistics [24] at each site), the case  $V = 13E_r$  shows first traces of insulator phase (integer occupation of sites with van-

ishing variance, the transition is completed for significant fraction of sites at  $V = 16E_r$  while for the deepest lattice  $V = 22E_r$  SF fraction is restricted to very narrow regions separating different integer occupations.

### III. TIME-DEPENDENT MEAN FIELD DYNAMICS

The results obtained for mean field ground state in realistic situations, shown in the previous Section, seem quite encouraging. Yet, in themselves they can say little about the dynamics of the system when the lattice depth  $V$  is varied. In an attempt to address this important issue we shall use a time-dependent version of the mean field approximation. To this end we employ the time-dependent variational principle [8] looking for the minimum of

$$\langle G(t) | \frac{\partial}{\partial t} H(t) + \hat{N} G(t) \rangle; \quad (5)$$

with  $H(t)$  being now the time dependent Hamiltonian. The time dependence is implicit via the dependence of

the BH Hamiltonian  $H$  on  $U; J$  and  $W_i$ , that in turn depend on  $V$ . The chemical potential becomes also time dependent when system parameters are varied.  $|\Phi(t)\rangle$ , the variational wavefunction, is assumed in the standard Gutzwiller-type form (3), with  $f_n^{(i)}(t)$  now being time-dependent. The very same approach has been successfully applied recently to the formation of molecules [8, 9], the treatment of the disordered optical lattices [10] as well as for determining the phase diagram in Bose-Fermi mixtures [13].

The minimization of (5) yields the set of first order differential equations for  $f_n^{(i)}(t)$ :

$$i\frac{d}{dt}f_n^{(i)} = \frac{U}{2}n(n-1) + n(W_i) f_n^{(i)} - \sum_j \frac{J}{2} \left( f_{n+1}^{(i)} + f_{n-1}^{(i)} \right); \quad (6)$$

where  $\sum_i = \sum_{\langle i,j \rangle} < G(t) \hat{a}_j^\dagger \hat{a}_i(t) >$  (the sum, as indicated by subscript in brackets is over the nearest neighbors only). The nice feature of the evolution resulting from equations (6) is that the average number of particles  $N = \langle \hat{N} \rangle$  is an exact constant of the motion [8]. Naturally when the parameters of the BH model, e.g.  $U$  and  $J$ , change the chemical potential corresponding to the mean field solution with a given number of particles  $N$  also changes. The dynamics of cannot be obtained from (6) only. One can find it, however, following the evolution of two states  $|\Phi_1\rangle$  and  $|\Phi_2\rangle$  with slightly different average number of particles  $N_2 = \langle G_2 \hat{N} G_2 \rangle = N_1 + \Delta N = \langle G_1 \hat{N} G_1 \rangle + \Delta N$ . The chemical potential at given time may be then approximated by  $\mu(t) = \langle G_2(t) \hat{H}(t) G_2(t) \rangle - \langle G_1(t) \hat{H}(t) G_1(t) \rangle = \Delta N$  and adjusted at each time step [13]. This is the approach used in the numerical results presented below.

Since we want to follow as closely as possible the experiment [4] let us recall its main features. The experiment has three stages after loading the harmonic trap with Rb condensate – compare Fig. 4. Firstly, the optical lattice depth  $V(t)$  was increased in 80 ms (using exponential ramp with time constant  $\tau = 20$  ms) from the initial zero value (when the harmonic trap was present only) to  $V_{\max} = 22E_r$ , where  $E_r$  is the recoil energy of Rb atoms. The sample was then held for 20 ms at  $V_{\max}$ . Finally  $V(t)$  was decreased with the linear ramp to  $V_f = 9E_r$  with different speed. At any stage the experiment could be interrupted by rapidly switching off all laser beams building up the lattice as well as the magnetic trap. The freely expanding atomic cloud, after some delay, was recorded by a destructive absorption imaging, yielding the signal which reflects the momentum distribution [14, 16]. Since the absorption images are taken along two orthogonal axes the quantity measured is in fact the integrated momentum distribution [14]:

$$N(k_x; k_y) / \int dk_x n_k; \quad (7)$$




Figure 4: The experimental time profile (solid line) of the lattice potential depth  $V$  measured in the recoil energy  $E_r$  units. The initial exponential increase with a time constant  $\tau = 20$  ms is followed by a plateau and a subsequent decrease to  $V_f = 9E_r$  with a linear ramp with varying slope. Dashed line corresponds to the exponential time constant  $\tau = 40$  ms, dash-dotted  $\tau = 80$  ms. Thin dotted lines indicate particularly interesting values of time and  $V$  (see text for discussion).

For clouds released from low optical lattices when tunneling dominates and the superfluid behavior is expected the signal reflects Bragg peaks due to interferences of atoms coming from different lattice sites. At increased lattice depths above  $13E_r$  the interference maxima become immersed in an incoherent background disappearing practically at  $20E_r$ . This behavior was associated with the quantum phase transition from SF to Mott phase [4]. Most interestingly the coherence of the sample may be rapidly recovered when the lattice depth is decreased (third stage of the experiment) as measured by the width of the central interference peak which decreases almost to its original value at  $V = 9E_r$  in about 4 ms.

In the former applications of the time-dependent mean field approach [8, 9, 10, 13] the time-dependence of system's parameters was assumed to be sufficiently slow to assure adiabaticity. In effect the mean field ground state has been followed by applying the time-dependent equations for  $f_n^{(i)}(t)$ 's (6). Here we have a similar situation since it is claimed [4] that the changes in time of  $V(t)$  are made sufficiently slow to keep the system in the many-body ground state. Having the mean field ground states for different  $V$  values we can (within the mean approximation) test this adiabaticity assumption.

Looking again at the time profile depicted in Fig. 4 one may notice that the ramp used in the experiment leads indeed to a very slow increase of  $V(t)$  initially, however changes of  $V(t)$  become relatively rapid about and above  $V = 9E_r$ , i.e. in the region where the transition from




Figure 5: Atom density distribution (on-site filling factor) after a mean field evolution starting from the mean field ground state at  $V = 9E_r$  at the final value of  $V = 22E_r$  for different  $V(t)$  time dependence. (a) corresponds to the exponential time scale of 20 ms, (b) to 40 ms, (c) to 80 ms. The longer the time scale, the slower the change of  $V(t)$  in the considered range as can be seen in Fig. 4. Thick dashed lines in (a) to (c) present twice the variance of the on-site occupation showing that the insulator regions in (b) and (c) are much larger than in (a). Panel (d) repeats, after Fig. 3, the mean field ground state at  $V = 22E_r$  for comparison.

SF to MI is supposed to take place. Taking as the initial state the mean field ground state at  $V = 9E_r$ , our simulations show that to assure adiabaticity a small change of  $V$  on the superfluid side (say from  $V = 9E_r$  to  $V = 9.1E_r$ ) requires about 20 ms (one needs 40 ms for a loop from  $V = 9E_r$  to  $V = 9.1E_r$  and back to keep the overlap on the initial state of the order of 99 percent). That strongly indicates that a much longer time is needed to traverse adiabatically the whole interesting region from  $V = 9E_r$  to  $V = 22E_r$ . And that change is realized in about 20 ms in the experiment.

To test the adiabatic issue further we shall concentrate in the following on the regime above  $V = 9E_r$  containing the quantum phase transition. Starting again from the Gutzwiller mean field ground state at  $V = 9E_r$  we simulate the time evolution up to  $V = 22E_r$  (with experimental time profile). We may compare the dynamically obtained wavefunction plotted in Fig. 5(a) with the mean field ground state at  $V = 22E_r$  (bottom right panel (d) in the figure). While the ground state has an insulator character almost everywhere in the trap with  $s_F = 0.01$ , the dynamically evolved wavefunction, by comparison, seems to reflect an excited wavepacket and

it has rather small regions where the occupation of sites is close to integer with vanishing number variance. The corresponding  $s_F = 0.12$  confirms the presence of a relatively large superfluid region.

To show that the effect is really due to the too fast increase of the lattice depth we have modified the experimental time profile slightly, by changing the exponential constant from 20 ms to 40 ms (or 80 ms). That makes the initial rise of the laser intensity (and the lattice depth) more uniform in time – compare Fig. 4. Observe that while the full duration of the first stage remains the same, the interval of time spent on the increase of the lattice depth from  $V = 9E_r$  to  $22E_r$  increases from below 20 ms (experimental profile), to about 30 ms (for  $\tau = 40$  ms) or about 37 ms (for  $\tau = 80$  ms). Starting again from the mean field ground state at  $V = 9E_r$  we obtain the atom density profiles shown in Fig. 5(b) and Fig. 5(c), respectively. Observe that the regions of insulator behavior for both cases are much larger than observed previously. The corresponding superfluid factors are  $s_F = 0.062$  for  $\tau = 40$  ms and  $s_F = 0.048$  for  $\tau = 80$  ms. While the final distributions still show signs of significant excitations, the insulator character becomes dominant for (c)

and also for (b) case.

Keeping the overall duration of the laser intensity turn-on at 80 ms and enlarging the final stage comes at the price that the initial rise from  $V = 0E_r$  to  $V = 9E_r$ , very smooth in the experiment [4] for  $\tau = 20$  ms, becomes sharper for larger  $\tau$  (compare Fig. 4). Thus larger  $\tau$  may lead to some excitation at the initial creation of the lattice, not apparent in our simulations since we start from the ground state at  $V = 9E_r$ . Trying to keep the total duration of the experiment as short as possible (to avoid, for example, the decoherence) one can still imagine a slightly more sophisticated pulse rise with say  $\tau = 20$  ms initially up to say  $V = 9E_r$  and further increase with a larger time constant, say 40 ms. While the duration of the experiment increases by 10% only, the degree of the excitation of the final wavepacket becomes much smaller and the insulator character much more pronounced.

In the experiment [4] the atomic density distribution is not measured directly. The presence of the Mott insulator layer has been detected by observing the resonance in the excitation spectrum around the interaction energy  $U$ . Clearly the size of the corresponding peak is related to the number of atoms in the insulator layers. Our results indicate that an appropriate suggested change in the time profile of  $V(t)$  should increase the size of insulator regions and thus enhance the resonant peak in the excitation spectrum.

Further evidence presented below also points out that significant nonadiabatic excitation in the experiment is created due to the relatively fast changes of the lattice depth. In the experiment, after reaching  $V = 22E_r$  the lattice height is kept constant for 20 ms and then rapidly decreased back to  $V = 9E_r$ . It is shown that the time needed to restore the coherence is of the order of 4 ms. These data are obtained measuring the widths of the observed interference patterns (momenta distribution). If, indeed no excitation occurred during the lattice height increase stage then at  $V = 22E_r$  the system would be in the corresponding ground state and the final stage could be reproduced starting from this state. Our simulations fail to reproduce this fact. Depending on the slope of the final decrease the height of the pattern changes but not the halfwidth of the central peak (evaluated by a Lorentzian fit as in [4]).

Maybe our mean field simulations are not sufficient to reproduce the experimental results? It is not so, as shown in Fig. 6. We make a simulation, starting from the static solution in the SF regime (taken for the convenience at  $V = 9E_r$  again) increasing exponentially the lattice height as in the experiment [4], the subsequent delay of 20 ms at  $V = 22E_r$  and a linear ramp-down with various slopes. Note that the shape of the curve as well as the time scale of restoring the coherence is in quite good agreement with the experiment. While the experimental data could be fit with a double exponential decay with two time scales, our mean field data are reasonably reproduced with a single exponential decay with time scale  $\tau = 1.45$  ms. This nicely corresponds with the shorter




Figure 6: Half-width of the central interference peak for different ramp down times  $t_r$  obtained by a Lorentzian fit of the integrated momentum distribution (compare (7)). Filled circles are connected by a line to guide the eye. Dashed line is a single exponential decay with a time constant  $\tau = 1.45$  ms.

time scale of the experiment (0.94 ms). The obtained time scale is also of the order of a typical single tunneling time ( $1=J$  in appropriate units) to the nearby site. On the other hand, if we associate the second time scale with long range correlation between sites it becomes clear why this time scale does not manifest itself in our mean field simulations (the Gutzwiller wavefunction neglects entanglement between sites).

The observed quite good agreement of the obtained widths of the momentum distribution with the experiment [4] seems to be quite a spectacular success of the dynamical mean field simulation bearing in mind its simplicity. The fact that the mean field approach works so well may be, in our opinion, attributed to the fact that the dynamics takes place in the regime where superfluid fraction remains significant. Then the mean atomic field  $\phi_i$  does not vanish allowing for semiclassical (mean-field) description. Our results suggest that the system has quite a long memory and remembers that it was originally a SF. This fits nicely with the excited wavepacket-like character of dynamically obtained wavefunction clearly visible in Fig. 5(a).

#### IV. CONCLUSIONS

To summarize, it has been shown that the mean field Gutzwiller approximation allows one to simulate a dynamics of inhomogeneous Bose-Hubbard model taking into account realistic experimental conditions. The accuracy of the approximation cannot be controlled which is the major drawback of the present approach (a comparison with exact dynamics for small systems will lead us nowhere since then the mean field approach is known to fail). On the other hand a comparison with the available data seems quite encouraging. Accepting mean field pre-

dictions we may confirm that indeed the transition from superfluid to Mott insulator takes place in the experiment [4]. On the other hand the claim that the first stage of the experiment is performed adiabatically assuring that the system remains in its many body ground state (and thus a genuine textbook quantum phase transition [3] is realized) seems questionable.

We suggest that optimization of the lattice depth time dependence (i.e. laser intensity profile) may help to enlarge the insulator regions making the transition more adiabatic. That may be detected by measuring the size

of the peak in the excitation spectrum of the system.

Lastly, let us mention, that a very recent preprint [25] reports a study of exact dynamics of the model using the method of [20, 21]. However, the results consider at most 49 atoms in 40 sites of 1D lattice.

Participation of D. Delande at the early stage of this work is appreciated as well as discussions with B. Damski, Lewenstein and K. Sacha. This work was supported by Polish Committee for Scientific Research Grant Quantum Information and Quantum Engineering, PBZ-MIN-008/P03/2003.

- 
- [1] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, & P. Zoller, *Phys. Rev. Lett.* 81, 3108 (1998).
  - [2] M. P. A. Fisher, P. B. Weichman, G. G. Grinstein, & D. S. Fisher, *Phys. Rev. B* 40, 546 (1989).
  - [3] S. Sachdev *Quantum Phase Transitions* (Cambridge Univ. Press, Cambridge 2001).
  - [4] M. Greiner, O. Mandel, T. Esslinger, T. W. Hansch, & I. Bloch, *Nature* 415, 39 (2002).
  - [5] D. van Oosten, P. van der Straten, and H. T. C. Stoof, *Phys. Rev. A* 63, 053601 (2001).
  - [6] D. B. M. Dickerscheid, D. van Oosten, P. J. H. Denteneer, and H. T. C. Stoof, *Phys. Rev. A* 68, 043623 (2003).
  - [7] C. Menotti, A. Serenzi, and A. Trombettoni, *New J. Phys.* 5, 112 (2003).
  - [8] D. Jaksch, V. Venturi, J. I. Cirac, C. J. Williams, & P. Zoller, *Phys. Rev. Lett.* 89, 040402 (2002).
  - [9] B. Damski et al. *Phys. Rev. Lett.* 90, 110401 (2003).
  - [10] B. Damski, J. Zakrzewski, L. Santos, P. Zoller, & M. Lewenstein, *Phys. Rev. Lett.* 91, 080403 (2003).
  - [11] A. Albus, F. Illuminati, and J. Eisert, *Phys. Rev. A* 68, 023606 (2003).
  - [12] H. P. Buchler and G. Blatter, *Phys. Rev. Lett.* 91, 130404 (2003).
  - [13] Fehrmann, H., Baranov, M. A., Damski, B., Lewenstein, M., & Santos, L. *cond-mat/0307635* (2003).
  - [14] Kashumikov V. A., Prokof'ev N. V., & Svistunov B. V., *Phys. Rev. A* 66, 031601 (2002).
  - [15] G. G. Batrouni et al, *Phys. Rev. Lett.* 89, 117203 (2002).
  - [16] W. Zwerger, *J. Opt. B: Quantum Semiclass. Opt.* 5, S9 (2003).
  - [17] S. Wessel, F. Alet, M. Troyer, and G. G. Batrouni, *cond-mat/0404552*.
  - [18] D. C. Roberts and K. Burnett, *Phys. Rev. Lett.* 90, 150401 (2003).
  - [19] L. Amico and V. Penna, *Phys. Rev. Lett.* 80, 2189 (1998).
  - [20] G. Vidal, *Phys. Rev. Lett.* 91, 147902 (2003).
  - [21] A. J. Daley, C. Kollath, U. Schollwöck, and G. Vidal, *J. Stat. Mech.: Theor. Exp.* P04005 (2004).
  - [22] W. H. P. Reiss, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in Fortran* Cambridge University Press, Cambridge 1996.
  - [23] Bogdan Damski as very kind to supply the appropriate numerical code.
  - [24] C. Orzel, A. K. Tuchman, M. L. Fenselau, M. Yasuda, and M. A. Kasevich, *Science* 291, 2386 (2001).
  - [25] S. R. Clark and D. Jaksch, *cond-mat/0405580*.