

Bose-Einstein condensation of magnons in TiCuCl_3

J. Sirker*, A. Weiße and O.P. Sushkov

School of Physics, The University of New South Wales, Sydney 2052, Australia

Abstract

A quantitative study of the field-induced magnetic ordering in TiCuCl_3 in terms of a Bose-Einstein condensation (BEC) of magnons is presented. It is shown that the hitherto proposed simple BEC scenario is in quantitative and qualitative disagreement with experiment. It is further shown that even very small Dzyaloshinsky-Moriya interactions or a staggered g tensor component of a certain type can change the BEC picture qualitatively. Such terms lead to a nonzero condensate density for all temperatures and a gapped quasiparticle spectrum. Including this type of interaction allows us to obtain good agreement with experimental data.

Key words: Spin-gap systems, Bose-Einstein condensation, spin-orbit coupling

PACS: 75.10.Jm, 03.75.Hh, 71.70.Ej

TiCuCl_3 is a spin-dimer system with an excitation gap $\Delta \approx 0.7$ meV in zero magnetic field. The dimers in this compound are formed by the $S = 1/2$ spins of the Cu^{2+} ions and superexchange interactions are mediated by the Cl^- ions. The lowest excited state is a triplet of massive bosons (magnons). The density n of the boson mode is directly related to the magnetization per site $m = g\mu_B n$. The magnon dispersion has been measured by inelastic neutron scattering [1] revealing a considerable dispersion in all spatial directions. TiCuCl_3 therefore has to be considered as a three-dimensional interacting dimer system.

A magnetic field H causes a Zeeman splitting of the triplet with the lowest mode crossing the singlet state at a critical field $H_c = \Delta/g\mu_B \sim 5.6$ T. The ground state for $H > H_c$ becomes a BEC of this low-energy boson. The condensation of magnons corresponds to a phase transition into a state with long-range antiferromagnetic order perpendicular to the applied field. In Ref. [2] it has been shown that this transition and the overall shape of the magnetization curves as a function of temperature are in rough qualitative agreement with the BEC picture. However, we find that the simple magnon dispersion $\Delta + \mathbf{k}^2/2m$ used in [2] is only justified for temperatures $T \ll 1$ K (see inset of Fig. 1) well below the experimental temperature range. Therefore

the universal power-law dependence between critical density and temperature $n_c \sim T_c^\alpha$ with $\alpha = 3/2$ expected within the BEC picture will only be observable at extremely low T thus offering a simple explanation for the deviations observed in [2]. Following Ref. [3] we have therefore first calculated the real triplet dispersion using a bond operator formalism [4]. The starting point is the strong coupling ground state $|s\rangle$ where each dimer at site i is in singlet configuration $|i, s\rangle$. One then expands the Hamiltonian in terms of operators $t_{i\alpha}^\dagger$ which create local triplet excitations with spiral index α . In first order a Hamiltonian bilinear in the operators $t^{(\dagger)}$ is derived. Only the most important exchange pathways between the dimers are considered so that the limited exchange parameter set can be determined by a fit to the measured dispersion [1]. In agreement with [3] we find that terms quartic in the triplet operators derived in next order yield only small corrections. The most important renormalization of the triplet dispersion is due to the hard-core constraint for the triplets, which can be taken into account by introducing an infinite on-site potential

$$\mathcal{H}_U = U \sum_{i,\alpha,\beta} t_{i\alpha}^\dagger t_{i\beta}^\dagger t_{i\alpha} t_{i\beta}, \quad U \rightarrow \infty. \quad (1)$$

The corresponding two-particle scattering vertex $v(\mathbf{k}, \omega)$ can be calculated exactly in the dilute limit by a summation of ladder diagrams [5]. Doing so we find a considerable

* Corresponding author: sirker@phys.unsw.edu.au

renormalization of the exchange parameters compared to the mean-field treatment in [3]. For the scattering vertex at the band minimum q_0 we find $v_0 = v(q_0, 0) \approx 10$ meV [4]. The density of states $\rho(\Omega)$ calculated with the renormalized dispersion Ω_k is shown in Fig. 1.




Fig. 1. Density of states $\rho(\Omega)$. The inset shows $\rho(\Omega)$ near the gap and a fit (dashed line) $\rho(\Omega) \sim \sqrt{\Omega - \Delta}$ yielding an effective mass $m_{\text{eff}} \sim 0.2$ meV $^{-1}$.

For magnetic fields $H \gtrsim H_c$ and temperatures $T < \Delta$ it is sufficient to take only the lowest triplet mode into account. The Hamiltonian is then given by

$$\mathcal{H} = \sum_k (\epsilon_k - \mu_0) t_k^\dagger t_k + \frac{v_0}{2} \sum_{k, k', q} t_{k+q}^\dagger t_{k'-q}^\dagger t_k t_{k'} \quad (2)$$

where $\epsilon_k \equiv \Omega_k - \Delta$ and $\mu_0 = g\mu_B(H - H_c)$. In TiCuCl₃ typical magnon densities in magnetic fields $H \sim 6-7$ T are of the order $n \sim 10^{-3}$ [2]. This allows us to calculate the thermodynamic properties systematically in an expansion in n (gas approximation). Practically it turns out to be sufficient to treat the interaction in (2) in the one-loop approximation. After diagonalizing the resulting bilinear Hamiltonian we can directly calculate the Boson density (magnetization) as a function of temperature [4] using the density of states from Fig. 1. The result is shown in Fig. 2a. Compared to




Fig. 2. Experimental $M(T)$ (symbols) for $H = 5.9, 6.0, \dots, 7$ T taken from Ref. [2]. (a) The theoretical $M(T)$ (solid lines) with $\Delta = 0.67$ meV and $v_0 = 9.8$ meV. (b) Interaction (3) included with $\gamma = 10^{-3}$ meV, $\Delta = 0.72$ meV and $v_0 = 27$ meV.

experiment the theoretically calculated magnetizations are too large and the T_c 's too high although $n_c \sim T_c^{2.1}$ yields a good fit with the exponent being close to the observed one.

Finally, we want to discuss reasons for this shortcoming. First, apart from a small temperature interval $|T - T_c| \lesssim$

$n^{1/3} a T_c$ with $n^{1/3} a \lesssim 0.1$ the one-loop approximation is reliable and higher order corrections are small. Therefore there must be some physics we missed so far. An interaction which we ignored and which is known to be important in TiCuCl₃ is spin-phonon coupling [6]. Its main effect will be a reduction of the bare magnon bandwidth and therefore an effective increase of v_0 . With $v_0 \sim 25$ meV we can indeed obtain good agreement with experimental data for $T \geq T_c$, however, the calculated magnetizations at low temperatures are still 50% too large [4]. In recent ESR measurements [7] a direct singlet-triplet transition has been observed which suggests that a perturbation of the form

$$\mathcal{H}' = i\gamma (t_{q_0} - t_{q_0}^\dagger) \quad (3)$$

might be present mixing the singlet and the triplet state. Here γ is a small parameter. Such a perturbation is equivalent to a staggered magnetic field perpendicular to the applied field. It can result from a staggered g tensor or Dzyaloshinsky-Moriya interactions which are both induced by spin-orbit coupling. With this term included there is no longer a sharp phase transition from a condensed into a non-condensed phase but instead the number of condensed magnons decreases exponentially with temperature. Furthermore there is no longer a gapless mode usually characteristic for the condensed phase [4]. In TiCuCl₃ the interaction (3) requires breaking of the inversion symmetry within a dimer. This might be due to small static distortions as discussed in Ref. [4]. We therefore expect γ to be quite small. It turns out that it is indeed possible to obtain excellent agreement with experiment with $\gamma = 10^{-3}$ meV as shown in Fig. 2b.

References

- [1] N. Cavadini, et al., Phys. Rev. B 63 (2001) 172414.
- [2] T. Nikuni, et al., Phys. Rev. Lett. 84 (2000) 5868.
- [3] M. Matsumoto, et al., Phys. Rev. Lett. 89 (2002) 077203; M. Matsumoto, et al., Phys. Rev. B 69 (2004) 054423.
- [4] J. Sirker, et al., cond-mat/0403311.
- [5] V. N. Kotov, et al., Phys. Rev. Lett. 80 (1998) 5790.
- [6] K.-Y. Choi, et al., Phys. Rev. B 68 (2003) 174412; E. Y. Sherman, et al., Phys. Rev. Lett. 91 (2003) 057201.
- [7] V. N. Glazkov, et al., Phys. Rev. B 69 (2004) 184410.